

**Dietary Intake, Nutritional Status of Lactating  
Women and their 6-23-Months-Old Children in  
*Genta Afeshum* District, Rural Ethiopia;  
Adaptation and Validation of Calculator for  
Inadequate Micronutrient Intake (CIMI)**

**Dissertation submitted in the fulfillment of the requirements for the  
Doctoral Degree of Natural Sciences (Dr. rer. nat. / Ph.D)**

**Faculty of Natural Sciences**

**University of Hohenheim**

**Institute of Nutritional Science**

**Submitted by**

**Beruk Berhanu Desalegn**

**Born in Awassa, Ethiopia**

**Stuttgart-Hohenheim 2020**

Dean:	Prof. Dr. Uwe Beifuß
1st reviewer:	Prof. Dr. Med. Hans Konrad Biesalski
2nd reviewer:	Prof. Dr. Jan Frank
3 <sup>rd</sup> examiner:	Prof. Dr. Tegene Negese

Submitted on: October 22, 2019

Oral examination on: September 11, 2020

This work was accepted by the Faculty of Natural Sciences at the University of Hohenheim on **December 27, 2019** as “Dissertation for Obtaining the Doctoral Degree of Natural Sciences”.

## General Information on Publication and Participation in the Scientific Workshops

Part of this dissertation are published and/ or submitted in internationally peer-reviewed journals, and/or presented in the international scientific workshop and/or accepted to be presented in the future.

The papers published are the following:

- Desalegn, B. B., Lambert, C., Riedel, S., Negese, T., and Biesalski, H. K. (2018). Ethiopian Orthodox Fasting and Lactating Mothers : Longitudinal Study on Dietary Pattern and Nutritional Status in Rural Tigray , Ethiopia. *Int. J. Environ. Res. Public Health*, 15(8),1767. <https://doi.org/10.3390/ijerph15081767>
- Desalegn, B., Lambert, C., Riedel, S., Negese, T., and Biesalski, H. (2019). Feeding Practices and Undernutrition in 6–23-Month-Old Children of Orthodox Christian Mothers in Rural Tigray, Ethiopia: Longitudinal Study. *Nutrients*, 11(1), 138. <https://doi.org/10.3390/nu11010138>

The paper submitted to journal is:

- Desalegn, B., Lambert, C., Gola, U., Riedel, S., Negese, T., and Biesalski, H.K. Calculator for Inadequate Micronutrient Intake (CIMI): Validation of the software in 12-23-months-old children and lactating women of rural Ethiopia

The papers presented or accepted for presentation in international scientific conferences are:

- Desalegn, B.B., Lambert, C., Riedel, S., Negesse, T., Biesalski, H. K. (2018). Ethiopian Orthodox Fasting and Lactating Mothers: Longitudinal Study on Dietary Pattern and Nutritional Status in Rural Tigray, Ethiopia. In *Global food security and food safety: The role of universities* (Vol. 30, p. 2018). Ghent, Belgium: Ghent University.
- Riedel, S., Lambert, C., Desalegn, B.B., Borko, T.B., Biesalski, H. K. (2019). Development and Validation of Calculator for Inadequate Micronutrient Intake (CIMI) Program for Ethiopian Population. In 4<sup>th</sup> International Hidden Hunger Congress: Hohenheim-Stuttgart, Germany: University of Hohenheim.
- Desalegn, B., Lambert, C., Widmer, C., Gola, U., Riedel, S., Negese, T., and Biesalski, H.K. Dietary Nutrient Intake of 12-23-Months-Old Children and Lactating Women in Rural Ethiopia; Development and Validation of Calculator for Inadequate Micronutrient Intake (CIMI) Program for Ethiopian Population. (This was accepted for poster presentation on the Nutrition 2019: Where the Best in Science and Health Meet: Baltimore: Maryland, USA: American Society for Nutrition, and the abstract was published on *Current Developments in Nutrition*, 3(S1). DOI: [10.1093/cdn/nzz036.P13-008-19](https://doi.org/10.1093/cdn/nzz036.P13-008-19)).

## List of Abbreviations

AOR	Adjusted Odd Ratio
ANC	Antenatal Care
AUC	Area Under Curve
BMI	Body Mass Index
BMZ	Bundesministerium Für Wirtschaftliche Zusammenarbeit (German Federal Ministry for Economic Development Cooperation)
COR	Crude Odd Ratio
CIMI	Calculator for Inadequate Micronutrient Intake
CDI	Centre for Development Innovation
CSA	Central Statistical Agency, Ethiopia
DAAD	German Academic Exchange Service
DALYs	Disability-adjusted life-years
DDS	Diet Diversity Score
DHS	Demographic and Health Survey
DRMFSS	Disaster Risk Management and Food Security Sector
EDHS	Ethiopian Demographic Health Survey
EPHI	Ethiopian Public Health Institute, formerly EHNRI
ENA	Essential Nutrition Action
ENGINE	Empowering the New Generation to Improve Nutrition and Economic Opportunities
EOTC	Ethiopian Orthodox Tewahedo Church
ERSS	Ethiopian Rural Socio-economic Survey
FAO	Food and Agriculture Organization of the United Nations
FANTA	Food and Nutrition Technical Assistance
FFQs	Food Frequency Questionnaires
GDP	Gross Domestic Product
GIZ	Deutsche Gesellschaft für Internationale Zusammenarbeit
HDL	High Density Lipoprotein
HFIAP	Household Food Insecurity Access Prevalence
HFIAS	Household Food Insecurity Access Scale
IMF	International Monetary Fund
IYCF	Infant and Young Child Feeding
LAZ	Length-for-Age
LDL	Low Density Lipoprotein
MAD	Minimum Acceptable Diet
MDD	Minimum Diet Diversity
MDD-W	Minimum Diet Diversity for Women at Reproductive Age
MMF	Minimum Meal Frequency
MOFED	Ministry of Finance and Economic Development, Ethiopia
NBS	Nutrition Baseline Survey
NNP	National Nutrition Program
NNS	National Nutrition Strategy
NS	NutriSurvey
NSA	Nutrition Sensitive Agriculture
PNC	Postnatal Care

PCA	Principal Component Analysis
RNI	Recommended Nutrient Intake
SDGs	Sustainable Development Goals
SPSS	Statistical Package for Social Science
UOH	University of Hohenheim
UNICEF	United Nations Children's Fund
UNU	United Nations University
USAID	United States of America Aids for International Development
USDA	United States Department of Agriculture
VIF	Variance Inflation Factor
WASH	Water Supply, Sanitation and Hygiene
WAZ	Weight-for-Age
WHO	World Health Organization
WLZ	Weight-for-Length

## **Acknowledgments**

"... All things were made by him; and without him was not anything made that was made..." John 1:3. No words for me to thank the blessing of God and his mother St. Virgin Marry since I come to this world.

Without the support of many people and institutions, this PhD work might not be real and successfully accomplished. My supervisor, Professor Dr. Biesalski, thank you very much for choosing me to be your student and gave me an opportunity to do my research in an area, where I did not dream, but extremely important for me to serve the community where I was born and grown up. Furthermore, your guidance, support and the comments you were forwarding from the design of the proposal until the end of the write-up of this dissertation were exceptionally important. Professor Tegene, my local supervisor, your mentorship, shaping me to improve the scientific writing skill and your valuable comments and corrections you are providing me for the manuscripts and this dissertation are immense. You have taught me a lot how to help people in the situation where I was stressed in getting local supervisor from Ethiopia. Dr. Christine, my immediate mentor for this PhD study is thanked for your close mentorship, guidance and technical support in the CIMI project activities and for the whole this PhD work, especially on statistical analysis, manuscript preparation and correction of the drafted manuscripts was exceptional, in addition to your patience and immediate feedbacks whenever I need your support in spite of your very busy schedule. Dr. Simon, project manager of Micro-gap Ethiopia, who was very friendly and supportive for the whole CIMI project period, and for the successful and timely delivery of the CIMI software adapted for Ethiopia population in the project schedule. Your frequent travel to Ethiopia, moreover to the hard to reach research site and home to home interview while we were identifying the local standard units for CIMI software in Ethiopia is unforgettable. Thanks to Dr. Jenny, Dr. Nicole and Dr. Heinrich of the Food Security Center and Mrs. Kasper, at the University of Hohenheim for your administrative support during my study period.

My appreciation goes to my friends (Dr. Tibebelesassie, Birhanu, Dr. Shimelis and Habtamu) in the University of Hohenheim and staffs (Dr. Kebede Abegaz and Dr. Samson Gebremedhin) and friends in the Hawassa University (Alemneh, Anbisa, Dr. Temesgen, Dr. Alemayehu, Dr. Yonas, Dr. Yonas, Biruk Ayalew, Dereje, Dagem, Wondessa and Yetnebersh) for your kind support you

provided me during the course of this PhD study. Thomas, your very special friendship and direction in both of my academic and non-academic carriers are gigantic.

I would like also to appreciate the German Federal Ministry for Economic Cooperation and Development (BMZ) for financing me this PhD scholarship through the University of Hohenheim, Germany. I appreciate the GIZ Nutrition Sensitive Agriculture (NSA) project in Ethiopia for providing me office in Mekele during my stay in field work. I would like also to thank the mothers and children who are participated in this study. The GIZ NSA experts, *Genta Afeshum woreda* Nutrition coordinator, data collectors, community facilitators, data encoders involved in this study also highly appreciated.

Abaye and Zewdeya, your question ‘How is your work?’ is always alarming me when I met you, beside your love and support you are providing me. Gashe Admasu and Dr. Birhanu, without you and your families support, it might not be easy to me to be at this success. Assie, Emie, Mesay, Engocha, Fraa, Biruk and Dagi, your care for my wife and my daughter was grateful. My special thanks go to Addisalem, my wife for your love, support, patience and taking care of our daughters: Aksale and Bethel, and handling a lot of issues during this doctoral study period.

Beruk Berhanu Desalegn

# TABLE OF CONTENTS

<b>General Information on Publication and Participation in the Scientific Workshops .....</b>	<b>III</b>
<b>List of Abbreviations .....</b>	<b>IV</b>
<b>Acknowledgments .....</b>	<b>VI</b>
<b>1. General Introduction .....</b>	<b>1</b>
1.1. Background .....	2
1.2. Outline of the Thesis .....	13
1.3. References .....	16
<b>2. Ethiopian Orthodox Fasting and Lactating Mothers: Longitudinal Study on Dietary Pattern and Nutritional Status in Rural Tigray, Ethiopia .....</b>	<b>29</b>
Abstract .....	31
2.1. Introduction .....	32
2.2. Materials and Methods .....	34
2.2.1. Study Area, Design, Participants, and Sampling .....	34
2.2.2. Data Collection .....	35
2.2.3. Minimum Women Diet Diversity Score .....	36
2.2.4. Household Food Insecurity Information .....	37
2.2.5. Wealth Index .....	37
2.2.6. Anthropometry .....	38
2.2.7. Data Management and Analysis .....	38
2.3. Results .....	39
2.4. Discussion .....	50
2.4.1. Nutritional Status of Lactating Mothers .....	50
2.4.2. Dietary Patterns of Lactating Mothers .....	54
2.5. Conclusions .....	58
2.6. References .....	59
<b>3. Feeding Practices and Undernutrition in 6–23-Month-Old Children of Orthodox Christian Mothers in Rural Tigray, Ethiopia: Longitudinal Study .....</b>	<b>71</b>
Abstract .....	73
3.1. Introduction .....	74
3.2. Materials and Methods .....	76
3.2.1. Study Area, Design, Participants and Sampling .....	76
3.2.2. Data Collection .....	77



3.2.3. Data Management and Analysis .....	78
3.3. Results .....	80
3.3.1. Socio-Demographic Characteristics .....	80
3.3.2. Nutritional Status and Consumption Pattern .....	81
3.3.3. Factors Associated with Child Undernutrition .....	83
3.3.4. Trend of Food Consumption Pattern and Nutritional Status of 6–23-Month-Old Children .....	84
3.4. Discussion .....	86
3.4.1. Food Consumption Pattern of 6–23-Month-Old Children .....	87
3.4.2. Nutritional Status of 6–23-month-Old Children.....	90
3.5. Conclusions .....	93
3.6. References .....	95
<b>4. Calculator for Inadequate Micronutrient Intake (CIMI): Validation of the Software in 12-23-Months-Old Children and Lactating Women of Rural Ethiopia .....</b>	<b>106</b>
Abstract .....	107
4.1. Introduction .....	108
4.2. Materials and Methods .....	110
4.2.1. Study Area, Design, Participants, Sampling and Ethical Clearance .....	110
4.2.2. Data Collection .....	111
4.2.3. Development of Calculator for Inadequate Micronutrient Intake (CIMI) for Ethiopia .....	111
4.2.4. Data Management and Analyses .....	112
4.3. Results .....	114
4.4. Discussion .....	126
4.4.1. Validation of Calculator for Inadequate Micronutrient Intake (CIMI) .....	127
4.4.2. Dietary Nutrient Intakes of 12-23-Months-Old Children and Lactating Women in Rural Ethiopia.....	131
4.5. Conclusions .....	135
4.6. References .....	136
<b>5. General discussion .....</b>	<b>144</b>
5.1. Undernutrition in Lactating Women and their 6-23-Months-Old Children.....	144
5.1.1. Prevalence of undernutrition in lactating women and their 6-23-months-old children .....	144

5.1.2. Factors associated with undernutrition of lactating women and their 6-23-months-old children .....	146
5.2. Feeding Practices and Nutrient Intake of Lactating Women and their 6-23-Months-Old Children.....	150
5.3. CIMI: A Promising Dietary Assessment Application for Rural Ethiopian Population ....	154
5.4. Conclusion.....	159
5.5. References .....	162
<b>6. Summary.....</b>	<b>172</b>
<b>Zusammenfassung.....</b>	<b>175</b>
<b>List of All References.....</b>	<b>179</b>
<b>Affidavit .....</b>	<b>214</b>
<b>Curriculum Vitae .....</b>	<b>215</b>

# **Chapter 1**

## **General Introduction**

# **1. General Introduction**

## **1.1. Background**

Globally, a lot of efforts have been undertaken in reducing malnutrition, even though the achievements are far less than the global goals targeted [1]. As a result, it remains unacceptably high and a serious public health problem in the globe. This categorizes malnutrition as the leading cause for more illness than others [2]. Worldwide, the estimated prevalence of undernourished people increased from 784 million (10.6%) in 2015 to 821 million (10.9%) in 2017 [3]. However, children and women at the reproductive age are more affected by malnutrition than men. The number of stunted, wasted and overweight children under five years of age were about 151 million, 50 million and over 38 million in 2017, respectively, while about 170 million underweight and 613.2 million anemic women at reproductive age, and more than 2 billion adult overweight (1.02 billion women vs. 984.6 million men) existed in 2016 [2]. Beside these, deficiency of micronutrients affects more than 2 billion people living in this planet [4]. It is also the principal cause for the 3.5 million deaths, 11% of the total global disability-adjusted life-years (DALYs), 35% of the disease burden in young children and 20% of the maternal mortality [5]. However, the highest prevalence of undernutrition (20.4%) is registered in Africa, but the least improvement (0.8%) is shown between 2005 and 2017 [3].

Despite the high and rapid economic development achieved in Ethiopia since 2005/06, the prevalence of food and nutrition insecurity remains to be high [6–8]. The number of food insecure Ethiopian people were about 7.8 million [9]. Since 2008, Ethiopian government has been exerting a lot of efforts to accelerate the reduction of undernutrition by developing and launching National Nutrition Strategy (NNS), and National Nutrition Programs (NNP) I and II, and fully integrating nutrition issues in the health sector program of the country [10–12]. However, the

achievement is still much lower than the ambitious ‘Seqota declaration’ goal set by 2030, which is elimination of all forms of malnutrition among the children below the age of two years [13]. The recent Ethiopian demographic and health survey report showed that the prevalence of stunted, wasted, underweight and anemic children under the age of 5 years were 38%, 10%, 24% and 57%, respectively. Likewise, 22% and 24% of women between the age of 15-49 years was thin ( $BMI < 18.5\text{kg/m}^2$ ) and anemic, respectively [6]. These burdens also co-existed with the ever increasing over-nutrition problem such as 1% overweight in children and 8% overweight or obese in women [6]. But, the distribution of the existing undernutrition problems vary by region, geographical locations, people and time [14–20]. Similar studies in three sub-Saharan African countries and Nigeria also confirmed these evidences [21,22,23]. According to UNICEF’s nutrition scale up approach for mothers and their children, inadequate dietary intake and disease are the immediate causes for the maternal and child undernutrition, which are affected by the underlying causes like household food insecurity (lack of availability, access and/or utilization of diverse diet), inadequate care and poor feeding practices for children, unhealthy household and surrounding environments, and inaccessible and often inadequate health care [24].

Food taboo is one of the socio-cultural practices in countries like Ethiopia, where more than 85 ethnic groups and different religious beliefs exist, and could affect the nutritional status of women and children [24]. Food taboos refer to those foods which are strictly forbidden for health, cultural, and religious reasons [25–31]. In different African countries including Ethiopia, food taboos are thought to have been established more often during pregnancy than other physiological stages of women, as a means of protecting the health of women and their babies. As a result, most studies in Ethiopia and other African countries focused on the exploration of food taboos during pregnancy and its association with their nutritional status and feeding

practices [25,26,28,31–38]. Accordingly, some studies in Ethiopia showed that 27-60% of pregnant women are subject to food taboo [25,26,36]. They also identified that linseed, honey, dairy products, meat, eggs, fruits and vegetables, coffee, tea, porridge, wheat bread, pimento, groundnut, salty diet, nug, sugarcane, pumpkin and coca drinks were among the food items not allowed for pregnant women [25,31,36]. The main reasons for avoidance of these food items were fear of plastering on fetal head, getting a big baby which is difficult for delivery, fear of abortion, evil eye and fetal abnormality. However, lactating mothers who required greater amount of energy and most nutrients even by pregnant mothers, are also the victim of food taboo although little is known in this regard [40–42]. However, if the practice of food taboos is not identified and intervened, then it could be passed from one generation to the next generation and may sustain the intergenerational cycle of malnutrition [43,44].

Fasting is a partial or complete abstention from all foods and water or refrain from some selected and or/ prohibited foods for some fixed period of time in the year [45,46]. Religious fasting is among the different types of dietary or food taboos, which may affect the dietary intake and nutritional status of children and women. Apart from the conventional food taboos, religious fasting has a momentary nature. In these fasting periods, abstaining from animal source foods and/or water is done in the fixed period of time in the year and is a religious obligation [45].

Ramadan is in the ninth month of Islamic lunar calendar and is when Muslims fast for 29 to 30 days. In this period, Muslims are expected to abstain from eating, drinking, smoking, sexual practices and other activities for an average of 12-hrs in a day from the dawn to dusk [47,48]. Beside this, the type and amount of food to be eaten during the night of Ramadan is also significantly altered than during the rest of the year [49]. Even if fasting during Ramadan is an obligation for Muslims; pregnant women, lactating women, children, the elderly, travelers and

some people with specific conditions like female at menstrual cycle or persons who cannot withstand for longer time without food or water due to some acute or chronic diseases are exempted from this practice [50,51]. However, usually breastfeeding and pregnant women and other exempted people prefer to fast with their family members to show their spiritual solidarity [50–53]. Accordingly, studies in Lebanon and Turk, pregnant women showed that the weight gain was lower in fasting compared to non-fasting groups during Ramadan [52,54]. Similarly, the % of recommended daily allowance of most nutrients fulfilled by the Turkish pregnant women who were fasting were lower compared to those pregnant women who did not [54]. The cohort study in Lebanon also showed that the mean birth weight of the women who fasted during their pregnant period were significantly lower than the average mean birth weight of the women who did not fast during their pregnancy period. Likewise, a study on Turkish lactating women confirmed that the nutrient quantity of the breast milk and weight and their child's weight were lower in Ramadan fasting period than after one month non-fasting period [53]. A study on patients with a cancer-related fecal stoma revealed that the weight loss after Ramadan fasting was about 9.5kg compared to the time right before Ramadan [50]. A systematic review and meta-analysis in West Asia, Africa, East Asia and North America/Europe confirmed that the weight loss was significantly higher, regardless of gender, but this loss was regained within a few weeks after Ramadan [48].

In Orthodox Christian religion, followers are expected to fast for at least 180 days in a year [46,55–58]. Fasting includes abstention from all animal source foods (meat, milk, butter and eggs), but restriction of fish in Ethiopian Orthodox church is not uniform [45]. Studies in Greek Orthodox Christian monks revealed that the consumption of dairy products, meat and eggs, and the dietary intake of calcium, and BMI were lower in fasting weeks compared to week following

fasting. However, the profile of individual health biomarkers like total cholesterol, LDL-cholesterol and HDL-cholesterol in serum were lower during fasting than non-fasting weeks [60]. Annual longitudinal study on 120 Greek Orthodox Christians (60 fasting vs. 60 non-fasting as control) also confirmed that a significant reduction in the total and LDL-cholesterol, and BMI was observed in an end line survey compared to control group [59]. Similar authors in the Greek Orthodox followers also evidenced that the dietary cholesterol, protein, energy and calcium intakes were also low in fasting groups compared to the control in the end-holidays than pre-fasting period [57]. Two review papers focused on the health related impact of Orthodox fasting also confirmed that BMI, lipid profile including cholesterol level of fasting individuals decreased, and these improved their health status [46,61]. Relatively, a recent systematic review on the effect of Orthodox fasting on human health confirmed that few researches were conducted in this specific topic. But the results of these studies indicated that the total energy and fat intakes are restricted, but carbohydrate and fiber intakes increased. Beside this, the overall effect of Orthodox fasting on reduction of total cholesterol and LDL-C levels were somewhat optimal. According to Kaufakis and colleagues, studies on the impact of Orthodox fasting are limited, but these studies are also conflicting with each other's, as a result a consensus has not been reached to draw conclusion. Therefore, further study on this regard is recommended [62]. In general, many of the researches focused on the effect of fasting on human health specifically chronic diseases. However, knowing the fasting effect on dietary pattern and nutritional status in countries with high prevalence of maternal and child undernutrition is very important.

Follower of Ethiopian Orthodox and Muslim religion make up to three-fourth of people in Ethiopia, who practice a strict fasting during their religious fasting periods [63]. But, their fasting practices including number of fasting days, type of fasting, number of fasting hours and type of



food items restricted are some of the differences between these two religions [64]. According to Ethiopian Orthodox church rule, there are seven official fasting periods and a total of these periods cover approximately 250 out of 365 days in a year. In these days, children less than seven years old, soldiers, severely ill or weak people, pregnant, and lactating mothers are permitted to eat both animal source foods and other foods including water, without abstention during the religious fasting periods or days [45,58,65]. However, little studies in Ethiopia highlight the religious fasting in relation with feeding practices and nutritional status in nutritionally vulnerable groups (women and children) who are the follower of the two dominant religion. Despite these nutritionally needy groups (lactating and pregnant mothers, and children) who are exempted from the religious fasting, yet the consumption of animal source foods during the fasting periods are very low for different reasons [43,66]. For example, studies in Amhara and Tigray regions explored that some mothers were not happy to prepare food for their children from animal sources due to fear of utensils contamination during cooking family food, and this reason increased the likelihood of feeding less diverse food for their children by 1.5 times compared with the economic related reasons [44,67,68]. Other studies in Ethiopia also showed that the demand of cattle meat during Orthodox fasting periods was observed to be low, resulting in closure of abattoirs or minimizing the service provided. A further study also reported that more than 85% of butcher houses were closed during Wednesday and Friday, which are Orthodox Christians fasting days of the week, in Addis Ababa, a capital city of Ethiopia [70,71].

The first 1000 days of life, between the conception of a woman until her child celebrates his/her two years' birthday, is a critical window of opportunity to build a foundation for optimal health and development, and success for the child across his/her life time. Proper nutrition and care during this window period improve the survival rate, his or her growth and learning ability, and

later contribute to societies long term health, stability and prosperity [72]. A mother's diet and her nutrient stores are the only source of nutrition for the developing baby in her womb. When a pregnant woman does not get the calories, key nutrients or essential proteins she needs to support her baby's development, her baby is placed at risk for developmental delays, birth defects and cognitive deficits [73]. Undernutrition in pregnant women increase the risk of maternal morbidity and mortality and also prone to poor pregnancy outcome, which include obstruction during labor, preterm or low birth child and postpartum hemorrhage [74]. Beside this, folate is critical to the early development of the brain and spinal cord of the fetus. When a woman lacks this before becoming pregnant and in the early weeks of her pregnancy, the development of the neural tube is affected, leading to birth defects of the brain and spinal cord (anencephaly and spinal bifida) that can cause death or lifelong disability [73]. Lactation is the second physiological stage, which cover three-fourth of the 1000 window days. Even except iron and folate, energy and most micronutrients need of lactation is greater compared to pregnancy [40,42,75–77]. During lactation, the energy, macro- and micronutrients in breast milk come from the mother's diet and/or her own body stores, which make the lactating women in high risk of malnutrition unless she consumes adequate nutrient and energy rich foods [51,53]. The quality of breast milk could be altered, as a result the nutritional status of the breastfed child could also change [51,78]. Furthermore, if the lactating mother is severely malnourished, then her lactation performance will decrease and contribute to child morbidity and mortality [79].

Exclusive breast feeding during the first six months after delivery and continued breast feeding up to one to two years in combination with introduction of appropriate, adequate and safe complementary feeding are among the recommended IYCF practices [80]. Appropriate complementary feeding can prevent about 6% of deaths of under-five aged children [5,81]. The

transition period from exclusive breastfeeding to two years is critical for optimal growth and development of a child [81]. The prevalence of “ever breastfeeding” in rural Ethiopia was about 97.8% versus 95.2% of recommended IYCF practices, which is a positive practice [ 82,83]. Likewise, many rural mothers continue breastfeeding until their child reaches the age of at least 24 months and in some cases even for a longer time. According to national study in Ethiopia, 76% of 2 years old children were continually breastfed [6]. Previous studies in Tigray, northern Ethiopia showed that 25% of the children were breastfed for more than 3 years [ 84]. Therefore, assessing the dietary adequacy of energy, macro- and micronutrients is very useful in the whole population groups specially children and women. For this, a number of both qualitative and quantitative dietary assessment techniques have been developed and implemented globally. However, most of the quantitative dietary nutrient intake assessment techniques like food frequency questionnaire (FFQs), food weight record and 24-h recall are time consuming, need well trained field workers during the surveys and an expert to calculate dietary intake using nutrition software and identify undernutrition. Therefore, the application of these methods has been less in rural poor settings in developing countries including Ethiopia. Alternatively, many assessment and intervention studies use a qualitative method known as diet diversity scores, which is a proxy indicator for assessing the dietary quality at household and individual levels [85]. Accordingly, national and different studies, elsewhere in Ethiopia showed that the consumption of foods from different food groups in children and women (pregnant, non-pregnant-non-lactating, and lactating women) were low [40,81–83,86–88]. However, this technique identifies the score of the consumption of diet from different standardized food groups regardless of the amount consumed, as this technique doesn’t allow the serving size ideally. Therefore, unless small amount of consumption is excluded in the count, then the validity of the

DDS will be in question. For this, many scholars suggested that consumption of at least 15 g from the group should be the minimum to be considered as consumed in women at reproductive age, otherwise the relationship between food group diversity and micronutrient adequacy is weaker [85,90–92]. As a solution, defining the food and ingredients to count and not to count during the construction of the questionnaire for MDD-W indicator should be done, and during the training of enumerators to avoid biased-decisions [85]. But, still the controversy remains on how quantification is practical in qualitative diet diversity assessment technique, unless the diet diversity data is extracted from other quantitative dietary intake data. Minimum acceptable diet, which is a combination of minimum diet diversity and minimum meal frequency is used to assess the quality and quantity of children's diet, in addition to other IYCF indicators [80]. Many studies at district and national levels in Ethiopia confirmed that the proportion of 6-23-months-old children who met the minimum number of meals from minimum diversified complementary food for their age and breastfeeding was very low (<10%) [6,43,93,94].

Human being needs, both nutrients which is called macronutrients (fat, protein and carbohydrate) and micronutrients which are needed in small amount, but are important for the metabolic process and function of the macronutrients are necessary [4,95]. Unlike macronutrients, micronutrients neither substitute each other nor are synthesized in our body. Therefore, supply of these nutrients through our diet is mandatory, although the amount required differ due to age, life style, hormonal activity or exercise and bioavailability related with food to be consumed [4]. If these micronutrients are inadequate, it leads to the invisible form of undernutrition called micronutrient deficiency or 'Hidden hunger'. If it persists for some longer period will increase the risk of disease, growth and cognitive development impairment in the early stage of childhood, and later maternal illness and decrease the life expectancy. Thus, early identification

of micronutrient deficiencies will be an important step to design and initiate appropriate prevention and intervention activities [96]. However, these are flouted, as the quantitative dietary data are not available most often [2]. As a result, the hidden hunger might not be addressed properly in Ethiopia and other African countries, where this occurrence remains high and persistent [2,6]. Therefore, an assessment tool which can estimate the quantitative dietary intake of an individual or a community, but easy to use, less costly and applicable method for developing countries including Ethiopia is urgently needed to achieve the national and international goals set for eradicating malnutrition. The Calculator for Inadequate Micronutrient Intake (CIMI) is a simple, easy-to-use, informative, web-based application of quantitative dietary assessment method. The first version of CIMI was developed in Indonesia using commonly consumed Indonesian foods for Indonesian population. Thus, CIMI estimates energy and nutrient intake correctly, and identifies nutrient inadequacy according to FAO/WHO recommended nutrient intake (RNI) regarding age, sex and physiological stage [96].

*Genta Afeshum* is one of the thirty-five district's in the Tigray regional state in Ethiopia. According to CSA (2007) projection, the district has a total population of 99,112, and almost all (99%) people are Orthodox Christianity follower. The district is known with cultivation of majorly cereals, followed by pulses, oil seeds, fruit trees and vegetables in decreasing order of importance. Of these, wheat, barley, legumes, maize, fenugreek, teff, sorghum and orange fleshed sweet potato are mentionable. However, Genta Afeshum is one of the hotspot districts for food insecurity in the region. It is also known with major disaster risks like drought, hail storms and livestock diseases; followed by human diseases, crop diseases and pests and flooding. Deforestation, water pollution and soil erosion are the major environmental problems; whereas, high dependency syndrome, poor economic condition, land shortage, severe shortage of drinking

water and poor saving are among the major vulnerability factors at household level [97]. But, to the best of our knowledge, there was no study which assessed the dietary pattern and nutritional status of women and children in Genta Afeshum district.

Therefore, the purpose of this study was assessing and comparing the nutritional status and dietary intake of lactating women and their 6-23months old children in fasting and non-fasting periods at Genta Afeshum District, rural Tigray in Northern Ethiopia, and to develop and validate the CIMI program for Ethiopian population. For this, the following specific objectives were set to address the three papers written as chapter two, three and four of this Dissertation,

1. Assessing and comparing the nutritional status of lactating mothers and their 6-23-months old children in Ethiopian Orthodox lent fasting and non-fasting periods in Genta Afeshum district, Rural Tigray, Ethiopia.
2. Assessing factors associated with undernutrition ( $BMI < 18.5 \text{ kg/m}^2$ ) in lactating mothers of Genta Afeshum district, Rural Tigray, Ethiopia.
3. Assessing factors associated with undernutrition (stunting, underweight and wasting) in 6-23-months old children of Genta Afeshum district, Rural Tigray, Ethiopia.
4. To compare dietary pattern of lactating mothers and their 6-23-months old children during fasting and non-fasting periods in Genta Afeshum district, Rural Tigray, Ethiopia.
5. To adapt CIMI program and compare with NutriSurvey for its ability to analyze dietary nutrients and energy intakes in Genta Afeshum district, Rural Tigray, Ethiopia.

## 1.2. Outline of the thesis

This Thesis/Dissertation has six chapters and the biography at the end. Chapter 2 of this Thesis/Dissertation is entitled “Ethiopian Orthodox Fasting and Lactating Mothers: Longitudinal Study on Dietary Pattern and Nutritional Status in Rural Tigray, Ethiopia “. The chapter focused on identifying underweight ( $BMI < 18.5\text{kg/m}^2$ ) and food consumption pattern of fasting and non-fasting mothers in Ethiopian Orthodox lent fasting and non-fasting periods. It also identified whether there was a difference in each food groups consumed, diet diversity and animal source consumption score, and anthropometric status between the lent fasting and non-fasting periods, separately for fasting and non-fasting mothers. For this, McNemar’s test and Wilcoxon Signed Ranks Tests were used to check the statistical difference. Furthermore, the chapter addressed the factors associated with the underweight of lactating mothers in the study area. The factors included in the logistic regression to identify associated factors were from the socio-demographic and economic characteristics, maternal and child characteristics, water, sanitation and hygiene (WASH), feeding practices and household food security indicators. The underlying publication was published in *International Journal of Environmental Research and Public Health* in 2018, Vol 15(8), page 1-20.

Chapter 3 of the Thesis dealt with “Feeding Practices and Undernutrition in 6–23-Month-Old Children of Orthodox Christian Mothers in Rural Tigray, Ethiopia: Longitudinal Study”. Like chapter 2, this chapter focused on differences in the consumption pattern between the general 6-23-months-old children in the lent fasting and non-fasting periods, regardless of their mothers fasting status, and separately both for the 6-23-months-old children of the fasting and non-fasting mothers. It also highlighted differences between mean standard deviations of length-for-age, weight-for-length and weight-for-age z-scores between children of fasting and non-fasting

mothers during lent fasting period. It also stressed on the trend of the weight for age and diet diversity score of the whole population between the lent fasting and non-fasting periods. On top of these, the prevalence of child undernutrition (underweight, wasting and stunting) and associated factors were also addressed. McNemar's test, independent sample t-test, Wilcoxon Signed Ranks Tests and binary logistic regression were used to analyze the data. This article was published in *nutrients* in 2019, Vol 11, page 1-15.

Chapter 4 holds a paper 'Dietary Nutrient Intake of 12-23-Months-Old Children and Lactating Women in Rural Ethiopia; Adaptation and Validation of Calculator for Inadequate Micronutrient Intake (CIMI)' and was submitted also to *nutrients* journal in May 22, 2019. It focuses on two major issues. The first part was on the development of simple and rapid app for assessment of dietary nutrient intake using commonly consumed Ethiopian food items for Ethiopian population, followed by assessing the validity of the results produced by this software compared to internationally established nutritional assessment software called NutriSurvey. The second part of this chapter focused on identifying the estimated dietary energy and nutrient intakes and the prevalence of inadequacy of the 12-23-months-old children and lactating mothers during the lent fasting and non-fasting periods. Statistical analyses were done using descriptive statistics and Pearson correlation test to check the relationship of the results produced by CIMI and NutriSurvey, of each nutrients and energy for the 12-23-months old children and lactating women.

In chapter 5, the results presented in the three chapters (2, 3 and 4) are merged and discussed together. In this discussion, recent evidences in relation with the dietary pattern and nutritional status of children and women; in regard to religious fasting were found and compared and argued with/against our findings. Additionally, the development of CIMI software, and the assessment of



its validity compared to NutriSurvey as a reference are discussed with previous findings in the CIMI software developed for other countries, and other dietary assessment tools in its applicability. At the end of this chapter, the general conclusion drawn from the whole research findings in this study are presented.

Chapter 6 encompasses the general summary of the whole findings in the Thesis/Dissertation and is presented in both German and English languages.

### 1.3. References

1. Development Initiatives. Global Nutrition Report 2017: Nourishing the SDGs. Bristol, UK; 2017.
2. Development initiatives. 2018 Global nutrition report: Shining a light to spur action on nutrition [Internet]. Bristol, UK; 2018. 261 p. Available from: <http://ebrary.ifpri.org/cdm/ref/collection/p15738coll2/id/130399%5Cnhttps://www.ifpri.org/publication/hidden-hunger-approaches-tackling-micronutrient-deficiencies>
3. FAO/IFAD/UNICEF/WFP/WHO. The State of Food Security and Nutrition in the World 2018. Building climate resilience for food security and nutrition. Rome; 2018.
4. Biesalski Hans K, Jana T. Micronutrients in the life cycle: Requirements and sufficient supply. NFS J. 2018;11[March]:1–11. <https://doi.org/10.1016/j.nfs.2018.03.001>
5. Black RE, Allen LH, Bhutta ZA, Caulfield LE, de Onis M, Ezzati M, et al. Maternal and child undernutrition: global and regional exposures and health consequences. Lancet. 2008;371[9608]:243–60.
6. CSA. Ethiopia Demographic and Health Survey 2016. Addis Ababa; 2017.
7. Endalew B, Muche M, Tadesse S. Assesment of food security situation in Ethiopia: A review. Asian J Agric Res. 2015;9[2]:55–68.
8. IMF. The Federal Democratic Republic of Ethiopia : Poverty Reduction Strategy Paper : Growth and transformation Plan 2010/11-2014/15-Volume I. Washington D.C; 2011.
9. FAO. Ethiopia Situation Report - May 2017. Addis Ababa, Ethiopia; 2017.
10. FMOH. The Federal Democratic Republic of Ethiopia: National Nutrition Strategy. Addis

- Ababa, Ethiopia; 2008.
11. FMOH. National Nutrition Programme Republic of Ethiopia. Addis Ababa, Ethiopia; 2013.
  12. FMOH. National Nutrition Program 2016-2020. Addis Ababa, Ethiopia; 2016.
  13. Federal Democratic Republic of Ethiopia [FDRE]. SEQOTA declaration implementation plan [ 2016 – 2030]: Summary programme approach document. Addis Ababa, Ethiopia; 2016.
  14. Ahmed A, Abdulahi A, Shab-bidar S, Rezaei S. Nutritional Status of Under Five Children in Ethiopia : A Systematic Review and Meta-Analysis. *Ethiop J Health Sci.* 2017;27(2):175-188. doi: 10.4314/ejhs.v27i2.10.
  15. Akombi BJ, Agho KE, Merom D, Hall JJ, Renzaho AM. Multilevel analysis of factors associated with wasting and underweight among children under-five years in Nigeria. *Nutrients.* 2017;9[1]:44. doi: 10.3390/nu9010044.
  16. Alemu ZA, Ahmed AA, Yalew AW, Birhanu BS. Non random distribution of child undernutrition in Ethiopia: Spatial analysis from the 2011 Ethiopia demographic and health survey. *International Journal for Equity in Health.* 2016;15:198. DOI 10.1186/s12939-016-0480-z
  17. Ali Naser I, Jalil R, Wan Muda WM, Wan Nik WS, Mohd Shariff Z, Abdullah MR. Association between household food insecurity and nutritional outcomes among children in Northeastern of peninsular Malaysia. *Nutr Res Pract.* 2014;8(3):304-11. doi: 10.4162/nrp.2014.8.3.304.

18. Hagos S, Hailemariam D, WoldeHanna T, Lindtjørn B. Spatial heterogeneity and risk factors for stunting among children under age five in Ethiopia: A Bayesian geo-statistical model. *PLoS ONE*, 12(2): e0170785. doi:10.1371/journal.pone.0170785
19. Sohnesen TP, Ambel AA, Fisker P, Andrews C, Khan Q. Small area estimation of child undernutrition in Ethiopian woredas. *PLoS ONE* 12(4): e0175445. <https://doi.org/10.1371/journal.pone.0175445>
20. Haile D, Azage M, Mola T, Rainey R. Exploring spatial variations and factors associated with childhood stunting in Ethiopia: Spatial and multilevel analysis. *BMC Pediatr* [Internet]. 2016;16[1]:1–14. Available from: <http://dx.doi.org/10.1186/s12887-016-0587-9>
21. Akombi BJ, Agho KE, Hall JJ, Wali N, Renzaho AMN, Merom D. Stunting, wasting and underweight in Sub-Saharan Africa: A systematic review. *Int J Environ Res Public Health*. 2017; 14 [863]. doi:10.3390/ijerph14080863
22. Kandala N, Fahrmeir L, Klasen S, Priebe J. Geo-additive Models of Childhood Undernutrition in Three Sub-Saharan African Countries 1. *Popul Space Place*. 2009;473[December 2008]:461–73.
23. Tolulope A, Olubukola B. Spatial Pattern of Household Food Insecurity and Childhood Malnutrition in Akinyele Local Government Area, Nigeria. *International Journal of Health Sciences* 2015; 3(1):235-250. DOI: 10.15640/ijhs.v3n1a15
24. UNICEF. UNICEF ' s approach to scaling up nutrition. 2015.
25. Biza Zepro N. Food Taboos and Misconceptions Among Pregnant Women of Shashemene District, Ethiopia, 2012. *Sci J Public Heal* [Internet]. 2015;3[3]:410. Available from:

- <http://www.sciencepublishinggroup.com/journal/paperinfo.aspx?journalid=251&doi=10.11648/j.sjph.20150303.27>
26. Demissie T, Muroki NM, Kogi-Makau W. Food aversions and cravings during pregnancy: prevalence and significance for maternal nutrition in Ethiopia. *Food Nutr Bull*. 1998;19[1]:20–6.
  27. Hadush Z, Birhanu Z, Chaka M, Gebreyesus H. Foods tabooed for pregnant women in Abala district of Afar region, Ethiopia: an inductive qualitative study. *BMC Nutr* [Internet]. 2017;3[1]:40. Available from: <http://bmcnutr.biomedcentral.com/articles/10.1186/s40795-017-0159-x>
  28. Kariuki LW, Lambert C, Purwestri RC, Maundu P, Konrad H, Kariuki LW, et al. Role of food taboos in energy, macro and micronutrient intake of pregnant women in western Kenya. *Nutrition & Food Science*, 47(6):795-807. <https://doi.org/10.1108/NFS-09-2016-0146>
  29. Meyer-Rochow VB. Food taboos: Their origins and purposes. *Journal of Ethnobiology and Ethnomedicine* 2009, 5:18. doi:10.1186/1746-4269-5-18
  30. Riang’a RM, Broerse J, Nangulu AK. Food beliefs and practices among the Kalenjin pregnant women in rural Uasin Gishu County, Kenya. *Journal of Ethnobiology and Ethnomedicine* 2017;13:29. DOI 10.1186/s13002-017-0157-8.
  31. Zerfu TA, Umata M, Baye K. Dietary habits, food taboos, and perceptions towards weight gain during pregnancy in Arsi, rural central Ethiopia: a qualitative cross-sectional study. *J Health Popul Nutr* [Internet]. 2016;35[1]:22. Available from: <http://dx.doi.org/10.1186/s41043-016-0059-8>

32. McGovern ME, Krishna A, Aguayo VM, Subramanian S V. A review of the evidence linking child stunting to economic outcomes. *Int J Epidemiol.* 2017;46[4]:1171-1191. doi: 10.1093/ije/dyx017
33. Arzoaquoi SK, Essuman EE, Gbagbo FY, Tenkorang EY, Soyiri I, Laar AK. Motivations for food prohibitions during pregnancy and their enforcement mechanisms in a rural Ghanaian district. *Journal of Ethnobiology and Ethnomedicine* 2015; 11:59. DOI 10.1186/s13002-015-0044-0.
34. Herrador Z, Perez-Formigo J, Sordo L, et al. Low Dietary Diversity and Intake of Animal Source Foods among School Aged Children in Libo Kemkem and Fogera Districts, Ethiopia. *Plos one.* 2015;10(7):e0133435. DOI: 10.1371/journal.pone.0133435.
35. Ekwochi U, Osuorah CDI, Ndu IK, Ifediora C, Asinobi IN, Eke CB. Food taboos and myths in South Eastern Nigeria: The belief and practice of mothers in the region. *J Ethnobiol Ethnomed.* 2016; 12: 7. doi: 10.1186/s13002-016-0079-x
36. Getnet W, Aycheh W, Tessema T. Determinants of Food Taboos in the Pregnant Women of the Awabel District , East Gojjam Zone , Amhara Regional State in Ethiopia. *Adv Public Heal.* 2018;2018:6. Article ID 9198076, 6 pages.  
<https://doi.org/10.1155/2018/9198076>
37. Hm HT, Ahmed EM, N AAM. Food Taboos among Pregnant Women in Health Centers, Khartoum State Sudan, 2016. *International Journal of Science and Healthcare Research,* 2018; 3(1):13-25.
38. Maliwichi-nyirenda CP, Maliwichi LL, Maliwichi-nyirenda CP, Maliwichi LL. Studies on Ethno-Medicine Food Related Taboos Observed During Pregnancy in Malawi. *Stud*

- ethno-Medicine. 2017;10[2]:263–8.
39. Herrador Z, Perez-Formigo J, Sordo L, et al. Low Dietary Diversity and Intake of Animal Source Foods among School Aged Children in Libo Kemkem and Fogera Districts, Ethiopia. *Plos one*. 2015;10(7):e0133435. DOI: 10.1371/journal.pone.0133435.
  40. Hailelassie K, Mulugeta A, Girma M. Feeding practices, nutritional status and associated factors of lactating women in Samre Woreda, South Eastern Zone of Tigray, Ethiopia. *Nutr J* [Internet]. 2013;12[28]. <http://www.nutritionj.com/content/12/1/28>
  41. Picciano MF. Dietary Supplement Use in Women : Current Status and Future Directions Pregnancy and Lactation : Physiological Adjustments , Nutritional Requirements and the Role of Dietary Supplements 1. *J Nutr*. 2003;133:1997–2002.
  42. Roba KT, O'Connor TP, Belachew T, O'Brien NM. Seasonal variation in nutritional status and anemia among lactating mothers in two agro-ecological zones of rural Ethiopia: A longitudinal study. *Nutrition* [Internet]. 2015;31[10]:1213–8. Available from: <http://dx.doi.org/10.1016/j.nut.2015.03.007>
  43. Desalegn B, Lambert C, Riedel S, Negese T, Biesalski H. Feeding Practices and Undernutrition in 6–23-Month-Old Children of Orthodox Christian Mothers in Rural Tigray, Ethiopia: Longitudinal Study. *Nutrients* [Internet]. 2019;11[1]:138. Available from: <http://www.mdpi.com/2072-6643/11/1/138>
  44. Kumera G, Tsedal E, Ayana M. Dietary diversity and associated factors among children of Orthodox Christian mothers/caregivers during the fasting season in Dejen District, North West Ethiopia. *Nutrition & Metabolism*, 2018; 15:16. <https://doi.org/10.1186/s12986-018-0248-0>

45. Karl Knutsson E, Selinus R. Fasting in Ethiopia: An anthropological and nutritional study. *Am J Clin Nutr.* 1970;23[7]:956–69.
46. Trepanowski JF, Bloomer RJ. The impact of religious fasting on human health. *Nutrition Journal* 2010, 9(57). Available from: <http://www.nutritionj.com/content/9/1/57>.
47. Ali Z, Abizari A. Ramadan fasting alters food patterns, dietary diversity and body weight among Ghanaian adolescents. *Nutr J.* 2018; 17(75). <https://doi.org/10.1186/s12937-018-0386-2>
48. Sadeghirad B, Motaghipisheh S, Kolahdooz F, Zahedi MJ, Haghdoost AA. Islamic fasting and weight loss: A systematic review and meta-analysis. *Public Health Nutr.* 2014;17[2]:396–406.
49. Leiper J., Molla AM, Molla AM. Effects on health of fluid restriction during fasting in Ramadan. *Eur J Clin Nutr.* 2003;57[2]:S30–8.
50. Altuntas YE, Gezen FC, Sahoniz T, Kement M, Aydin H, Sahin F, et al. Ramadan fasting in patients with a stoma: a prospective study of quality of life and nutritional status. *Ostomy Wound Manag [Internet].* 2013;59[5]:26–32. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/23669258>
51. Faris MA-IE, Al-Holy MA. Implications of Ramadan intermittent fasting on maternal and fetal health and nutritional status: A review. *Med J Nutrition Metab.* 2014;7:107–18.
52. Awwad J, Usta IM, Succar J, Musallam KM, Ghazeeri G, Nassar AH. The effect of maternal fasting during Ramadan on preterm delivery: A prospective cohort study. *BJOG An Int J Obstet Gynaecol.* 2012;119[11]:1379–86.



53. RAKICIOĞLUNESLIŞAH, SAMUR G, TOPÇU A, AYAZ T. The effect of Ramadan on maternal nutrition and composition of breast milk. *Pediatr Int.* 2006;48[3]:278–83. doi: 10.1111/j.1442-200X.2006.02204.x.
54. Kiziltan G, Karabudak E, Tuncay G, Avsar F, Tuncay P, Mungan O, et al. Dietary intake and nutritional status of Turkish pregnant women during Ramadan. *Saudi Med J.* 2005;26[11]:1782–7.
55. Fasts FW, Fasts TP. Fasting and abstinence of the Coptic Orthodox Church of Alexandria [Internet]. Wikipedia. 2019 [cited 2019 Mar 19]. p. 3–7. Available from: [https://en.wikipedia.org/wiki/Fasting\\_and\\_abstinence\\_of\\_the\\_Coptic\\_Orthodox\\_Church\\_of\\_Alexandria](https://en.wikipedia.org/wiki/Fasting_and_abstinence_of_the_Coptic_Orthodox_Church_of_Alexandria)
56. Sarri KO, Kafatos AG, Higgins S. Is religious fasting related to iron status in Greek Orthodox Christians ? *Br J Nutr.* 2005;2001:198–203.
57. Sarri KO, Linardakis MK, Bervanaki FN, Tzanakis NE, Kafatos AG. Greek Orthodox fasting rituals: a hidden characteristic of the Mediterranean diet of Crete. *Br J Nutr* [Internet]. 2004;92:277–84. Available from: [http://www.journals.cambridge.org/abstract\\_S000711450400159X](http://www.journals.cambridge.org/abstract_S000711450400159X)
58. Sellassie SH, Mikael B. Worship in the Ethiopian Orthodx church [Internet]. The Ethiopian orthodox Tewahedo Church. 2019 [cited 2019 Mar 12]. p. 4–7. Available from: <https://www.ethiopianorthodox.org/english/ethiopian/worship.html>
59. Sarri KO, Tzanakis NE, Linardakis MK, Mamalakis GD, Kafatos AG. Effects of Greek orthodox christian church fasting on serum lipids and obesity. *BMC Public Health.* 2003;3[16].

60. Papadaki A, Vardavas C, Hatzis C, Kafatos A. Calcium , nutrient and food intake of Greek Orthodox Christian monks during a fasting and non-fasting week. Public. 2007;11[10]:1022–9.
61. Lazarou C, Matalas A. A critical review of current evidence , perspectives and research implications of diet-related traditions of the Eastern Christian Orthodox Church on dietary intakes and health consequences. Int J Food Sci Nutr. 2010; 61(7):739-758, DOI: 10.3109/09637481003769782
62. Koufakis T, Karras S, Antonopoulou V, Angeloudi E, Zebekakis P, Kotsa K. Effects of Orthodox religious fasting on human health: a systematic review. Eur J Nutr. 2017;56[8]:2439–55.
63. CSA. Population and Housing Census of Ethiopia. 2007.
64. Id ANB, Potts KS, Mulugeta A. How do pregnant and lactating women , and young children , experience religious food restriction at the community level ? A qualitative study of fasting traditions and feeding behaviors in four regions of Ethiopia. PLoS ONE 2018; 13(12): e0208408. <https://doi.org/10.1371/journal.pone.0208408>
65. EOTC/USAID. Patriarch of the Ethiopian Orthodox Tewahedo Church Endorses Nutrition Sermon Guide Encouraging Better Nutrition for Mothers and Children . 2016 Jan 12;1–2.
66. Desalegn BB, Lambert C, Riedel S, Negese T, Biesalski HK. Ethiopian Orthodox Fasting and Lactating Mothers : Longitudinal Study on Dietary Pattern and Nutritional Status in Rural Tigray , Ethiopia. Int J Environ Res Public Heal. 2018;15[1767]:1–20.
67. Alive and Thrive. IYCF practices, beliefs, and influences in Tigray Region, Ethiopia.

Addis Ababa, Ethiopia; 2010.

68. Mulat E, Alem G, Woyraw W, Temesgen H. Uptake of minimum acceptable diet among children aged 6 – 23 months in orthodox religion followers during fasting season in rural area, Dembecha, north West Ethiopia. BMC Nutr. 2019;5[18]:1–10.
69. Kumera G, Tsedal E, Ayana M. Dietary diversity and associated factors among children of Orthodox Christian mothers/caregivers during the fasting season in Dejen district, north west Ethiopia. Nutrition & Metabolism 2018; 15:16. <https://doi.org/10.1186/s12986-018-0248-0>
70. TEWODROS D. Beef Cattle Production System and Opportunities for Market Orientation in Borena Zone, Southern. MSc Thesis, Department of Animal Science, Haromaya University; 2008. Pp.139.
71. Avery A. Red Meat and Poultry Production and Consumption in Ethiopia and Distribution in Addis Ababa. International Livestock Research Institute 2004, Addis Abeba, Ethiopia.Pp.64.
72. DSM. Nutrition during the first 1 , 000 days. In: Vitamins in Motion for a brighter world. 2015. p. 1–24.
73. Williams J, Mai CT, Mulinare J. National Birth Defects Prevention Month and Folic Acid Awareness Week — January 2015 Updated Estimates of Neural Tube Defects Prevented by Mandatory Folic Acid Fortification — United States , 1995 – 2011. MMWR. 2015;64[1]:1–5.
74. Gebre B, Biadgilign S, Taddese Z, Legesse T, Letebo M. Determinants of malnutrition

- among pregnant and lactating women under humanitarian setting in Ethiopia. *BMC Nutr.* 2018;4[11]:1–8.
75. FAO/WHO. Human Vitamin and Mineral Requirements. Rome, Italy; 2001.
  76. FAO/WHO/UNU. Human energy requirements. Rome, Italy; 2001.
  77. FAO/WHO/UNU. Protein and Amino acid requirements in Human Nutrition. a of Joint a Joint. Geneva, Switzerland; 2001.
  78. National Academy of Sciences. Nutrition During Lactation. 4th Editio. Washington, D.C: National Academy Press; 1991. 326 p.
  79. Demissie T, Mekonen Y, Haider J. Agroecological comparison of leveles and correlates of nutritional status of women. *Ethiop J Heal Dev.* 2003;17[3]:189–96.
  80. WHO. Indicators for assessing infant and young child feeding practices. Part 3 Ctry Profiles. 2010;1–47.
  81. Roba KT, Connor TPO, Belachew T, Brien NMO. Infant and Young Child Feeding [ IYCF ] Practices Among Mothers of Children Aged 6 – 23 Months in Two Agro-ecological Zones of Rural Ethiopia. *International Journal of Nutrition and Food Sciences* 2016;5[3]:185–94.
  82. CSA. Ethiopia Demographic and Health Survey 2011. Addis Ababa; 2012.
  83. Stoecker BJ, Abebe Y. Context for food-based approaches for improved feeding of infants and children in Ethiopia. *Food Nutr Bull.* 2014;35[4]:S209–13.
  84. Mulugeta A, Hagos F, Kruseman G, Linderhof B, Stoecker B, Abraha Z, et al. Factors Contributing to Child Malnutrition in Tigray , Northern Ethiopia. *East Afr Med J.* 2010;87[6]:248–54.

85. FAO and FHI 360. Minimum Dietary Diversity for Women A Guide to Measurement. Rome; 2016. 82 p.
86. Dangura D, Gebremedhin S. Dietary diversity and associated factors among children 6-23 months of age in Gorche district, Southern Ethiopia: Cross-sectional study. BMC Pediatr [Internet]. 2017;17[1]:1–7. Available from: <http://dx.doi.org/10.1186/s12887-016-0764-x>
87. Roba KT, O'Connor TP, Belachew T, O'Brien NM. Variations between post- and pre-harvest seasons in stunting, wasting, and infant and young child feeding [IYCF] practices among children 6-23 months of age in lowland and midland agro-ecological zones of rural Ethiopia. Pan Afr Med J. 2016;24:1–9.
88. Weldehaweria NB, Misgina KH, Weldu MG, Gebregiorgis YS, Gebrezgi BH, Zewdie SW, et al. Dietary diversity and related factors among lactating women visiting public health facilities in Aksum town, Tigray, Northern Ethiopia. BMC Nutr [Internet]. 2016;2[38]. Available from: <http://bmcnutr.biomedcentral.com/articles/10.1186/s40795-016-0077-3>
89. Stoecker BJ, Abebe Y. Context for food-based approaches for improved feeding of infants and children in Ethiopia. Food and Nutrition Bulletin 2014;35[4]:209–13.
90. Arimond M, Ruel MT. Dietary diversity is associated with child nutritional status : Evidence from 11 demographic and health surveys. JNutr. 2004;134:2579–85.
91. FAO. Moving forward on choosing a standard operational indicator of women`s dietary diversity. 2015. 226 p.
92. Gewa CA, Murphy SP, Weiss RE, Neumann CG. Determining minimum food intake amounts for diet diversity scores to maximize associations with nutrient adequacy : an analysis of schoolchildren` s diets in rural Kenya. Public Health Nutrition 2014; 17(12), 2667–2673. doi:10.1017/S1368980014000469

93. Ersino G, Henry CJ, Zello GA. Suboptimal Feeding Practices and High Levels of Undernutrition among Infants and Young Children in the Rural Communities of Halaba and Zeway, Ethiopia. *Food Nutr Bull.* 2016;37[3]:409–24.
  
94. Tassew AA, Tekle DY, Belachew AB, Id MA. Factors affecting feeding 6 – 23 months age children according to minimum acceptable diet in Ethiopia : A multilevel analysis of the Ethiopian Demographic Health Survey. *PLoS ONE* 2019; 14(2): e0203098.  
<https://doi.org/10.1371/journal.pone.0203098>
  
95. Bhandari S, Sayami JT, Thapa P, Sayami M, Kandel BP, Banjara MR. Dietary intake patterns and nutritional status of women of reproductive age in Nepal: Findings from a health survey. *Arch Public Heal* [Internet]. 2016;74[1]:1–11. Available from:  
<http://dx.doi.org/10.1186/s13690-016-0114-3>
  
96. Jati IRAP, Widmer C, Purwestri RC, Wirawan NN, Gola U, Lambert C, et al. Design and validation of a program to identify inadequate intake of iron, zinc, and vitamin A. *Nutrition* [Internet]. 2014;30[11–12]:1310–7. Available from:  
<http://dx.doi.org/10.1016/j.nut.2014.03.015>
  
97. Federal Republic of Ethiopia DRM and FS. Disaster risk management and [ DRMFSS ] woreda disaster risk profile [ WDRP ] region : zone : woreda: eastern Tigray Ganta Ahfeshum February 2015 Addis Ababa Contents. Addis Ababa; 2015.

# **Chapter 2**

## **Paper I**

## **2. Ethiopian Orthodox Fasting and Lactating Mothers: Longitudinal Study on Dietary Pattern and Nutritional Status in Rural Tigray, Ethiopia**

Beruk Berhanu Desalegn<sup>1,2,\*</sup>, Christine Lambert<sup>2</sup>, Simon Riedel<sup>2</sup>, Tegene Negese<sup>1</sup> and Hans Konrad Biesalski<sup>2</sup>

<sup>1</sup>College of Agriculture, Hawassa University, Postal code: 05, Hawassa, Ethiopia; tegeengss38@gmail.com

<sup>2</sup>Institute of Biological Chemistry and Nutrition, University of Hohenheim, Garbenstr. 30, 70593 Stuttgart, Germany; christine.lambert@uni-hohenheim.de (C.L.); simon.riedel@uni-hohenheim.de (S.R.); biesal@uni-hohenheim.de (H.K.B.)

\* Correspondence: berhanuberuk@gmail.com Tel.: +251-941-048-918

**A version of this chapter has been published as**

Desalegn, B., Lambert, C., Riedel, S., Negese, T., Biesalski, H.K. (2018). Ethiopian Orthodox Fasting and Lactating Mothers: Longitudinal Study on Dietary Pattern and Nutritional Status in Rural Tigray, Ethiopia. *International Journal of Environmental Research and Public Health*, 15(1767). doi:10.3390/ijerph15081767.



# **Ethiopian Orthodox Fasting and Lactating Mothers: Longitudinal Study on Dietary Pattern and Nutritional Status in Rural Tigray, Ethiopia**

## **Abstract**

About half of Ethiopians belong to the Orthodox Tewahedo religion. Annually, more than 200 days are dedicated to religious fasting, which includes abstaining from all types of food, animal source foods, and water. However, the association of fasting with undernutrition remains unknown in Ethiopia. Therefore, dietary pattern and nutritional status of lactating women during lent fasting and non-fasting periods were studied, and predictor variables for maternal underweight were identified. To achieve this, lactating mothers in lent fasting (N = 572) and non-fasting (N = 522) periods participated from rural Tigray, Northern Ethiopia. Average minimum diet diversity (MDD-W) was computed from two 24-h recalls, and nutritional status was assessed using body mass index (BMI). Binary logistic regression was used to identify potential predictors of maternal underweight. Wilcoxon signed-rank (WSRT) and McNemar's tests were used for comparison of the two periods. The prevalence of underweight in fasting mothers was 50.6%. In the multivariate logistic regression model, younger age, sickness in the last four weeks preceding the survey, fasting during pregnancy, lactation periods, grandfathers' as household decision makers, previous aid experience, non-improved water source, and not owning chicken were positively associated with maternal underweight. In WSRT, there was no significant ( $p > 0.05$ ) difference on maternal body weight and BMI between periods. The average number of meals, diet diversity, and animal source foods (ASFs), consumption scores were significantly increased in non-fasting compared to fasting periods in both fasting and non-fasting mothers ( $p < 0.001$ ,  $p$

$< 0.05$ , and  $p < 0.001$ , respectively). Consumption of dark green leafy vegetables was higher in the fasting period (11%) than non-fasting (3.6%), in the study population. As a conclusion, Ethiopian Orthodox fasting negatively affected maternal nutritional status and dietary pattern in rural Tigray, Northern Ethiopia. To reduce maternal malnutrition in Ethiopia, existing multi-sectoral nutrition intervention strategies, should include religious institutions in a sustainable manner.

Keywords: lactating mothers; Ethiopian Orthodox lent fasting; ASFs consumption; underweight; Ethiopia

## **2.1. Introduction**

Undernutrition is a serious public health problem worldwide. It is the underlying cause for 3.5 million deaths, 35% of the disease burden in children younger than 5 years old, 11% of total global disability-adjusted life-years (DALYs) and accounted for at least 20% of maternal mortality [1–4]. According to Food and Agriculture Organization of the United Nations (FAO) estimates, the global prevalence of chronically undernourished people increased from 777 million (10.6%) in 2015 to 815 Million (11%) in 2016; however, the deterioration was most severe in sub-Saharan Africa. Eastern Africa is one of the four sub-regions in sub-Saharan Africa, where one-third (33.9%) of the population was estimated to be undernourished in 2016 [3].

Good nutritional status of women is important for their good health and working capacity, as well as for the health of their offspring [1]. During pregnancy and lactation, women are more

vulnerable to undernutrition than others at reproductive age, due to increased energy and nutrient requirements [5–7]. Globally in 2011, the prevalence of anemia in pregnant women was 38.2% vs. 29% in non-pregnant women [8]. According to the Demographic and Health Survey (DHS) in Ethiopia, the prevalence of anemia in pregnant or lactating women (19% and 29%) was higher than in non-lactating-non-pregnant women (15% and 21%) [DHS 2011, 2016, respectively] [9,10]. Likewise, the prevalence of underweight ( $\text{BMI} < 18.5 \text{ kg/m}^2$ ) in lactating mothers was 25–55%, which was higher than non-pregnant-non-lactating women (22%) [5,9–12].

Food taboos refer to those foods which are strictly forbidden for health, cultural, and religious reasons [13–16]. In Ethiopia, food taboos are thought to have been established during pregnancy as a means of protecting the health of women and their babies. As a result, scholars in Ethiopia focused on the exploration of food taboos during pregnancy and its association with their nutritional status and feeding practices [15–18]. However, lactating mothers who even need more nutrients than pregnant women [5,6,12], are ignored in this regard. Religious fasting is one of the categories of dietary or food taboos, which may affect the dietary intake and nutritional status of lactating mothers, and these may lead to undernourished breastfed children. The main difference between religious fasting and the ordinary type of food taboo is its momentary nature, in which abstention from eating animal source foods and/or from eating certain foods is done for fixed periods of time [19]. According to previous studies, Ramadan fasting affected nutritional status, dietary nutrient intake, birth outcome, breast milk composition, and health status of women in reproductive age [20–28].

In Ethiopia, about half (44%) of the whole population are Ethiopian Orthodox Christians [9]. Religious fasting from any animal source foods and abstaining from any foods and water for some hours daily is mandatory. However, if it is practiced, it affects over 200 days annually [29].

Apart from this, children less than seven years old, soldiers, severely ill or weak people, pregnant, and lactating mothers are permitted to eat both animal source foods and other foods including water, without abstention during the religious fasting periods or days [19,29,30]. Nonetheless, information is lacking on the effect of religious fasting on the dietary and nutritional status of women in reproductive age. Therefore, the purpose of this study was to explore the dietary pattern and nutritional status of lactating women during Ethiopian Orthodox Lent fasting and non-fasting periods, and to identify potential predictors associated with maternal underweight in rural Tigray, Northern Ethiopia.

## **2.2. Materials and Methods**

### **2.2.1. Study Area, Design, Participants, and Sampling**

The study was conducted in the Genta Afeshum woreda of rural Tigray, Northern Ethiopia. The woreda covers an area of 1636 km<sup>2</sup> with a total population of 99,112, and almost all people (99%) are Orthodox Christians. The woreda reside at an altitude between 2045 and 3314 masl. The woreda is classified as a hotspot for food insecurity [31–33]. In the woreda, drought, hail storms, and livestock diseases are the major disaster risks; followed by human diseases, crop diseases, pests, and flooding. Additionally, deforestation, water pollution, and soil erosion are the major environmental problems; whereas, high dependency syndrome, poor economic conditions, land shortage, severe shortage of drinking water, and poor saving are among the major vulnerability factors at the household level [34].

The study had a community-based longitudinal survey design, to assess the nutritional status and dietary pattern of lactating mothers. The data was collected during the Ethiopian Orthodox lent fasting period (Fasting of Jesus Christ, 15 February 2017 to 15 April 2017) and non-fasting

periods (1 May 2017 to 30 May 2017). The sample size was calculated based on the prevalence of underweight in lactating mothers in the Tigray region, using the formula for estimating single population proportion and considering a 95% of confidence interval for true prevalence, and a relative precision (d) of 5%. In lactating mothers, the prevalence of underweight (BMI < 18.5 kg/m<sup>2</sup>) was 25% elsewhere in Tigray [5]. The total number of lactating mothers was estimated to be 3369, which was less than 10,000; therefore, the finite source population size correction formula was used. Additionally, 10% was considered as non-responses and dropout rates. Moreover, a 1.5 design effect was used on the final calculated sample size, and the final total sample size was 575.

Multi-stage systematic random sampling was applied to obtain representative samples for the study. At first, Genta Afeshum was randomly selected out of the three GIZ Ethiopia, Nutrition Sensitive Agriculture (NSA) project woreda's in Tigray region. Out of twenty rural kebeles in the Genta Afeshum woreda, seven were randomly selected. Then, the list of households which had lactating mothers with children aged between 6 to 23 months old, who fulfilled the inclusion criteria, was prepared for the seven kebeles (lowest local administrative unit) by health extension workers at the nearby health posts. Subsequently, the samples were chosen using systematic random sampling techniques.

### **2.2.2. Data Collection**

Ten trained and well experienced data collectors who were fluent in Tigrigna, Amharic, and English languages were recruited. Additionally, before conducting the main survey, the questionnaire was translated to Tigrigna by a professional translator and verified by data collectors. Then the translated questionnaire was pre-tested for its appropriateness, by

administering it to lactating mothers around Mekele, and corrections were made. Structured and semi-structured questionnaires were prepared to collect information on socio-demographic and economic characteristics, maternal and child characteristics, water, sanitation and hygiene (WASH), feeding practices, and household food security indicators [35].

Before conducting the study, the whole study protocol was ethically approved by the Institutional Review Board of the College of Health Sciences at Hawassa University and the Tigray Region Health Bureau in Ethiopia; and the ethical review committee of Landesärztekammer Baden-Württemberg, Germany. Permissions from Genta Afeshum Woreda Health Office were also obtained. After the purpose of the study was explained to the study participants, agreement to participate in the study was documented by signing the informed consent. Each participant was also told that the collected information was confidential, and whenever she wanted to discontinue, withdrawal from the study was possible.

### **2.2.3. Minimum Women Diet Diversity Score**

The minimum-diet diversity score (MDD-W) was obtained by (a) collecting two 24-h dietary recalls; (b) categorizing as consumed or not consumed of the food group, considering the minimum amount (15 g) of any food items or the sum of food items eaten under a given food group and giving a score of 1 if consumed, otherwise 0 if not; (c) calculating the diet diversity score for each two days using 10 food groups, as the summation of consumed food groups or scored 1 for each day separately; and (d) taking the average diet diversity score of the two days, as an individual MDD-W score. The 10 food groups used for calculating MDD-W score were grains, white roots and tubers, and plantains; pulses (beans, peas, and lentils); nuts and seeds;

dairy; meat, poultry, and fish; eggs; dark green leafy vegetables; other vitamin A-rich fruits and vegetables; other vegetables; and other fruits [36].

#### **2.2.4. Household Food Insecurity Information**

Household food insecurity data was collected using the household food insecurity access scale (HFIAS). It is the measure of the degree of food insecurity (access) in the household in the past 4 weeks (30 days). The questionnaire encompassed nine questions, which assess the occurrence of food insecurity in increasing level. Under each of the nine questions, frequency of occurrence questions were a follow up to determine how often the condition (1 = rarely, 2 = sometimes, 3 = often) occurred.

The Household Food Insecurity Access Prevalence (HFIAP) status indicator, was used to determine prevalence of food insecurity to report household food insecurity. Using the HFIAP indicators four categories, the households were categorized into four levels of household food insecurity (access). These were food secure, mild, moderately, and severely food insecure. After creating these four categories, the HFIAP was calculated as the number of households with a given food insecurity category divided by the total number of households with household food insecure access category, multiplied by 100 [35].

#### **2.2.5. Wealth Index**

Principal component analysis (PCA) was carried out to compute the wealth index. To achieve this, 17 variables: transport animals (horse/donkey/mule), goat and/or sheep, household owns kerosene or lamp, owns bed, chair, table, radio, electric-mitad, bicycle, mobile phone, non-mobile phone, animal drawn cart, motor bicycle, TV, electricity, windows and separate room for animal, which could indicate the living standard of the surveyed area were included in the

analysis. The first factor that explained most of the variation (86.3%) was used to group study households. Finally, the wealth tertile was performed and categorized as higher, medium, and lower [37].

#### **2.2.6. Anthropometry**

Weighing body mass of the mothers was conducted using a portable digital scale (Seca 770, Hanover, Germany), working with a powered battery and measured to the nearest 0.1 kg. For height measurement, a disassembling plastic height measuring board with a sliding head bar was used and measured to the nearest 0.1 cm. During weight and height measurements, the mothers were advised to remove their jackets until they had light clothes to minimize the weight due to clothes.

The measurements (height and weight) were carried out using standardized equipment and procedures in duplicate and the average values were used. Additionally, the BMI of the mothers was calculated as the weight of the mothers in kilograms divided by the square of their height in meters. The BMI values of mothers were classified in three categories as underweight, normal, and overweight ( $<18.5$ ,  $18.5\text{--}24.99$  and  $25\text{ kg/m}^2$ ), respectively [38].

#### **2.2.7. Data Management and Analysis**

Before submitting the data, variable coding was conducted in SPSS version 20. Following this, the data was entered, cleaned, and analyzed. First, frequency and crosstab were conducted to determine completeness of data and to present the results in descriptive statistics (frequency and percent). Association between outcome and potential explanatory variables, was assessed using bivariate analysis with a confidence level of 95% to declare the statistical significance. Out of all the independent variables entered in bivariate logistic regression, seventy variables with p-values



of less than 0.25 were entered for multivariable logistic regression to identify predictor variables for maternal underweight ( $\text{BMI} < 18.5 \text{ kg/m}^2$ ).  $p\text{-value} < 0.05$  was used to declare the variables as predictors for the outcome variable. Hosmer and Lemeshow test and C-statistics (AUC) were conducted to assess fitness of the final model. Meanwhile, multi-collinearity was checked using the variance inflation factor (VIF) and standard error with  $<10$  and  $<2$  as a cutoff point, respectively. Maternal nutritional status was defined as underweight ( $\text{BMI} < 18.5 \text{ kg/m}^2$ ) and normal if  $\text{BMI} \geq 18.5 \text{ kg/m}^2$ , since the interest of this study was being underweight for logistic regression. Normality of continuous data was checked using the Kolmogorov-Smirnov test. Non-normally distributed data was analyzed using the Wilcoxon signed-rank test; whereas the dichotomous data was analyzed using McNemar's test to detect a difference between fasting and non-fasting periods, of fasting and non-fasting mothers for their dietary pattern, separately.

### **2.3. Results**

A total of 572 lactating mothers who were inhabitants of Genta Afeshum district of rural Tigray, Northern Ethiopia participated during the 1st round of data collection (Ethiopian Orthodox lent fasting) period, with a response rate of 99.5%. Out of those participants in the 1st survey, 522 mothers were again involved during the non-fasting period, with a response rate of 90.8%. The lost for follow-up was due to age ( $>23$  months) of breastfeeding child, migration to other areas, and absence during the re-visit at the 2nd round survey period.

In this study, all mothers were from the Tigray ethnic group (100%) and almost all were Ethiopian Orthodox Christians (99.7%). Majority of mothers were married (84.8%); whereas, the rest were single, widowed, and divorced mothers. More than sixty percent of mothers were thirty years old and above in the study. However, about eight out of ten mothers were married before

the age of 21 years. The proportion of mothers who were illiterate and housewives were 35.1% and 79.2%, respectively. Nearly sixty percent of mothers had children of one year and above. Majority of the households were food insecure (71%), male-headed (80%), and taking aid (food, cash, in kind) (89%) before the survey period, from local government or humanitarian organizations. Likewise, 61.4% and 75% of households had family members of more than five and resided in an area more than 15 km away from the largest market in the district, respectively (Table 2.1).

**Table 2.1** Socio-demographic and economic characteristics of the study participants (n = 572) in rural Tigray, Northern Ethiopia (February–June, 2017).

Characteristics (n = 572)		Number	Percent
Ethnicity of mother	Tigray	572	100
Religion of mother	Orthodox	570	99.7
	Other	2	0.3
Current maternal age (years)	>30	227	39.7
	≤30	345	60.3
Mother age at marriage (years)	>20	116	20.3
	≤20	456	79.7
Mother age at giving 1st birth (years)	>20	279	48.8
	≤20	293	51.2
Family size	≤5	221	38.6
	>5	351	61.4
Household previous aid experience	No	64	11.2
	Yes	508	88.8
Household main income decision maker	Husband	129	22.6
	Jointly husband and wife	352	61.5
	Wife	74	12.9
	Grand father	17	3.0
Age of child (months)	≤12	241	42.1
	13–18	200	35.0
	>18	131	22.9
Mother previous credit experience from local	No	438	76.6

institutes	Yes	134	23.4
Distance from woreda market (km)	≤15	143	25.0
	>15	429	75.0
Marital status	Married	485	84.8
	Others	87	15.2
Mother education	Illiterate	201	35.1
	Literate	371	64.9
Mother occupation	Housewives	453	79.2
	Farmer	81	14.2
	Daily laborer	15	2.6
	Business owner	15	2.6
	Employee	8	1.4
Household head	Father	458	80.1
	Mother	114	19.9
Household food security status	Food secured	166	29.0
	Mildly food insecure	181	31.6
	Moderately food insecure	213	37.3
	Severely food insecure	12	2.1
	Higher	149	26.1
Wealth tertile	Medium	237	41.4
	Lower	186	32.5
Household chicken owning	No	247	43.2
	Yes	325	56.8

As shown in Table 2.2, 28% and 31% of mothers were fasting during the pregnancy and lactation period of the indexed child, respectively. Beside this, 65% of mothers did not change their food intake during their lactation period. Very small proportion of mothers (13.8%), had illness within four weeks preceding the fasting period. Most of the mothers (91%), used antenatal care (ANC) services four times or more during pregnancy of the indexed child. However, mothers who received post-natal care (PNC) services after delivery were 43%. (Table 2.2).

Mothers less than or equal to thirty years of age, were about 1.7 times more likely to be underweight than those above thirty years (Adjusted Odds Ratio (AOR) = 1.73). Conversely, mothers who had children between 13 and 18 months of age had two times higher odds of being underweight, compared to those mothers who had children between six to twelve months of age

(AOR = 2.01). Households having no access to improved water and not owning chickens, were more likely to have underweight mothers compared to their counterparts (AOR = 1.57, 1.73, respectively). However, family size, marital status, family planning method used, distance from woreda market, number of ANC visits, PNC attendance, toilet presence, and wealth index had no association with maternal underweight (Table 2.3).

**Table 2.2** Maternal health and feeding practices (n = 572) in rural Tigray, Northern Ethiopia (February–June 2017).

Characteristics (n = 572)		Numbers	Percent
Fasting during lactation	No fasting	394	68.9
	Fasting	178	31.1
Fasting during pregnancy	No fasting	414	72.4
	Fasting	158	27.6
Change of food intake	No	374	65.4
	Yes	198	34.6
Number of visits in antenatal care	≥4	518	90.6
	<4	54	9.4
Attendance of postnatal care service after delivery	No	329	57.5
	Yes	243	42.5
Household water source	Improved	379	66.3
	Non-improved	193	33.7
Household toilet presence	No	112	19.6
	Yes	460	80.4
Household garbage pit presence	No	74	12.9
	Yes	498	87.1
Family planning methods	No	274	47.9
	Yes	298	52.1
Sickness last 4 weeks	No	493	86.2
	Yes	79	13.8

The odds of being underweight for mothers fasting during their pregnancy and lactation period of the indexed child, were 1.7 and 2.8 times higher than those who were not fasting in both periods (AOR = 1.75, 2.82), respectively. Additionally, mothers who had sickness within four weeks preceding the fasting period had 3.6 times higher odds of being underweight, compared to those who were healthy (AOR = 3.62). Similarly, households whose decision makers were

grandfathers and had received aid were 6.0 and 2.9 times more likely to have underweight mothers, compared to households with husbands as decision makers and those which did not receive aid (AOR = 6.02, 2.86, respectively).

**Table 2.3** Association of some socio-demographic, health, and feeding practice variables with maternal underweight (BMI) during Ethiopian Orthodox Lent fasting period in Rural Tigray, Northern Ethiopia (n = 572).

Variables	Underweight		COR (95% CI)	AOR (95% CI)
	No	Yes		
	Number (%)	Number (%)		
Mother age				
>30	158 (69.6)	69 (30.4)	1	1
≤30	222 (64.3)	123 (35.7)	1.27 (1.11, 2.71)	1.73 (1.11, 2.71)*
Mother fasting status during pregnancy period of indexed child				
No	301 (72.7)	113 (27.3)	1	1
Yes	79 (50.0)	79 (50.0)	2.66 (1.82, 3.89)*	1.75 (1.11, 2.75) *
Mother fasting status during lactation period of indexed child				
No	292 (74.1)	102 (25.9)	1	1
Yes	88 (49.4)	90 (50.6)	2.93 (2.02, 4.24)*	2.82 (1.80, 4.42)*
Family size				
≤5	159 (71.9)	62 (28.1)	1	1
>5	221 (63.0)	130 (37.0)	1.51 (1.05, 2.17)	1.42 (0.87, 2.32)
Decision maker on main household income				
Husband	83 (64.3)	46 (35.7)	1	1
Jointly husband and wife	246 (69.90)	106 (30.1)	0.78 (0.51, 1.19)	0.65 (0.39, 1.09)
Wife	46 (62.2)	28 (37.8)	1.10 (0.61, 1.98)	0.94 (0.35, 2.47)
Grand father	5 (29.4)	12 (70.6)	4.33 (1.44,13.06)*	6.02 (1.47, 24.80)*
Marital status				
Married	330 (68.0)	155 (32.0)	1	1
Others	50 (57.5)	37 (42.5)	1.58 (0.99, 2.51)	1.28 (0.52, 3.17)
Age of indexed child				
≤12 months	171 (71.0)	70 (29.0)	1	1
13–18	120 (60.0)	80 (40.0)	1.63 (1.10, 2.42)*	2.01 (1.27, 3.18)*
>18	89 (67.9)	42 (32.1)	1.15 (0.73, 1.83)	1.14 (0.67, 1.96)
Household previous experience to aid				

No	53 (82.8)	11 [17.2]	1	1
Yes	327 (64.4)	181 [35.6]	2.67 (1.36, 5.23)*	2.86 (1.32, 6.20)*
Family planning use status				
Yes	205 (68.8)	93 (31.2)	1	1
No	175 (63.9)	99 (36.1)	1.25 (0.88, 1.76)	1.12 (0.74, 1.69)
Distance from woreda market				
≤15	101 (70.6)	42 (29.4)	1	1
>15	279 (65.0)	150 (35.09)	1.29 (0.86, 1.95)	1.26 (0.72, 2.21)
ANC number visited during the pregnancy period of indexed child				
≥4	349 (67.4)	169 (32.6)	1	1
<4	31 (57.4)	23 (42.6)	1.53 (0.87, 2.71)	0.97 (0.49, 1.94)
Mother PNC				
Yes	169 (69.5)	74 (30.5)	1	1
No	211 (64.1)	118 (35.9)	1.28 (.90, 1.82)	1.01 (0.66, 1.55)
Household consumption water source				
Improved	263 (69.4)	116 (30.6)	1	1
Non-improved	117 (60.6)	76 (39.4)	1.47 (1.03, 2.12)*	1.57 (1.02, 2.43)*
Toilet presence in the household				
Yes	311 (67.6)	149 (32.4)	1	1
No	69 (61.6)	43 (38.4)	1.30 (0.85, 1.99)	1.48 (0.90, 2.43)
Household chicken owning status				
Yes	229 (70.5)	96 [29.5]	1	1
No	151 (61.1)	96 [38.9]	1.52 (1.07, 2.15)*	1.73 (1.15, 2.61)*
Mother sickness last one month preceding the survey				
No	350 (71.0)	143 (29.0)	1	1
Yes	30 (38.0)	49 (62.0)	3.99 (2.44, 6.55) *	3.62 (2.00, 6.54) *
Wealth status tertile				
Higher	134 (72.0)	52 (28.0)	1	1
Medium	153 (64.6)	84 (35.4)	0.64 (0.41, 1.02)	0.98 (0.53, 1.83)
Lower	93 (62.4)	56 (37.6)	0.91 (0.60, 1.40)	0.86 (0.52, 1.41)

COR = Crude Odds Ratio; AOR = Adjusted Odds Ratio; CI = confidence interval; Hosmer and Lemeshow test showed  $p = 0.197$ ; C-statistic: Area under the Curve (AUC) = 0.766; 95% CI (0.724–0.809); \*=significantly associated at  $p < 0.05$ .

Variables included in the adjusted model logistic regression: Mother age, Mother fasting status during pregnancy period of indexed child, Mother fasting status during lactation period of indexed child, Family size, Decision maker on main household income, Marital status, Age of

indexed child, Household previous experience to aid, Family planning use status, Distance from woreda market, ANC number visits during the pregnancy period of indexed child, Mother PNC, Household consumption water source, Toilet presence in the household, Household chicken owning status, Mother sickness last one month preceding the survey, and Wealth status tertile.

A small fraction of mothers (1.4%) had height < 145 cm, indicating stunting. Nearly one-third of mothers had body weight < 45 kg (32.5%) and BMI < 18.5 kg/m<sup>2</sup> (33.6%), in the study population. However, the prevalence was higher in fasting mothers than non-fasting mothers, both in the fasting and non-fasting period. However, the prevalence of overweight (BMI ≥ 25 kg/m<sup>2</sup>) was lower in fasting mothers than non-fasting in both periods, as shown in Table 2.4.

**Table 2.4** Food consumption and anthropometric status of lactating mothers during Ethiopian Orthodox lent fasting and non-fasting periods at rural Tigray, Northern Ethiopia.

Variables		Lent Fasting Period (n = 572)			Non-Fasting Period (n = 522)		
		Non-Fasting Mothers	Fasting Mothers	Total	Non-Fasting Mothers	Fasting Mothers	Total
		n (%)	n (%)	n (%)	n (%)	n (%)	n (%)
Height (cm)	<145	7 (1.8)	1 (0.6)	8 (1.4)			
	≥145	387 (98.2)	177 (99.4)	564 (98.6)			
Weight (kg)	≤45 kg	106 (26.9)	80 (44.9)	186 (32.5)	97 (26.8)	71 (44.4)	168 (32.2)
	>45 kg	288 (73.1)	98 (55.1)	386 (67.5)	265 (73.2)	89 (55.6)	354 (67.8)
BMI (kg/ m <sup>2</sup> )	<18.5	102 (25.9)	90 (50.6)	192 (33.6)	90 (24.9)	81 (50.6)	171 (32.8)
	18.5–24.99	277 (70.3)	86 (48.3)	363 (63.4)	259 (71.5)	77 (48.1)	336 (64.3)
	≥25	15 (3.8)	2 (1.1)	17 (3.0)	13 (3.6)	2 (1.2)	15 (2.9)
Diet diversity score (MDD-W)	<3	199 (50.5)	101 (56.7)	300 (52.5)	163 (45.0)	73 (45.6)	236 (45.2)
	3	161 (40.9)	65 (36.5)	226 (39.5)	149 (41.2)	65 (40.6)	214 (41.0)
	>3	34 (8.6)	12 (6.7)	46 (8.0)	50 (13.8)	22 (13.8)	72 (13.8)
Number of meals	<3	119 (30.2)	64 (36.0)	183 (32.0)	29 (8.0)	15 (9.4)	44 (8.4)
	3	226 (57.4)	92 (51.7)	318 (55.6)	299 (82.6)	131 (81.9)	430 (82.4)
	>3	49 (12.4)	22 (12.4)	71 (12.4)	34 (9.4)	14 (8.8)	48 (9.2)
Number of cups of coffee	<2	208 (52.8)	105 (59.0)	313 (54.7)	215 (59.4)	94 (58.8)	309 (59.2)
	≥2	186 (47.2)	73 (41.0)	259 (45.3)	147 (40.6)	66 (41.2)	213 (40.8)

Grain consumption	394 (100.0)	178 (100.0)	572 (100.0)	362 (100.0)	160 (100.0)	522 (100.0)
Pulse consumption	383 (97.2)	170 (95.5)	553 (96.7)	355 (98.1)	157 (98.1)	512 (98.1)
Nuts and seeds consumption	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)
Dairy products consumption	8 (2.0)	0 (0.0)	8 (1.4)	23 (6.4)	9 (5.6)	32 (6.1)
Meat, poultry and fish consumption	2 (0.5)	0 (0.0)	2 (0.3)	60 (16.6)	19 (11.9)	79 (15.1)
Eggs consumption	4 (1.0)	2 (1.1)	6 (1.0)	19 (5.2)	12 (7.5)	31 (5.9)
Dark green leafy vegetables consumption	45 (11.4)	18 (10.1)	63 (11.0)	10 (2.8)	9 (5.6)	19 (3.6)
Vitamin A-rich fruits and vegetables consumption	4 (1.0)	1 (0.6)	5 (0.9)	0 (0.0)	1 (0.6)	1 (0.2)
Other vegetables consumption	294 (74.6)	130 (73.0)	424 (74.1)	265 (73.2)	116 (72.5)	381 (73.0)
Other fruits consumption	0 (0.0)	2 (1.1)	2 (0.3)	0 (0.0)	0 (0.0)	0 (0.0)

Our findings demonstrated that the number of mothers who were in the fasting sub-group and had a diet diversity score (MDD-W) less than three, were slightly more in the fasting (56.7%) than non-fasting period (45.6%). However, the proportion of fasting and non-fasting mothers who scored a diet diversity greater than three, increased from the fasting period (6.7% and 8.6%, respectively) to non-fasting period (13.8%). Fasting mothers reduced consumption of coffee in the fasting period (41%), whereas non-fasting mothers (47.2%) did not. In both study periods, all mothers consumed grains, but there was no consumption of nuts and seeds. Almost all mothers ate pulses in fasting (96.7%) and non-fasting (98.1%) periods. Conversely, except two fasting mothers in fasting period, there was no consumption of other fruit groups in both periods. Furthermore, mothers who consumed vitamin A-rich fruits and vegetables preceding the survey days in both periods were less than 1%. The overall consumption of dark green, leafy vegetables was lower in the non-fasting (3.6%) than fasting period (11%). The consumption of other vegetable groups was almost the same during non-fasting (73%) and fasting (74.1%) periods. Dairy products; meat, poultry and fish; and eggs food groups were consumed by a very low



number of both fasting (0%, 0% and 1.1%) and non-fasting mothers (1.4%, 0.3% and 1%) during the fasting period, respectively. However, consumption of the three ASFs significantly increased in both fasting and non-fasting mothers, in the non-fasting period compared to fasting period, as shown in Table 2.4. According to McNemar's test, there was no significant difference on the proportion of fasting mothers who consumed pulses ( $p = 0.289$ ), dark green leafy vegetables ( $p = 0.078$ ), vitamin A-rich fruits and vegetables ( $p = 1.000$ ), and other vegetables groups ( $p = 1.000$ ), between fasting and non-fasting periods. However, the proportions of fasting and non-fasting mothers who consumed eggs, were significantly higher in non-fasting than fasting periods ( $p = 0.006$ ,  $p = 0.003$ , respectively). Similarly, the consumption of pulses and other vegetable groups by non-fasting mothers were not significantly different ( $p = 0.332$ ,  $p = 0.716$ , respectively), between fasting and non-fasting periods. However, the number of non-fasting mothers who consumed dairy; meat, poultry and fish products; eggs; and dark green leafy vegetables were significantly different ( $p = 0.002$ ,  $p = 0.001$ ,  $p = 0.003$  and  $p = 0.001$ , respectively), between fasting and non-fasting periods (Table 2.5).

Using the Wilcoxon signed-rank test, the median BMI, body weight of fasting mothers, and number of cups of coffee consumed were not significantly different between fasting and non-fasting periods. Similarly, the median BMI and body weight of non-fasting mothers were not statistically different between the two periods (Tables 2.6 and 2.7). However, in both fasting and non-fasting mothers, the number of meals eaten ( $p = 0.001$ ), MDD-W ( $p = 0.037$  and  $p = 0.014$ ), and ASFs consumption scores ( $p = 0.001$ ) were significantly increased in the non-fasting than fasting periods, respectively. However, the number of cups of coffee consumed by non-fasting mothers was significantly ( $p = 0.009$ ) higher in fasting period than in the non-fasting period (Tables 2.6 and 2.7).

**Table 2.5** Food consumption status of fasting and non-fasting mothers, during Ethiopian Orthodox lent fasting and non-fasting periods in rural Tigray, Northern Ethiopia.

Parameters	Fasting Mothers (a, n = 160)			Non-Fasting Mothers (b, n = 362)		
	Lent Fasting Period	Non-Fasting Period	Significance	Lent Fasting Period	Non-Fasting Period	Significance
	n (%)	n (%)	p-value	n (%)	n (%)	p-value
Grain consumption	160 (100.0)	160 (100.0)	NA	362 (100.0)	362 (100.0)	NA
Pulse consumption	153 (95.6)	157 (98.1)	0.289	351 (97.0)	356 (98.3)	0.332
Nuts and seeds consumption	0 (0.0)	0 (0.0)	NA	0 (0.0)	0 (0.0)	NA
Dairy products consumption	0 (0.0)	9 (5.1)	NA	7 (1.9)	23 (6.4)	0.002 *
Meat, poultry and fish consumption	0 (0.0)	19 (11.9)	NA	2 (0.6)	60 (16.6)	≤0.001 *
Eggs consumption	2 (1.2)	12 (7.5)	0.006 *	4 (1.1)	19 (5.2)	0.003 *
Dark green leafy vegetables consumption	18 (11.2)	9 (5.6)	0.078	41 (11.3)	10 (2.8)	≤0.001 *
vitamin A-rich fruits and vegetables consumption	1 (0.6)	1 (0.6)	1.000	4 (1.1)	0 (0.0)	NA
Other vegetables consumption	116 (72.5)	116 (72.5)	1.000	270 (74.6)	265 (73.2)	0.716
Other fruits consumption	1 (0.6)	0 (0.0)	NA	0 (0.0)	0 (0.0)	NA

Data analysis using McNemar's test, significant level at  $p < 0.05$ , a < b, NA-the data was not appropriate for analysis; \*=significantly associated at  $p < 0.05$ .

**Table 2.6** Comparison of fasting mothers' food consumption and anthropometric status between Orthodox lent fasting and non-fasting periods in rural Tigray, Northern Ethiopia.

Variables (N = 160)	Data Collection Period						
	Lent Fasting (a)			Non-Fasting (b)			
	Mean (SD)	Median (IQR)	Range (Min, Max)	Mean (SD)	Median (IQR)	Range (Max, Min)	Sign
Weight (kg)	46.18 (5.55)	45.55 (42.23, 49.48)a	(35.40, 65.60)	45.98 (5.49)	45.65 (42.23, 48.90)a	35.30, 65.40	0.054
BMI (kg/m <sup>2</sup> )	18.86 (2.14)	18.47 (17.55, 20.05)a	(14.79, 27.99)	18.79 (2.18)	18.48 (17.49, 20.0)a	14.75, 28.21	0.051
Diet Diversity Score (MDD-W)	2.57 (0.54)	2.50 (2.0, 3.0)a	(1.00, 4.0)	2.69 (0.58)	3.0 (2.0, 3.0)b	1.50, 4.0	0.037
ASFs Score	0.01 (0.06)	0 (0, 0)a	(0, 0.50)	0.13 (0.27)	0 (0, 0)b	0, 1.0	≤0.001
Number of meals	2.79 (0.49)	3.0 (2.5, 3.0)a	(1.50, 4.0)	3 (0.29)	3.0 (3.0, 3.0)b	2.0, 4.50	≤0.001
Number of cup of coffee	2.64 (3.4)	1.5 (0, 3.0)a	(0, 12.0)	1.74 (1.67)	1.5 (0, 3.0)a	0, 7.50	0.058

Data analysis using Wilcoxon Signed Ranks Test significant level at  $p < 0.05$ ,  $a < b$ .

**Table 2.7** Comparison of non-fasting mothers' food consumption and anthropometric status between Orthodox Lent fasting and non-fasting periods in rural Tigray, Northern Ethiopia.

Variables (N = 362)	Data Collection Period						
	Lent Fasting (a)			Non-Fasting (b)			
	Mean (SD)	Median (IQR)	Range (Min, Max)	Mean (SD)	Median (IQR)	Range (Max, Min)	Sign
Weight (kg)	49.30 (6.41)	48.50 (45.10, 53.10)a	35.50, 72.00	49.20 (6.61)	48.50 (44.60, 53.13)a	35.70, 74.00	0.092
BMI (kg/m <sup>2</sup> )	20.08 (2.36)	19.68 (18.53, 21.34)a	15.87, 29.91	20.04 (2.43)	19.64 (18.50, 21.48)a	15.57, 29.79	0.086
Diet Diversity Score (MDD-W)	2.63 (0.53)	2.50 (2.00, 3.00)a	1.00, 4.00	2.73 (0.55)	3.00 (2.00, 3.00)b	1.50, 4.50	0.014
ASFs Score	0.02 (0.12)	0.00 (0.00, 0.00)a	0.00, 1.00	0.16 (0.31)	0.00 (0.00, 0.00)b	0.00, 1.50	≤0.001
Number of meals	2.84 (0.50)	3.00 (2.50, 3.00)a	1.00, 4.50	3.01 (0.27)	3.00 (3.00, 3.00)b	2.00, 4.50	≤0.001
Number of cup of coffee	2.12 (1.96)	1.5 (0.00, 3.00)b	0.00, 9.00	1.76 (1.57)	1.5 (0.00, 3.00)a	0.00, 6.00	0.009

Data analysis using Wilcoxon Signed Ranks Test significant level at  $p < 0.05$ ,  $a < b$ .

## **4. Discussion**

This study assessed whether nutritional status and dietary pattern of lactating mothers were different between Ethiopian Orthodox lent fasting and non-fasting periods. It also determined factors associated with maternal underweight in rural Tigray, Northern Ethiopia.

### **4.1. Nutritional Status of Lactating Mothers**

Adult stature is the collective outcome of the interaction between environment and inheritances, over the critical growing period of a person [39]. Prior evidence has demonstrated that short maternal height was associated with increased offspring mortality, underweight, and stunting in infancy, childhood, and later in an adult age [39,40]. In our study, prevalence of maternal stunting was 1.4%, which was less than previous findings in Ethiopia [5,9,10,41]. Shorter women are believed to have reduced protein and energy stores, smaller size of their reproductive organs, and smaller pelvis diameter. This may limit fetal development in the uterus, increases risk for mother and child complication during delivery, and later infant growth through reduced breast milk quantity and quality, resulting in stunted children. Thus, appropriate feeding behavior is important for pregnant women health, and later for bearing healthier and well grown new born babies. This finding suggests that health extension agents working in the rural communities should advise mothers on appropriate feeding behavior during the pregnancy period.

The overall prevalence of maternal underweight in this study was between 32.6 and 33.6%. The prevalence was lower than previous findings in Tigray region (34–55%), and Dedo and Seqa-Chekorsa districts (41%), in South-west Ethiopia [9–12]. Conversely, the prevalence was higher than in other studies in the Tigray and Oromia regions of Ethiopia [5,41–44]. The latter could be related to climate phenomenon ‘El Nino’, which caused the strongest famine in Ethiopia, where

the impact seriously affected Tigray region. It could also be related to differences in feeding practices, study population, and period [45].

Shockingly, prevalence of underweight was 51% in fasting mothers compared with non-fasting mothers (25%). The result was consistent with studies conducted on lactating mothers living in the midland agro-ecology of Tigray region, which was 57% [12]. Similarly, the BMI was significantly lower among fasting adults than non-fasting adults in Greek Orthodox Christians [28]. This could be related to the almost 317 kcal difference between fasting and non-fasting adults, in end-holy days of fasting periods [46].

In the present study, prevalence of overweight was between 1.1–3.8%, which was relatively comparable with the Ethiopian Demographic and Health Survey (EDHS) 2011 report (2.9%) in Tigray, and elsewhere (1.3–1.8%) in the region [5,24,38], but lower than the EDHS (2016) report at national level (6%) and Tigray region (4.9%). In urban women, it was reported (6.2–25.3%) in Ethiopia, Bengal district in India (5.4%), and Nepal (6.3–24.8%) [44,47–52]. This might be because we attributed this finding to rural people [10], where most could engage in heavy physical activities and walking over long distances to access services due to the mountainous topography [53,54].

Mothers thirty years of age and younger were 1.7 times more likely to be underweight than those above thirty. That conforms with other studies conducted in Ethiopia and Nepal, resulting in higher prevalence of underweight in younger women [10,47]. In the current study, more than 27% of mothers fasted during their pregnancy period, and these mothers had 1.7 times more odds to be underweight than not fasting during the same period. In a previous study, more than one-third of mothers were fasting during their pregnancy period in Oromia region, Ethiopia [15]. Similarly, those mothers fasting during lactation period were 2.9 times more exposed to

underweight, than those who did not practice. Mothers who had children between the age of 13 and 18 months, were twice more likely to be underweight compared to those who had a child aged 6–12 months. Similar result was observed in a study conducted by Hailelassie and his colleagues in Northern Ethiopia [5]. This might be due to increased nutritional requirements of the growing child, effort for child care in connection with food intake by the mother that is not increased or even decreased.

Disease is one of the immediate causes of maternal and child undernutrition [1]. Mothers who had any illness in the last four weeks preceding the survey were 3.6 times more frequently underweight than healthy mothers. In a study conducted in the Limu area of Southern Ethiopia, maternal sickness was positively associated with maternal underweight [55]. This could be related to decreased food intake and absorption, alteration of metabolism, and increment in nutritional requirements. Marital status of lactating mothers was not significantly associated with maternal underweight. Prior evidence has also demonstrated that marital status was not associated with Ethiopian women [56]. Otherwise, birth spacing has important implications for the health and nutritional status of mothers and their children [57]. In our study, family planning use was not associated with maternal underweight, which is inconsistent with the study conducted in the Tena district of Oromia region, Ethiopia. In the latter, the prevalence of family planning utilization was higher (65%) than our finding, which was 52% [58].

Good care during pregnancy is important for the health of the mother and development of the unborn baby. The findings of this study indicated that a major proportion (91%) of mothers had at least four ANC services during their pregnancy period, and were not significantly associated with maternal underweight. This result was equal with the regional coverage of Tigray (91%), but higher than the national prevalence (62%) [10]. The postnatal period is a critical phase in the

lives of a mother and the newborn baby. In this period, major changes occur, but it is the most neglected time for the provision of quality services. As a result, the rates of provision of skilled care are lower after childbirth when compared to rates before and during childbirth [59]. In this study, prevalence of PNC coverage was 42.5%, which is comparable with previous findings in Tigray region (45.4%); however, PNC attendance was not significantly associated with maternal undernutrition. Household size was not significantly associated with maternal underweight in the study. This coincides with studies in Southern Ethiopia and Tigray region [5,41,55]. However, it was inconsistent with one study conducted in Nekemte town, Oromia region in Ethiopia [44]. The difference might be a higher proportion of lactating mothers who lived in rural households with many family members, in our case. Grandfathers as a decision maker for the household were associated with maternal underweight. This could be related to sharing the household income to more family members or to the loss (death, departure) of the husband resulting in lower working capacity and income, or due to most lactating mothers in grandfather headed households being younger, which is associated with maternal underweight in our case. It has also been reported that mothers who had more children, decreases the resources allocated including food, resulting in underweight [60]. Access to safe drinking water, sanitation and hygiene (WASH) services is a fundamental element of healthy communities and has an important positive impact on child and maternal nutrition [61,62].

One-third of households included in this study, had non-improved water sources for household consumption. The odds of being underweight for mothers from non-improved water sources were 1.6 higher than those from households with improved water sources. This may be due to the fact of frequent illness related to water borne diseases and contamination. Accordingly, child undernutrition was associated with source of drinking water in Iraq and sub-Saharan Africa

[63,64]. In contrast, toilet presence in the household was not associated with maternal underweight. This might apply to the majority of households included in the study, one of the successes of the health extension program in Ethiopia. In our study, households not owning chickens were 1.8 times more likely to have underweight mothers, than those who owned chickens. Similarly, the proportion of mothers who ate more diversified foods were higher in households which owned chicken, than those from households not owning chicken. Prior research in Ethiopia, indicated that owning livestock in the household was associated with a higher diet diversity score [65]. Studies in three East African countries also evidenced that, in households owning livestock, the prevalence of child stunting was low [66]. Thus, promoting chicken husbandry may also improve the low consumption of animal source foods, and the diversity of food to be eaten at large. Among households involved in the study, more than a quarter (32.5%) were at the lowest wealth tertile. According to the EDHS report, a lower proportion of households (23%) was at the lowest wealth quantile [10]. Of thirty-four woreda's in Tigray region, thirty-one were food insecure, including our study district [33]. The results from the present study showed that more than two-thirds (71%) of households were food insecure. Comparably, the prevalence of food insecurity was 76% in East Bedawacho district of Southern Ethiopia [67]. However, the prevalence was lower than in a study conducted in two agro-climatic zones in Sidama, Southern Ethiopia, which was 82% [68].

#### **4.2. Dietary Patterns of Lactating Mothers**

According to the essential nutrition action (ENA), mothers are recommended to take at least two additional meals during their lactation period [69]. In the present study, nearly two-thirds of the mothers (65.4%) did not change the food intake during their lactation period. This result is lower than a study conducted in Samre district which reported (71%), but higher than findings in Raya



area (59%) of Tigray region. This could be related to the interval in the study periods and study area [5,41].

Lactating mothers who ate more than three times a day were 9–12%. This result agreed with findings in the Tigray, Oromia, and Southern regions in Ethiopia [11,41]. One-third of lactating mothers ate less than three times on average of the two days preceding the survey during fasting period, which is lower than the expected three meals to have in a day of a normal adult in real context. Prior research in Samre district of Northern Ethiopia, showed that the proportion of lactating mothers who had less than three meals in the last 24-h preceding their survey was 27% [5], which is comparable with our findings. However, this prevalence reduced to 8.4% in the non-fasting period, which is comparable with the prevalence (7.5%) of pregnant women in Gambela town, Western Ethiopia [70]. Similarly, the number of meals eaten both by fasting and non-fasting mothers were significantly increased after two months of fasting period. The latter should be further studied for explanation.

Diet diversity is one important dimension of diet quality, and a proxy indicator for higher micronutrient adequacy [36]. Majority of mothers had a diet diversity score of three and less in the fasting period (92%) and non-fasting period (87%). This could indicate that dietary micronutrient inadequacy is high in the study population. The median diet diversity score for both sub-groups of fasting and non-fasting mothers, increased significantly in non-fasting period. This was associated with higher consumption of ASFs and a higher frequency of meals in our and related studies [65,71,72]. Thus, in the Ethiopian Orthodox fasting period, the feeding practice of lactating mothers is sub-optimal. This finding suggests that nutrition education, which will improve feeding practices should involve religious leaders in a sustainable manner. The number of mothers who had consumed more cups of coffee was higher in the fasting than non-

fasting period, in the whole study population. Likewise, non-fasting mothers took significantly more coffee in the fasting period than non-fasting. However, consuming more coffee during the fasting period, where plant-based foods are the sole source of minerals, could reduce their bioavailability, leading to increased risk for micronutrient deficiencies, especially of iron, calcium, and zinc [73]. However, many fasting mothers consumed more cups of coffee in the non-fasting than fasting period. This could be referred to some Ethiopian Orthodox religion monarchist, who preached to followers not to consume coffee during fasting periods.

In this study, cereals, pulses, and other vegetables were the main food groups commonly consumed by lactating mothers. This result agrees with previous studies in Ethiopia [65,71,74]. These plant foods usually contain dietary components that compromise digestion and inhibit absorption of vital nutrients. For example, phytic acid chelates multivalent ions such as zinc, calcium, and iron; therefore, their bioavailability reduced [75]. This finding suggests that traditional processing techniques which can improve the bioavailability of minerals should be promoted in the community.

Dark green, leafy vegetables are important plant sources of micronutrients like iron, calcium, and vitamin A [76]. However, their consumption was very low in the non-fasting period, both in fasting (5.6%) and non-fasting (2.8%) mothers. However, the result was lower than that previously reported for Axum town (19%) in Northern Ethiopia. The discrepancy between these two findings could be related to better market access, which is less in this study due to the rural area. However, the number of non-fasting mothers who consumed dark green, leafy vegetables was significantly higher in the fasting period than non-fasting period. This could be related to a potential to choose and consume more diversified food, especially ASFs which were less consumed in fasting period. Furthermore, almost all lactating mothers did not consume nuts and

seeds, fruits, and vitamin A rich fruits and vegetables food groups at all in both study periods, because of a lack of cultivation on local farms and unavailability in the market [77]. Thus, activities which can improve the dietary diversity including nuts and seeds, dark green leafy vegetables, and vitamin A and C rich fruits consumption, as well as meal frequency are inevitable. Moreover, the consumption of fruits which are rich in vitamin C should be taught to improve the mineral bioavailability of plant foods, which are the predominate source of most nutrients for a given community.

The proportion of lactating mothers who consumed ASFs was significantly lower in the fasting than non-fasting period. In Northern Ethiopia, particularly in rural Tigray, most strict Ethiopian Orthodox Christians abstain from eating animal source foods during the lent and other fasting periods or days, including Wednesday and Friday. During these fasting periods, the demand for cattle meat was observed to be low, resulting in closure of abattoirs or minimizing the service provided [78]. A previous study, reported that more than 85% of butcher houses were closed in Addis Ababa during Wednesday and Friday, which are Orthodox Christians fasting days of the week [79].

This study has much strength and some limitations. It is the first study in Ethiopia, which assessed the effect of religious fasting on maternal nutrition, particularly lactating mothers, who need more nutrients than pregnant women. Moreover, the study was done in an area, where almost all people were Ethiopian Orthodox Christians to minimize research bias. In addition, the study had a longitudinal nature, which considered a large sample size, and two 24-h dietary recall data for each study period. On top of these, this study also determined the differences in dietary and nutritional status during fasting and non-fasting periods, for both fasting and non-fasting mothers' sub-groups, separately. However, the study only considered the long Ethiopian

Orthodox lent fasting period out of the seven official fasting periods. This study did not measure micronutrient levels or bio-markers in the blood, to assess micronutrient deficiencies reflecting the consequences of qualitative malnutrition. Underweight only indicates energetic malnutrition, representing the tip of the iceberg.

## **5. Conclusions**

It has been found that the prevalence of underweight among fasting, lactating mothers during Ethiopian Orthodox fasting periods is high and a serious public health problem in the district. The dietary pattern [diet diversity and number of meals eaten] of lactating mothers is sub-optimal, both in fasting and non-fasting periods. Cereals and pulses are the dominant food groups eaten in the district, known to have a high amount of anti-nutrients like phytic acid, which limit the bioavailability of minerals for absorption. Therefore, activities which will improve dietary diversity including nuts and seeds, dark green leafy vegetables, and vitamin A and C rich fruits consumption, as well as meal frequency are urgently needed. Additionally, consuming vitamin C rich fruits and use of traditional processing techniques, which can improve the bioavailability of minerals should be promoted. Furthermore, promoting chicken husbandry may also improve the low consumption of animal source foods. Though moderate intake of coffee is beneficial, consuming coffee closely to meals to be eaten will decrease the absorption of non-heme iron from plant source foods and should be discouraged. Thus, households with younger mothers and non-improved water sources should get priority in nutritional intervention. Generally, multi-sectoral nutrition intervention strategies should include religious institutions to reduce maternal malnutrition in Ethiopia. Otherwise, nutrition education activities will not lead to sustainable improvements.

Author Contributions: B.B.D., C.L., S.R. and H.K.B. involved in research design; B.B.D. conducted the research; B.B.D. analyzed the data; B.B.D. wrote the paper; B.B.D., C.L., S.R., T.N. and H.K.B. edited draft manuscript; the final content was the primary responsibility of B.B.D. All authors read and approved the final manuscript.

Funding: This study was financially supported by German Federal Ministry for Economic Cooperation and Development (BMZ).

Acknowledgments: We would like to thank data collectors, Genta Afeshum Woreda Health Office Nutrition coordinator Gebretsadekane, and GIZ Nutrition Sensitive Agriculture project staff specially Habteab and Muez, in Tigray region, Ethiopia.

Conflicts of Interest: The authors declare no conflict of interest.

## **6. References**

1. Black, R.E.; Allen, L.H.; Bhutta, Z.A.; Caulfield, L.E.; De Onis, M.; Ezzati, M.; Mathers, C.; Rivera, J.; Maternal and Child Undernutrition Study Group. Maternal and child undernutrition: Global and regional exposures and health consequences. *Lancet* 2008, 371, 243–260. [CrossRef]
2. Caulfield, L.E.; De Onis, M.; Blössner, M.; Black, R.E. Undernutrition as an underlying cause of child deaths associated. *Am. J. Clin. Nutr.* 2004, 80, 193–198. [CrossRef] [PubMed]
3. FAO; IFAD; UNICEF; WFP; WHO. *The State of Food Security and Nutrition in the World*; FAO: Rome, Italy, 2017; p. 132.
4. Müller, O.; Krawinkel, M. Malnutrition and Health in Developing Countries. *Can. Med. Assoc. J.* 2005, 173, 279–286. [CrossRef] [PubMed]

5. Hailelassie, K.A.; Mulugeta, M.; Girma, M. Feeding practices, nutritional status and associated factors of lactating women in Samre Woreda, South Eastern Zone of Tigray, Ethiopia. *Nutr. J.* 2013, 12, 28. [CrossRef] [PubMed]
6. Picciano, M.F. Dietary Supplement Use in Women: Current Status and Future Directions Pregnancy and Lactation: Physiological Adjustments, Nutritional Requirements and the Role of Dietary Supplements 1. *J. Nutr.* 2003, 133, 1997–2002. [CrossRef] [PubMed]
7. Weldehaweria, N.B.; Misgina, K.H.; Weldu, M.G.; Gebregiorgis, Y.S.; Gebrezgi, B.H.; Zewdie, S.W.; Ngusse, H.A.; Gebrewa, H.G.; Alemu, W. Dietary diversity and related factors among lactating women visiting public health facilities in Aksum town, Tigray, Northern Ethiopia. *BMC Nutr.* 2016, 2, 38. [CrossRef]
8. WHO. The Global Prevalence of Anemia in 2011; World Health Organization: Geneva, Switzerland, 2015; pp. 1–48.
9. Central Statistical Agency (CSA). Ethiopia Demographic and Health Survey 2011; Central Statistical Agency: Addis Ababa, Ethiopia; ICF International Calverton: Calverton, MD, USA, 2012; p. 452.
10. Central Statistical Agency (CSA). Ethiopia Demographic and Health Survey 2016; Central Statistical Agency: Addis Ababa, Ethiopia; ICF International Calverton: Calverton, MD, USA, 2017; p. 551.
11. Alemayehu, M.; Argaw, A.; Geberemariam, A. Factors Associated with Malnutrition among Lactating Women in Subsistence Farming Households from Dedo and Seqa-Chekorsa Districts, Jimma Zone, 2014. *Dev. Ctry. Stud.* 2015, 5, 114–122.

12. Roba, K.T.; O'Connor, T.P.; Belachew, T.; O'Brien, N.M. Seasonal variation in nutritional status and anemia among lactating mothers in two agro-ecological zones of rural Ethiopia: A longitudinal study. *Nutrition* 2015, 31, 1213–1218. [CrossRef] [PubMed]
13. Demissie, T.; Muroki, N.M.; Kogi-Makau, W. Food aversions and cravings during pregnancy: Prevalence and significance for maternal nutrition in Ethiopia. *Food Nutr. Bull.* 1998, 19, 20–26. [CrossRef]
14. Meyer-Rochow, V.B. Food taboos: Their origins and purposes. *J. Ethnobiol. Ethnomed.* 2009, 5, 1–10. [CrossRef] [PubMed]
15. Zepro, N.B. Food Taboos and Misconceptions among Pregnant Women of Shashemene District, Ethiopia, 2012. *Sci. J. Public Health* 2015, 3, 410–416. [CrossRef]
16. Zerfu, T.A.; Umeta, M.; Baye, K. Dietary habits, food taboos, and perceptions towards weight gain during pregnancy in Arsi, rural central Ethiopia: A qualitative cross-sectional study. *J. Health Popul. Nutr.* 2016, 35, 22. [CrossRef] [PubMed]
17. Hadush, Z.; Birhanu, Z.; Chaka, M.; Gebreyesus, H. Foods tabooed for pregnant women in Abala district of Afar region, Ethiopia: An inductive qualitative study. *BMC Nutr.* 2017, 3, 40. [CrossRef]
18. Vasilevski, V.; Carolan-olah, M. Ethiopian pregnancy food taboos. *J. Clin. Nurs.* 2016, 25, 3069–3075. [CrossRef] [PubMed]
19. Knutsson, E.K.; Selinus, R. Fasting in Ethiopia: Anthropological and nutritional study. *Am. J. Clin. Nutr.* 1970, 23, 956–969. [CrossRef] [PubMed]
20. Altuntas, Y.E.; Gezen, F.C.; Sahoniz, T.; Kement, M.; Aydin, H.; Sahin, F.; Okkabaz, N.; Oncel, M. Ramadan fasting in patients with a stoma: A prospective study of quality of life and nutritional status. *Ostomy Wound Manag.* 2013, 59, 26–32.

21. Almond, D.; Mazumder, B. Health Capital and the Prenatal Fasting during Pregnancy Health Capital and the Prenatal Environment: The Effect of Maternal Fasting during Pregnancy; Working paper 2007-22; Federal Reserve Bank of Chicago: Chicago, IL, USA, 2011; p. 74.
22. Awwad, J.; Usta, I.M.; Succar, J.; Musallam, K.M.; Ghazeeri, G.; Nassar, A.H. The effect of maternal fasting during Ramadan on preterm delivery: A prospective cohort study. *BJOG Int. J. Obstet. Gynaecol.* 2012, 119, 1379–1386. [CrossRef] [PubMed]
23. Faris, M.E.; Al-Holy, M.A. Implications of Ramadan intermittent fasting on maternal and fetal health and nutritional status: A review. *Med. J. Nutr. Metab.* 2014, 7, 107–118.
24. Fayet-Moore, F.; Kim, J.; Sritharan, N.; Petocz, P. Impact of breakfast skipping and breakfast choice on the nutrient intake and body mass index of Australian children. *Nutrients* 2016, 8, 487. [CrossRef] [PubMed]
25. Kiziltan, G.; Karabudak, E.; Tuncay, G.; Avsar, F.; Tuncay, P.; Mungan, O.; Meral, P. Dietary intake and nutritional status of Turkish pregnant women during Ramadan. *Saudi Med. J.* 2005, 26, 1782–1787. [PubMed]
26. Leiper, J.B.; Molla, A.M.; Molla, A.M. Effects on health of fluid restriction during fasting in Ramadan. *Eur. J. Clin. Nutr.* 2003, 57, S30–S38. [CrossRef] [PubMed]
27. Rakicio ğlu, N.; Samur, G.; Topçu, A.; Ayaz, T. The effect of Ramadan on maternal nutrition and composition of breast milk. *Pediatr. Int.* 2006, 48, 278–283. [CrossRef] [PubMed]
28. Sarri, K.O.; Tzanakis, N.E.; Linardakis, M.K.; Mamalakis, G.D.; Kafatos, A.G. Effects of Greek orthodox christian church fasting on serum lipids and obesity. *BMC Public Health* 2003, 3, 16. [CrossRef] [PubMed]



29. Karle, C.S. Fasting among Orthodox Christians in Ethiopia: The Symbolic and Nutritional Significance of the Fast for Reproductive Women. Bachelor's Thesis, Department of Public Health, University of Copenhagen, Copenhagen, Denmark, 2017; p. 35.
30. EOTC/USAID. Patriarch of the Ethiopian Orthodox Tewahedo Church Endorses Nutrition Sermon Guide Encouraging Better Nutrition for Mothers and Children; USAID: Addis Ababa, Ethiopia, 2019; pp. 1–2.
31. Ludwig, C. Nutrition Baseline Survey-Ethiopia-For the Global Programme Food and Nutrition Security and Enhanced Resilience; Bioversity International: Rome, Italy, 2017; p. 90.
32. Tigray Region Agriculture and Rural Development Bureau. Report on Food Security and Nutrition Status of 31 Woreda's of the Region; Tigray Region Agriculture and Rural Development Bureau: Mekele, Ethiopia, 2016.
33. WFP. Food Security and Vulnerability in Selected Towns of Tigray Region, Ethiopia: Vulnerability Assessment and Mapping (VAM); WFP-Ethiopia: Addis Ababa, Ethiopia, 2009; p. 52. Available online: <http://documents.wfp.org/stellent/groups/public/documents/ena/wfp221389.pdf?iframe> [accessed on 24 January 2018].
34. FDRE Disaster Risk Management and Food Security Sector [FDRE/DRMFSS]. Woreda Disaster Risk Profile (WDRP); Tigray region; Food Security Sector: Addis Ababa, Ethiopia, 2015; p. 276.
35. Coates, J.F.; Swindale, A.; Bilinsky, P. Household Food Insecurity Access Scale (HFIAS) for Measurement of Household Food Access: Indicator Guide (v. 3); Food and Nutrition

- Technical Assistance Project; Academy for Educational Development: Washington DC, USA, 2007; p. 36.
36. FAO and FHI 360. Minimum Dietary Diversity for Women a Guide to Measurement; FAO: Rome, Italy, 2016; p. 82.
37. Abate, K.H.; Belachew, T. Women's autonomy and men's involvement in child care and feeding as predictors of infant and young child anthropometric indices in coffee farming households of Jimma Zone, South West of Ethiopia. *PLoS ONE* 2017, 12, e0172885. [CrossRef] [PubMed]
38. Cogill, B. Anthropometric Indicators Measurement Guide; Food and Nutrition Technical Assistance [FANTA] Project; FHI: Washington, DC, USA, 2003; p. 93.
39. Addo, O.Y.; Stein, A.D.; Fall, C.H.; Gigante, D.P.; Guntupalli, A.M.; Horta, B.L.; Kuzawa, C.W.; Lee, N.; Norris, S.A.; Prabhakaran, P.; et al. Maternal height and child growth patterns. *J. Pediatr.* 2013, 163, 549–554. [CrossRef] [PubMed]
40. Özaltın, E.; Hill, K.; Subramanian, S.V. Association of Maternal Stature with Offspring Mortality, Underweight, and Stunting in Low- to Middle-Income Countries. *JAMA* 2010, 303, 1507–1516. [CrossRef] [PubMed]
41. Sitotaw, I.K.; Kiday, H.; Adama, Y. Comparison of nutritional status and associated factors of lactating women between lowland and highland communities of District Raya, Alamata. *BMC Nutr.* 2017, 3, 61. [CrossRef]
42. Biru, K.; Jima, A.; Abeya, S. Prevalence of Chronic Energy Malnutrition and Maternal Health Service Utilizations among Lactating Mothers in Adama District, Oromia Region, Eastern Ethiopia. *J. Food Process. Technol.* 2017, 8, 1–5.

43. Hailu, C.; Dejene, T.; Tebeje, B. Prevalence and Socio-Demographic/Economic Risk Factors of Low and High Body Mass Index of Urban versus Rural Pregnant Women at Booking for Antenatal Care in Governmental Health Centers of Jimma Zone, Southwest Ethiopia. *Open Access Libr. J.* 2015, 2, e1672.
44. Hundera, T.D.; Gemede, H.F.; Wirtu, D.; Kenie, D.N. Nutritional Status and Associated Factors among Lactating Mothers in Nekemte Referral Hospital and Health Centers, Ethiopia. *Food Sci. Qual. Manag.* 2015, 35, 64–72. [CrossRef]
45. Food and Agriculture Organization of the United Nations (FAO). *FAO in Ethiopia, El Niño Response Plan*; FAO: Rome, Italy, 2016.
46. Sarri, K.O.; Linardakis, M.K.; Bervanaki, F.N.; Tzanakis, N.E.; Kafatos, A.G. Greek Orthodox fasting rituals: A hidden characteristic of the Mediterranean diet of Crete. *Br. J. Nutr.* 2004, 92, 277–284. [CrossRef] [PubMed]
47. Bhandari, S.; Sayami, J.T.; Thapa, P.; Sayami, M.; Kandel, B.P.; Banjara, M.R. Dietary intake patterns and nutritional status of women of reproductive age in Nepal: Findings from a health survey. *Arch. Public Health* 2016, 74, 1–11. [CrossRef] [PubMed]
48. Awoke, A.; Awoke, T.; Alemu, S.; Megabiaw, B. Prevalence and associated factors of hypertension among adults in Gondar, Northwest Ethiopia: A community based cross-sectional study. *BMC Cardiovasc. Disord.* 2012, 12, 113. [CrossRef] [PubMed]
49. Abrha, S.; Shiferaw, S.; Ahmed, K.Y. Overweight and obesity and its socio-demographic correlates among urban Ethiopian women: Evidence from the 2011 EDHS. *BMC Public Health* 2016, 16, 636. [CrossRef] [PubMed]

50. Gali, N.; Tamiru, D.; Tamrat, M. The Emerging Nutritional Problems of School Adolescents: Overweight/Obesity and Associated Factors in Jimma Town, Ethiopia. *J. Pediatr. Nurs.* 2017, 35, 98–104. [CrossRef] [PubMed]
51. Tigga, P.L.; Sen, J. Maternal Body Mass Index Is Strongly Associated with Children Z-Scores for Height and BMI. *J. Anthropol.* 2016, 2016, 10. [CrossRef]
52. Tebekaw, Y.; Teller, C.; Colón-Ramos, U. The burden of underweight and overweight among women in Addis Ababa, Ethiopia. *BMC Public Health* 2014, 14, 1126. [CrossRef] [PubMed]
53. Alemu, E.; Atnafu, A. Prevalence of Overweight and/or Obesity and Associated Factors among High School Adolescents in Arada Sub city, Addis Ababa, Ethiopia. *J. Nutr. Food Sci.* 2014, 4, 2. [CrossRef]
54. Anteneh, Z.A.; Gedefaw, M.; Tekletsadek, K.N.; Tsegaye, M.; Alemu, D. Risk Factors of Overweight and Obesity among High School Students in Bahir Dar City, North West Ethiopia: School Based Cross-Sectional Study. *Adv. Prev. Med.* 2015, 2015, 294902. [CrossRef] [PubMed]
55. Demissie, T.; Mekonen, Y.; Haider, J. Agroecological Comparison of Levels and Correlates of Nutritional Status of Women. *Ethiopian. J. Heal. Dev.* 2003, 17, 189–196.
56. Bitew, F.H.; Telake, D.S. Undernutrition among Women in Ethiopia: Rural-Urban Disparity; DHS Working Papers No. 77; ICF Macro: Calverton, MD, USA, 2010.
57. Naik, R.; Smith, R. Linkages between Family Planning and Nutrition. Futures Group; Health Policy Project: Washington, DC, USA, 2015.
58. Tafa, M.; Haidar, J.; Fekadu, H. Modern Family Planning Use and its Relationship with the Nutritional Status of Women of Child Bearing Age at Tena District, Arsi Zone Oromia

- Regional State, Ethiopia 2013: A Community based Case Control Study. *Gynecol. Obstet.* 2015, 5, 3.
59. World Health Organization (WHO). WHO Recommendations on Postnatal Care of the Mother and Newborn 2013; WHO: Geneva, Switzerland, 2014; pp. 1–72.
  60. Wu, X.; Li, L. Family size and maternal health: Evidence from the One-Child policy in China. *J. Popul. Econ.* 2012, 25, 1341–1364. [CrossRef]
  61. Cunningham, K.; Headey, D.; Singh, A.; Karmacharya, C.; Rana, P. Maternal and Child Nutrition in Nepal: Examining drivers of progress from the mid-1990s to 2010s. *Glob. Food Secur.* 2017, 13, 30–37. [CrossRef]
  62. World Health Organization (WHO). Improving Nutrition Outcomes with Better Water, Sanitation and Hygiene: Practical Solutions for Policies and Programmers; World Health Organization: Geneva, Switzerland, 2015.
  63. Akombi, B.J.; Agho, K.E.; Hall, J.J.; Wali, N.; Renzaho, A.M.N.; Merom, D. Stunting, wasting and underweight in Sub-Saharan Africa: A systematic review. *Int. J. Environ. Res. Public Health* 2017, 14, 863. [CrossRef] [PubMed]
  64. Hasanain, F.G.; Jamsiah, M.; Zaleha, M.I.; Tamil, A.M.; Mohammed, A.A. Association between drinking water sources and diarrhea with malnutrition among kindergarten's children in Baghdad city, Iraq. *Malays. J. Public Health Med.* 2012, 12, 45–48.
  65. Workicho, A.; Belachew, T.; Feyissa, G.T.; Wondafrash, B.; Lachat, C.; Verstraeten, R.; Kolsteren, P. Household dietary diversity and Animal Source Food consumption in Ethiopia: Evidence from the 2011 Welfare Monitoring Survey. *BMC Public Health* 2016, 16, 1192. [CrossRef] [PubMed]

66. Mosites, E.M.; Rabinowitz, P.M.; Thumbi, S.M.; Montgomery, J.M. The Relationship between Livestock Ownership and Child Stunting in Three Countries in Eastern Africa Using National Survey Data. *PLoS ONE* 2015, 10, e0136686. [CrossRef] [PubMed]
67. Betebo, B.; Ejajo, T.; Alemseged, F.; Massa, D. Household Food Insecurity and Its Association with Nutritional Status of Children 6—59 Months of Age in East Badawacho District, South Ethiopia. *J. Environ. Public Health* 2017, 2017, 6373595. [CrossRef] [PubMed]
68. Regassa, N.; Stoecker, B.J. Household food insecurity and hunger among households in Sidama district, southern Ethiopia. *Public Health Nutr.* 2012, 157, 1276–1283. [CrossRef] [PubMed]
69. Essential Nutrition Action (ENA). Key Behaviours for Optimal Breastfeeding, Complementary Feeding and Maternal Nutrition at Critical Stages in the Life Cycle of Women and Children; ENA Counselor's guide—LINKAGES Project; World Health Organization: Geneva, Switzerland, 2008.
70. Nigatu, M.; Gebrehiwot, T.T.; Gemed, D.H. Household Food Insecurity, Low Dietary Diversity, and Early Marriage Were Predictors for Undernutrition among Pregnant Women Residing in Gambella, Ethiopia. *Adv. Public Health* 2018, 2018, 1350195. [CrossRef] [PubMed]
71. Mekuria, G.; Wubneh, Y.; Tewabe, T. Household dietary diversity and associated factors among residents of finote selam town, north west Ethiopia: A cross sectional study. *BMC Nutr.* 2017, 3, 28. [CrossRef]

72. Schwei, R.J.; Tesfay, H.; Asfaw, F.; Jogo, W.; Busse, H. Household dietary diversity, vitamin A consumption and food security in rural Tigray, Ethiopia. *Public Health Nutr.* 2017, 20, 1540–1547. [CrossRef] [PubMed]
73. Gebremedhin, S.; Enquselassie, F.; Umeta, M. Prevalence of prenatal zinc deficiency and its association with socio-demographic, dietary and health care related factors in Rural Sidama, Southern Ethiopia: A cross-sectional study. *BMC Public Health* 2011, 11, 898. [CrossRef] [PubMed]
74. Ethiopian Public Health Institute (EPHI). Ethiopian National Food Consumption Survey; EPHI: Addis Ababa, Ethiopia, 2013.
75. Schlemmer, U.; Frølich, W.; Prieto, R.M.; Grases, F. Phytate in foods and significance for humans: Food sources, intake, processing, bioavailability, protective role and analysis. *Mol. Nutr. Food Res.* 2009, 53, S330–S375. [CrossRef] [PubMed]
76. Amagloh, F.K.; Atuna, R.A.; Richard McBride, R.; Edward Ewing Carey, E.E.; Christides, T. Nutrient and Total Polyphenol Contents of Dark Green Leafy Vegetables, and Estimation of Their Iron Bioaccessibility Using the In-Vitro Digestion/Caco-2 Cell Model. *Foods* 2017, 6, 54. [CrossRef] [PubMed]
77. Demissie, T.; Ali, A.; Zerfu, D. Availability and consumption of fruits and vegetables in nine regions of Ethiopia with special emphasis to vitamin A deficiency. *Ethiop. J. Health Dev.* 2009, 23, 216–222. [CrossRef]
78. Tewodros, D. Beef Cattle Production System and Opportunities for Market Orientation in Borena Zone, Southern Ethiopia. Master's Thesis, Haromaya University, Dire Dawa, Ethiopia, 2008.

79. Avery, A. Red Meat and Poultry Production and Consumption in Ethiopia and Distribution in Addis Ababa; Borlaug-Ruan World Food Prize Intern, International Livestock Research Institute: Addis Ababa, Ethiopia, 2004.



# **Chapter 3**

## **Paper II**

### **3. Feeding Practices and Undernutrition in 6–23-Month-Old Children of Orthodox Christian Mothers in Rural Tigray, Ethiopia: Longitudinal Study**

Beruk Berhanu Desalegn<sup>1,2,\*</sup>, Christine Lambert<sup>2</sup>, Simon Riedel<sup>2</sup>, Tegene Negese<sup>1</sup> and Hans Konrad Biesalski<sup>2</sup>

<sup>1</sup>College of Agriculture, Hawassa University, Postal code: 05, Hawassa, Ethiopia; tegeengss38@gmail.com

<sup>2</sup>Institute of Biological Chemistry and Nutrition, University of Hohenheim, Garbenstr. 30, 70593 Stuttgart, Germany; christine.lambert@uni-hohenheim.de (C.L.); simon.riedel@uni-hohenheim.de (S.R.); biesal@uni-hohenheim.de (H.K.B.)

\* Correspondence: berhanuberuk@gmail.com Tel.: +251-941-048-918

**A version of this chapter has been published as**

Desalegn, B., Lambert, C., Riedel, S., Negese, T., Biesalski, H.K. (2019). Feeding Practices and Undernutrition in 6–23-Month-Old Children of Orthodox Christian Mothers in Rural Tigray, Ethiopia: Longitudinal Study. *Nutrients*, 11(138). doi:10.3390/nu11010138.

# **Feeding Practices and Undernutrition in 6–23-Month-Old Children of Orthodox Christian Mothers in Rural Tigray, Ethiopia: Longitudinal Study**

## **Abstract**

Fasting period and fasting status affect the feeding practices and nutritional status of Ethiopian Orthodox mothers. Even if children are exempted from fasting, some mothers do not prepare their food from animal sources as it could contaminate utensils for cooking family foods. Therefore, the objective of this study was to assess feeding practices and undernutrition in 6–23-months old children whose mothers are Ethiopian Orthodox religion followers during lent fasting and non-fasting periods in rural Tigray, Northern Ethiopia, and to identify associated factors. A community-based longitudinal study was carried out in Ethiopian Orthodox lent fasting and non-fasting periods. Using a multi-stage systematic random sampling technique, 567 and 505 children aged 6–23 months old participated in the fasting and non-fasting assessments, respectively. Statistical analyses were done using logistic regression, an independent sample t-test, Wilcoxon signed-rank (WSRT) and McNemar's tests. The prevalences of stunting, underweight and wasting were 31.6–33.7%, 11.7–15.7% and 4.4–4.8%, respectively. The weight-for-length (WLZ) and length-for-age (LAZ) values for children of fasting mothers were significantly lower ( $p < 0.05$ ) compared to those of non-fasting mothers. Likewise, the median weight-for-age (WAZ) and diet diversity score (DDS) of children of fasting mothers were also significantly higher in non-fasting than in fasting periods. A small proportion of children (2.3–6.7%) met the minimum acceptable diet (MAD) in the study population, but these measures were significantly increased ( $p < 0.001$ ) in the children of non-fasting mothers. Mother's fasting during lactation period of the indexed child was amongst the independent factors common in child stunting,

underweight and wasting. Nutritional status and feeding practices of 6–23-month-old children are affected by maternal fasting during the fasting period. Therefore, without involvement of religious institutions in the existing nutritional activities, reduction of undernutrition would not be successful and sustainable.

Keywords: stunting; wasting; undernutrition; minimum acceptable diet; minimum diet diversity; Tigray; Orthodox fasting; Ethiopia

### **3.1. Introduction**

Despite the efforts undertaken globally, little has been achieved in reducing undernutrition with a sizeable gap from the global targets and goals still to fill. Worldwide, the number of stunted and wasted children under five years of age were about 154.8 and 52 million, respectively, in 2016, while more than 40 million children are overweight [1]. About 38%, 27% and 24% of stunted, wasted and overweight children under five years of age, respectively, were from Africa. Even worse, the number of stunted and overweight children increased by 9 and 9.8 million, respectively, between 2000 and 2016 [1]. The long-term impact of malnutrition on people's lives, most notably in health, education, and productivity, highly affects the human capital of a country. For example, an estimated loss of 4.7 billion US\$ (equivalent to 16.5% of the national GDP) was recorded in 2009 due to child undernutrition in Ethiopia [2]. Therefore, the government of Ethiopia has been following different approaches to reduce maternal and childhood undernutrition significantly. Recently, the “Seqota” Declaration [2015–2030] was launched with the goal of eliminating all forms of malnutrition among children under age 2 by 2030 [3]. However, this cannot be achieved unless the socio-cultural and religious issues related with

feeding practices are deeply investigated and addressed in Ethiopia, where more than 85 ethnic groups with diversified religious practices exist. As these practices have been passed from one generation to the next generation, they might have resulted in an intergenerational cycle of malnutrition. This could be worse in physiologically nutrient needy groups like pregnant and lactating mothers, and young and growing children.

Previous studies in Ethiopia and other parts of the world showed that maternal nutritional status and feeding practices are associated with growth indices of their children [4–8]. A more recent study by Desalegn and colleagues showed that maternal Body Mass Index (BMI) and dietary pattern of lactating mothers were negatively affected by maternal fasting status, and during Ethiopian Orthodox lent fasting period regardless of fasting status in rural Ethiopia [9]. Even if children are exempted from fasting, some mothers said that they were not happy to prepare food for them from animal sources as it could contaminate utensils for cooking family food [10]. Furthermore, a study on 6–23-month-old children of Ethiopian Orthodox mothers from the Gojam district also showed that mothers/caregivers who did not feed a diet containing animal products to their children due to fear of utensil contamination for family food preparation were less likely to feed the recommended dietary diversity. Economic limitations were a minor reason [11]. However, to the best of our knowledge, the effect of maternal fasting status and feeding practices on the nutritional status of the children and how these differ between fasting and non-fasting periods are not known. Therefore, the purpose of this study was to assess and compare the feeding practices and nutritional status of 6–23-month-old children of fasting and non-fasting lactating mothers during Ethiopian Orthodox lent fasting and non-fasting periods in rural Tigray of Ethiopia, where the majority are Orthodox religion followers.

## **3.2. Materials and Methods**

### **3.2.1. Study Area, Design, Participants and Sampling**

A longitudinal community-based survey was conducted in lent fasting (15 February–15 April 2017) and non-fasting (1–30 May 2017) periods in rural Genta Afeshum woreda (the third-level administrative division in Ethiopia), in rural Tigray, Ethiopia. The district is one of the hot spot areas of food insecurity and has a total population of 99,112 and 19 health posts to serve the community. Almost all people in the woreda belong to the ethnic group of Tigray and are followers of Ethiopian Orthodox Christianity.

For this study, the sample size was calculated using a single population proportion formula and considering the prevalence of stunting (57.1%) elsewhere in Tigray region [12], 95% of confidence interval for true prevalence and a relative precision (d) of 5%. The total population of children aged between 6–23 months old in the district was 4906, so that the finite source population size correction formula was used. Additionally, a 1.5 design effect and 10% non-response rate was used in the calculated sample size so as to get the final sample size of 575.

To obtain representative samples, multi-stage systematic random sampling was used. First, of the three woredas where the Deutsche Gesellschaft für Internationale Zusammenarbeit (GIZ, Bonn, Germany) are implementing a nutrition sensitive agriculture (NSA) project in Ethiopia, Genta Afeshum was randomly selected. Then, seven kebeles (smallest administrative unit in Ethiopia) were selected randomly out of the twenty kebeles residing in the woreda. Children who were aged between 6–23 months, apparently healthy and breastfeeding during the study periods, were included, whereas children who were twins and whose mothers were not permanent residents of the study area during the study period were excluded from the study. Based on these, the list of

children was prepared by the health extension workers in the selected kebeles, and finally the samples were selected using systematic random sampling methods.

### **3.2.2. Data Collection**

Trained and experienced data collectors who are fluent in Tigrigna, Amharic and English languages were recruited. First, the questionnaire was prepared by the principal investigator, considering the information needed for the study, translated to the local language Tigrigna by a professional translator and pre-tested for its appropriateness. Using this, information on socio-demographic and economic characteristics, maternal and child characteristics, water, sanitation and health (WASH) and feeding practices was collected. The household food insecurity information was also collected using a household food insecurity access scale (HFIAS) standard questionnaire developed by Food and Nutrition Technical Assistance Project (FANTA) [13]. The weight of children was measured using a calibrated portable digital scale (Seca 770, Hanover, Germany) working with powered battery and measured to the nearest 0.1 kg. Likewise, the length of children was measured in a lying position with a wooden board to the nearest 0.1 cm. During weight and length measurements, the mothers were advised to remove their child's clothes and shoes until the child had only light clothes to minimize the weight and to get actual length, respectively. Duplicate measurements (length and weight) were carried out following standard procedures. Minimum diet diversity (MDD), minimum meal frequency (MMF) and minimum acceptable diet (MAD) of the children were computed from the two 24-h recall data points collected preceding the survey.

Ethical approval was obtained from the institutional review board of the College of Health Sciences at Hawassa University and Tigray region Health Bureau in Ethiopia and Ethics

Commission, Landesärztekammer Baden-Württemberg, Germany. Permission was also obtained from the Genta Afeshum woreda Health Office in the Tigray Province of Ethiopia. Additionally, the purpose of the study and the confidentiality of the information to be collected were explained to the mothers, and their agreement to participate with their indexed child in the study was documented by signing the informed consent. Each participating mother was also told that whenever the mothers felt they wanted to discontinue the participation with their child, withdrawal from the study was possible.

### **3.2.3. Data Management and Analysis**

Before submitting the data, variable coding was conducted in SPSS for window version 20 (IBM Corporation, Armonk, NY, USA). Following this, the data were entered, cleaned and analyzed. First, frequency and crosstab were conducted to check completeness of data and to present the results in descriptive statistics (frequency and percent).

In our case, the diet diversity score was calculated as the summation of the number of food groups consumed for each of the two days, and averaged. Seven food groups were created, which included: grains, roots and tubers; legumes and nuts; dairy products; flesh foods; eggs; vitamin A-rich fruits and vegetables; and other fruits and vegetables. Likewise, meal frequency was computed as sum of the number of solid, semi-solid, or soft foods eaten by the child within 24-h for each of the two days, and averaged. A child who ate an average of at least four food groups in their meals of the two days fulfilled MDD. In breastfed children, the MMF was two times if aged 6–8 months and three times for 9–23 months. If a child fulfilled both the criteria for MDD and MMF, then he/she was considered as having met MAD.



The nutritional status (stunting, underweight and wasting) was identified using length-for-age, weight-for-age, and weight-for-length Z scores and compared with WHO reference data. For this, the length, weight, age and sex data of the children were entered in WHO Anthro software version 3.2.2 (World Health Organization, Geneva, Switzerland). A child who was below -2 standard deviations (-2SD) for WAZ, WLZ or LAZ was considered as stunted, wasted or underweight, respectively.

First, potential predicting variables for the outcome variables (stunting, underweight and wasting) were assessed from previous studies and included in bivariate analysis, and those with  $p\text{-value} < 0.25$  were selected for multivariate logistic regression. Then, multi-collinearity was checked between the candidate variables separately for the three outcome variables using a variance inflation factor (VIF), and all were less than 10. Then, the candidate variables were entered in a multivariate logistic regression model with the stepwise forward Wald method. A  $p\text{-value} < 0.05$  was used to declare the variables as predictors of the outcome variables. Hosmer and Lemeshow test and C-statistics (Area under the curve (AUC)) were used for model tests and were verified. Normality of continuous data was checked using the Kolmogorov-Smirnov test. An independent sample t-test was used to identify the difference between the children of fasting and non-fasting mothers' LAZ, WLZ and WAZ score values during the lent fasting period. Non-normally distributed data were analyzed using the Wilcoxon signed-rank test, whereas dichotomous data were analyzed by McNemar's test to identify the difference between children of fasting and non-fasting mothers' sub-groups dietary patterns between fasting and non-fasting periods.

### 3.3. Results

A survey was conducted with 567 children during Ethiopian Orthodox lent fasting with a response rate of 98.6%. Of those children who participated in the lent fasting survey, 505 were again involved in the non-fasting period, with a response rate of 87.8%. The reasons for excluding some children of the non-fasting period for the follow-up study were that some were older than 23 months or migrated with their family to other areas or were absent during the 2nd round survey.

#### 3.3.1. Socio-Demographic Characteristics

In this study, almost all children belong to mothers who were from Ethiopian Orthodox Christians. Most of the mothers were house-wives, and more than half of the mothers were literate.

**Table 3.1** Socio-demographic and economic characteristics of the study participants (n = 567) in rural Tigray, Northern Ethiopia (February–June, 2017).

Characteristics (n = 567)		Number	Percent
Religion of mother	Orthodox	565	99.6
	Other	2	0.4
Mother fasting during pregnancy period of indexed child	Yes	156	27.5
Mother fasting during lactation period of indexed child	Yes	176	31.0
Mother's education	Literate	369	65.1
Mother's occupation	Housewives	449	79.2
	Farmer	80	14.1
	Other	38	6.7
Family size	≤5	218	38.4
Household food security status	Food secure	166	29.3
Household toilet presence	Yes	455	80.2

About one-third of the mothers included in the study were fasting while they were pregnant and lactating. Only about one-third of 6–23-month-old children were from food secured and small family size households (HH) although most of them were from HH that owned toilets (Table 3.1).

### **3.3.2. Nutritional Status and Consumption Pattern**

Almost all children included in this study consumed colostrum during the first few days after birth. More children were stunted (31.6–33.7%) than underweight (11.7–15.7%) and wasted (4.4–4.8%) in the study. Consumption of grains, roots and tubers, legumes and nuts, and fruits and vegetable foods was lower in children born of fasting than non-fasting mothers, both in fasting and non-fasting periods. Likewise, the proportion of 6–23-month-old children of non-fasting mothers who consumed dairy products, and food groups containing egg was higher compared to children of fasting mothers, both in fasting and non-fasting periods. Additionally, the proportion of 6–23-month-old children who consumed meat and vitamin A-rich fruits and vegetables was higher during non-fasting than fasting periods. Moreover, a lower proportion of 6–23-month-old children fulfilled the MMF, MDD and MAD standards during fasting (74.3%, 2.5% and 2.3%) than during non-fasting periods (75.0%, 6.9% and 6.7%), respectively (Table 3.2).

**Table 3.2** Basic information, anthropometric status and consumption pattern of 6–23-month-old children in lent fasting and non-fasting periods in rural Tigray, Ethiopia (February–June, 2017).

		Lent Fasting Period ( <i>n</i> = 567)			Non-Fasting Period ( <i>n</i> = 505)		
		Children of Non-Fasting Mothers	Children of Fasting Mothers	Total	Children of Non-Fasting Mothers	Children of Fasting Mothers	Total
		<i>N</i> (%)	<i>N</i> (%)	<i>N</i> (%)	<i>N</i> (%)	<i>N</i> (%)	<i>N</i> (%)
Age of child	≤12 months	168 (29.6)	71 (12.5)	239 (42.2)	103 (20.4)	45 (8.9)	148 (29.3)
	13–23 months	223 (39.3)	105 (18.5)	328 (57.8)	246 (48.7)	111 (22.0)	357 (70.7)
Sex of child	Male	215 (37.9)	104 (18.3)	319 (56.3)	190 (37.6)	93 (18.4)	283 (56.0)
	Female	176 (31.0)	72 (12.7)	248 (43.7)	159 (31.5)	63 (12.5)	222 (44.0)
Colostrum intake after birth	Yes	381 (67.2)	169 (29.8)	550 (97.0)			
Stunted child		104 (18.3)	75 (13.2)	179 (31.6)	110 (21.8)	60 (11.9)	170 (33.7)
Underweight child		42 (7.4)	47 (8.3)	89 (15.7)	27 (5.3)	32 (6.3)	59 (11.7)
Wasted child		7 (1.2)	20 (3.5)	27 (4.8)	10 (2.0)	12 (2.4)	22 (4.4)
Grains, roots and tubers consumption		357 (63.0)	164 (28.9)	521 (91.9)	339 (67.1)	147 (29.1)	486 (96.2)
Legumes and nuts consumption		257 (45.3)	112 (19.8)	369 (65.1)	250 (49.5)	108 (21.4)	358 (70.9)
Other fruits and vegetables consumption		104 (18.3)	49 (8.6)	153 (27.0)	138 (27.3)	52 (10.3)	190 (37.6)
Dairy products consumption		33 (5.8)	15 (2.6)	48 (8.5)	41 (8.1)	9 (1.8)	50 (9.9)
Flesh foods consumption		0 (0.0)	3 (0.5)	3 (0.5)	4 (0.8)	4 (0.8)	8 (1.6)
Eggs consumption		71 (12.5)	34 (6.0)	105 (18.5)	87 (17.2)	38 (7.5)	125 (24.8)
Vitamin A-rich fruits and vegetables consumption		0 (0.0)	5 (0.9)	5 (0.9)	4 (0.8)	2 (0.4)	6 (1.2)
MAD		10 (1.8)	3 (0.5)	13 (2.3)	25 (5.0)	9 (1.8)	34 (6.7)
MDD		9 (1.6)	5 (0.9)	14 (2.5)	25 (5.0)	10 (2.0)	35 (6.9)
MMF		291 (51.3)	130 (22.9)	421 (74.3)	258 (51.1)	121 (24.0)	379 (75.0)

MAD = minimum acceptable diet; MDD = minimum diet diversity; MMF = minimum meal frequency; Fasting: Abstaining from eating any animal source foods during the seven official fasting periods of the Ethiopian Orthodox Tewahedo Church with/without abstaining from any foods and water until 12:00 noon and beyond of the fasting days; Fasting mother: a mother who usually abstained from eating any animal source foods (flesh foods, eggs and dairy products) during the official fasting periods of the Ethiopian Orthodox Tewahedo Church with/without abstaining from any foods and water until 12:00 noon and beyond of the fasting days, after the baptism day (40 and 80 days for a baby boy and girl, respectively) until the breast feeding indexed child was aged 23 months; Children of fasting mothers: Those children who were born of fasting mothers; Children of non-fasting mothers: Those children who were born of mothers who were not practicing fasting.

### 3.3.3. Factors Associated with Child Undernutrition

Children between 13–23 months of age were about 1.8 times more likely to be underweight than those who were between 6–12 months (AOR = 1.79 (1.07, 3.0)). Children who were born of mothers fasting during their pregnancy and lactation periods had 2.1 and 2.6 times higher odds of being underweight than those born of mothers not fasting during their pregnancy and lactation periods (AOR = 2.12 (1.29, 3.51) and AOR = 2.57 (1.56, 4.23), respectively). Children from illiterate and farming mothers were 1.8 and 2.3 times more likely to be underweight compared to those whose mothers were literate and housewives (AOR = 1.79 (1.10, 2.90) and AOR = 2.28 (1.25, 4.16), respectively). Similarly, children between 13–23 months of age had 3.1 times higher odds of stunting than those between 6–12 months of age (AOR = 3.07 (2.04, 4.63)).

Children who did not consume colostrum after their birth were about 4.1 times as likely to be stunted compared to those who did (AOR = 4.10 (1.42, 11.82)). Additionally, children born of mothers fasting during their pregnancy and lactation periods were 1.8 times more likely to be stunted (AOR = 1.83 (1.21, 2.77)) than those children from mothers who did not fast during both periods (AOR = 1.81 (1.20, 2.72)). Furthermore, children living in households who did not own a latrine were 1.7 times more likely to be stunted compared to those owning a latrine (AOR = 1.67 (1.06, 2.64)). Children whose mothers were fasting during their lactation period and who were illiterate were 6.8 and 2.2 times more exposed to wasting when compared with the references (not fasting and literate, respectively) (Table 3.3).

**Table 3.3** Multivariate logistic regression: associated factors of malnutrition of 6–23-month-old children regarding underweight, stunting and wasting in rural Tigray, Ethiopia (N = 567).

Variables	Predicting Category	OR	95% CI	p-Value
Underweight				
Age of child	13–23 months	1.79	1.07, 3.00	0.028
Mother fasting during pregnancy period of indexed child	Fasting	2.12	1.29, 3.51	0.003
Mother fasting during lactation period of indexed child	Fasting	2.57	1.56, 4.23	0.000
Mother's education	Illiterate	1.79	1.10, 2.90	0.019
Mother's occupation	Farmer	2.28	1.25, 4.16	0.021
Hosmer-Lemeshow test: $p = 0.92$ , C-statistic: AUC = 0.73; 95% CI 0.68–0.79				
Stunting				
Age of child	13–23 months	3.07	2.04, 4.63	0.000
Child colostrum intake status	No colostrum after birth	4.10	1.42, 11.82	0.009
Mother fasting during pregnancy period of indexed child	Fasting	1.83	1.21, 2.77	0.005
Mother fasting during lactation period of indexed child	Fasting	1.81	1.20, 2.72	0.005
Toilet presence	No toilet in the household	1.67	1.06, 2.64	0.026
Hosmer-Lemeshow test: $p = 0.98$ , C-statistic: AUC = 0.70; 95% CI 0.66–0.75				
Wasting				
Mother fasting during lactation period of indexed child	Fasting	6.78	2.78, 16.40	0.000
Mother education	Illiterate	2.24	1.01, 4.98	0.047
Hosmer-Lemeshow test: $p = 0.47$ , C-statistic: AUC = 0.78; 95% CI 0.70–0.86				

AUC = Area under the curve; OR = Odd ratio; CI: Confidence interval.

### 3.3.4. Trend of Food Consumption Pattern and Nutritional Status of 6–23-Month-Old Children

Based on McNemar's test, there was no significant difference ( $p > 0.05$ ) in the proportion of children born of fasting mothers who consumed from all of the seven food groups between the fasting and non-fasting periods. Likewise, the proportion of children of fasting mothers who fulfilled the MMF, MDD, and MAD criteria were not also significantly different ( $p > 0.05$ ) between fasting and non-fasting period. However, the proportion of children of non-fasting mothers who consumed grains, roots and tubers, pulses and nuts, other fruits and vegetables, and egg food groups were significantly higher during the non-fasting period compared than the fasting period ( $p \leq 0.001$ ,  $p = 0.047$ ,  $p \leq 0.001$  and  $p = 0.017$ , respectively). Similarly, the proportion of children who fulfilled the MDD and MAD were significantly increased in the non-fasting period compared to the fasting period, for the children of non-fasting mothers' group ( $p \leq 0.001$ ) (Table 3.4).

**Table 3.4** Food consumption pattern of 6–23-month-old children of fasting and non-fasting mothers during lent fasting and non-fasting periods in rural Tigray, Northern Ethiopia.

Parameters	Children of Fasting Mothers ( <i>n</i> = 156)			Children of Non-Fasting Mothers ( <i>n</i> = 349)		
	Lent Fasting Period	Non-Fasting Period	Significance	Lent Fasting Period	Non-fasting Period	Significance
	<i>N</i> (%)	<i>N</i> (%)	<i>p</i> -Value	<i>N</i> (%)	<i>N</i> (%)	<i>p</i> -Value
Grains, roots and tubers consumption	145 (92.9)	147 (94.2)	0.815 <sup>ns</sup>	317 (90.8)	339 (97.1)	0.000 *
Pulses and nuts consumption	97 (62.2)	108 (69.2)	0.228 <sup>ns</sup>	228 (65.3)	250 (71.6)	0.047 *
Other fruits and vegetables	45 (28.8)	52 (33.3)	0.427 <sup>ns</sup>	91 (26.1)	138 (39.5)	0.000 *
Dairy products consumption	14 (9.0)	9 (5.8)	0.332 <sup>ns</sup>	31 (8.9)	41 (11.7)	0.203 <sup>ns</sup>
Flesh food consumption	3 (1.9)	4 (2.6)	1.000 <sup>ns</sup>	0 (0.0)	4 (1.1)	NA
Eggs consumption	30 (19.2)	38 (24.4)	0.302 <sup>ns</sup>	64 (18.3)	87 (24.9)	0.017 *
Vitamin A fruits and vegetables	5 (3.2)	2 (1.3)	0.453 <sup>ns</sup>	0 (0.0)	4 (1.4)	NA
MAD	3 (1.9)	9 (5.8)	0.146 <sup>ns</sup>	7 (2.0)	25 (7.5)	0.001 *
MDD	5 (3.2)	10 (6.4)	0.302 <sup>ns</sup>	6 (1.7)	25 (7.2)	0.001 *
MMF	114 (73.1)	121 (77.6)	0.349 <sup>ns</sup>	256 (73.4)	258 (73.9)	0.927 <sup>ns</sup>

Data analysis using McNemar's test, significant level at  $p < 0.05$ ; NA = the data were not appropriate for analysis; MAD: minimum acceptable diet, MDD = minimum diet diversity, MMF: minimum meal frequency; ns- not significantly different at  $p < 0.05$ ; \* = significantly different at  $p < 0.05$ .

Likewise, the WAZ, WLZ and LAZ of children of non-fasting mothers were significantly higher than those of children of the fasting sub-group during the lent fasting period ( $p \leq 0.001$ ,  $p \leq 0.001$  and  $p = 0.003$ , respectively) (Table 3.5). A Wilcoxon signed-rank test showed that except for the median of WAZ ( $p = 0.153$ ) for children of the non-fasting mothers' sub-group, the median WAZ ( $p = 0.034$ ) for the children of fasting mothers, and the DDS of both, the children of the fasting and non-fasting mothers' sub-groups were significantly increased in the non-fasting than in the fasting period ( $p \leq 0.001$ ) (Table 3.6).

**Table 3.5** Anthropometric measurements (WAZ, WLZ and LAZ) of 6–23-month-old children of fasting and non-fasting mothers in the lent fasting period in rural Tigray, Ethiopia.

Variables	Fasting Mother (N = 176)	Non-Fasting Mother (N = 391)	p-Value
	Mean (Standard Deviation)	Mean (Standard Deviation)	
WAZ	−1.25 (1.15)	0.77 (1.09)	0.000 *
WLZ	−0.58 (1.15)	−0.19 (1.06)	0.000 *
LAZ	−1.60 (1.35)	−1.24 (1.34)	0.003 *

LAZ = length for age; WLZ = weight for length; WAZ = weight for age; data analysis using independent sample t-test, significance level declared at  $p < 0.05$ ; \* = significantly different at  $p < 0.05$ .

**Table 3.6** Comparison of 6–23-month-old children’s WAZ and DDS values between lent fasting and non-fasting periods in rural Tigray, Ethiopia

Variables	Data Collection Period						
	Lent Fasting			Non-Fasting			Sign
	Mean (SD)	Median (IQR)	Range (Min, Max)	Mean (SD)	Median (IQR)	Range (Max, Min)	
Children WAZ of fasting mother	−1.25 (1.14)	−1.22 (−2.03, −0.54)	−4.33, 2.52	−1.16 (1.14)	−1.11 (−1.83, −0.46)	−4.37, 3.83	0.034 *
Children WAZ of non-fasting mother	−0.74 (1.08)	−0.79 (−1.48, −0.07)	−4.97, 2.70	−0.66 (1.02)	−0.69 (−1.32, −0.11)	−0.508, 2.32	0.153 <sup>ns</sup>
Children DDS of fasting mother	2.09 (1.02)	2.00 (1.50, 2.50)	0.00, 7.00	2.43 (0.93)	2.50 (2.00, 3.00)	0.00, 4.5)	0.000 *
Children DDS of non-fasting mother	2.04(0.88)	2.00(1.50, 2.50)	0.00, 5.00	2.50 (0.95)	2.50 (2.00, 3.00)	0.00, 7.00	0.000 *

Fasting mothers (n = 156); non-fasting mother (n = 349); data analyses using the Wilcoxon signed-rank test, significant level at  $p < 0.05$ ; WAZ = weight for age; DDS = diet diversity score; ns- not significantly different at  $p < 0.05$ ; \* = significantly different at  $p < 0.05$ .

### 3.4. Discussion

This is the first time a study has assessed feeding practices and nutritional status of 6–23-month-old children of fasting and non-fasting mothers during the Ethiopian Orthodox lent fasting and non-fasting periods in rural Tigray, Ethiopia. Socio-economic factors, which are possibly associated with stunting, underweight and wasting, were also determined.



### **3.4.1. Food Consumption Pattern of 6–23-Month-Old Children**

According to the UNICEF conceptual framework of malnutrition, inadequate dietary intake is one of the immediate causes of child malnutrition [14]. In our study, cereals, roots and tubers, and pulses and nuts are the food groups majorly prepared for the 6–23-month-old children in the rural Tigray in Ethiopia, regardless of fasting period. This result is consistent with previous studies in Ethiopia [11,15–20]. A similar finding was reported on the consumption of grains, roots and tubers for children from Vietnam, Bangladesh, Pakistan, Indian and the U.S. [21–25]. According to Demissie and his colleagues [21], consumption of fruits and vegetables is sub-optimal in the Tigray region and Ethiopia in general [26]. Supporting this evidence, we also found that only 1% of children consumed vitamin A-rich fruits and vegetables. The result is also comparable with a previous study in rural Tigray [27]. However, our finding was lower than findings of two national surveys [14.8, 27.7%] in Ethiopia and Pakistan (18.7%) [22,28,29]. This could be related to the fact that cultivation on the farm and availability in the markets are rare in the study area, probably resulting in a less diverse diet. Another explanation could be a difference in feeding practices [26]. However, the consumption of other fruits and vegetables food group was between 27.0–37.6% in the present study. This result is higher than national figures, which are 3.3% and 10.1% [28,29].

Animal source foods are both energy and nutrient dense, and an excellent source of high quality and easily digestible protein, in addition to being an efficient and easily absorbable source of micronutrients like iron, zinc, vitamin A, vitamin B12, riboflavin and calcium [30]. In the present study, the consumption of a diet composed of flesh foods was 0.5% and 1.6% in fasting and non-fasting periods, respectively. These results are comparable with a previous study during the Ethiopian Orthodox fasting period in the Dejen district (0.0%), and studies conducted in

Gobalafto district in North Wollo (0–5%), Wolaita district (1.6%), and findings of the Ethiopian national food consumption survey for all regions in the country (0.1–0.6%) [11,15,31,32]. However, our finding was lower than a study conducted in the Arsi-Negele district of Oromia region, in Ethiopia (11.3%) and Mongolia (4–14%) [33,34]. This might be due to the fact that the area of the later studies has animal husbandry potential and cultural and eating habit differences. In the current study, consumption of dairy products was between 8.5% and 9.9% in fasting and non-fasting periods, respectively. Almost comparable proportion of 6–23-month-old children (7.5%) of the Dejen district consumed dairy products in the fasting period in North-west Ethiopia [11]. However, the findings in the present study were lower than in studies conducted in Amhara (about 20.0%) and Oromia (52.4%) regions, and Addis Ababa City (70.4%) in Ethiopia and Pakistan (50.5%) [17,20,22,34]. An explanation for this could be that the studies were conducted in different periods, and dairy products were the second most common foods groups consumed by the Pakistan children. In the present study, the proportions of children who consumed egg in the days before the surveys of fasting and non-fasting periods were 18.5% and 24.8%. However, these were lower than in similar studies in the Amhara region, which was 98.0%, where fasting of Ethiopian Orthodox religion is routine in the fasting days of the year. This could be due to more chickens in the households included in the later study, which may result in availability of eggs for children during the fasting period and household consumption in general. The consumption of egg in our finding was higher than the national findings of the last two Ethiopian demographic and health survey (DHS) studies (8.3%, 17%) and study in Amhara region (<10.0%) in Ethiopia [15,28,29]. However, it was consistent with a study conducted in the Arsi-Negele district (22.3%) of Oromia region, in Ethiopia [34].

Minimum meal frequency is a proxy indicator to determine whether the energy requirement of a child is met and examines the frequency of meals where the children received foods other than breastmilk, considering specific age and breastfeeding status of the child [29]. In this study, three-fourths of children of fasting and non-fasting mothers had eaten the minimum number of meals in the days preceding the fasting and non-fasting surveys. This result is in line with studies conducted in the Dabat (69.3–75.0%) and Gorche (78.0%) districts of northern and southern Ethiopia, respectively [35,36]. However, our finding is higher than studies in Kemba (14.5–33.6%), Bale (44.7%), Arsi-Negele (67.3%), Wolaita (69.3%) districts, elsewhere in Tigray region (66.0%) and findings for the Tigray region (55.6%, 49.4%), and national averages (47.9%, 44.6%) in Ethiopian DHS studies of 2011 and 2016, respectively, and Pakistan (38.2%) [16,19,28,29,34,37,38]. However, it is lower than findings on children from the South Wollo district, in Ethiopia, which were 91.0–97.0% and Nepal (87.8–98.2%) [15,39].

Minimum dietary diversity is one of the indicators to assess the quality of food consumed by 6–23-month-old children. Consumption of at least four different food groups in a day is the minimum requirement [29,40]. In this study, the proportion of children who consumed foods prepared from at least four diversified food groups (MDD) was 2.5% and 6.9% in fasting and non-fasting periods, respectively. This finding indicates that children aged 6–23 months in the study area are consuming a poor quality complementary diet. A study conducted in 6–23-month-old children of Orthodox Christian mothers in Gojam indicated that 13.6% had four diversified food groups in the food they consumed preceding the survey during the fasting period [11]. Similarly, our findings are also lower than findings in children living in the Gorche (10.4%), Kemba (23.3%), Bale (28.5%), and Wollo (23.0–41.0%) districts in Ethiopia, India (38%), Bangladesh (48.4%) and Vietnam (83.2%) [15,16,25,35,41,42]. Generally, these findings

indicate that undernutrition specifically stunting and micronutrient deficiencies could be the public health problem in the area. This is supported by studies conducted in children from different parts of the world and ours [43–46].

For appropriate growth and development, infants and young children should be fed with at least four diversified food groups from the minimum number of meals (MAD) recommended according to specific to the age and breastfed status of the child [40]. In the present study, the proportion of 6–23-month-old children who met both the minimum diversity and the number of meals (MAD) preceding the surveys was 2.3–6.7% in the study population. The result is consistent with DHS 2011 and 2016 findings for the Tigray region (4.5% and 5.7%, respectively) and the national figure (4.0%) in DHS 2016 for Ethiopia [28,29]. However, it is lower than research findings conducted in three districts (17.0%) of the Tigray region, in northern Ethiopia, India (9%), Nepal (32%), Bangladesh (40%) and Sri Lanka (68%) [31,38,47]. Moreover, the proportion of 6–23-month-old children who fulfilled the criteria for MAD was higher for non-fasting mothers (2.0–7.5%) and in non-fasting period (6.7%) compared with their counterparts. This finding suggests that activities which can improve the child feeding practices should include improving maternal feeding behavior, as many scholars evidenced their association, including ours [9,21,48].

### **3.4.2. Nutritional Status of 6–23-month-Old Children**

In the present study, the prevalence of wasting, underweight and stunting in the 6–23 months old children was between 4.4–4.8%, 11.7–15.7% and 31.6–33.7%, respectively. These results are lower than the findings for the Tigray region and national figures in last two DHS studies in Ethiopia. For example, wasting, underweight, and stunting in under-five aged children were

10.3% and 9.7%, 35.1% and 28.7%, and 51.4% and 44.4% in DHS 2011 (Tigray region and national, respectively). Except for wasting, these figures further decreased to 23.0% and 23.6% for underweight, and 39.3% and 38.4% for stunting in the DHS 2016 study [28,29]. This could indicate an improvement in child nutritional status, both regionally, and at the national level.

In this study, the median WAZ of children of fasting mothers was lower compared to children of non-fasting mothers, both in fasting and non-fasting periods. The WAZ of children from non-fasting mothers' sub-group also improved during the non-fasting period. Similarly, the prevalence of underweight children from the fasting sub-group was higher than that of children from the non-fasting sub-group, in both periods. Likewise, the mean standard deviations (SD) of WAZ, WLZ and LAZ of children from the fasting mothers sub-group were lower compared to children from non-fasting mothers during the fasting period. Similarly, the proportion of wasted children was higher in the fasting sub-group compared to those children from non-fasting mothers during the fasting period. Conversely, relatively more children from non-fasting mothers were stunted than from fasting mothers. This could be due to the higher number of older children in the non-fasting group. Similar findings were observed in studies conducted in the Tigray region, northern Ethiopia [4]. Multivariate logistic regression showed that children of mothers who fasted during the pregnancy and lactation period were more likely to be undernourished (stunting, underweight, and wasting) than those children of mothers who were not fasting in the same periods. These indicate that, maternal fasting during pregnancy, and lactation not only affect the nutritional status of the mother themselves [9], but also their fetus/breastfed child. To increase the awareness on the effect of fasting practices on the undernutrition, the Ethiopian Orthodox Church (EOTC) with the United States Agency for International Development (USAID) multi-sectoral nutrition project, Empowering New Generations to Improve Nutrition

and Economic opportunities (ENGINE) held a consultative meeting with religious leaders and church scholars to clarify fasting practices for children, pregnant and lactating women in line with religious guidelines. This was done after a sermon guide was developed in 2016, based on the church teachings and outlining the laws and regulations related with fasting and nutrition. According to the joint press release by EOTC and USAID/ENGINE, the sermon guide highlighted the exemption of mothers and their babies from fasting during the first 1000 days [49]. The sermon guide also emphasized the cooking utensils to be used for complementary food preparation during fasting seasons, as many mothers were not happy to prepare complementary food from the animal sources during fasting seasons due to fear of contamination of family food [10]. Therefore, strengthening the activities started by the Church and the involvement of religious institutions and leaders in the existing national multi-sectoral approach activities would accelerate and sustain the reduction of nutrition-related problems in the country.

Previous studies in Africa and Southeast Asian countries demonstrated that children in the age between 13–23 months were more associated with underweight and stunting compared to those who were aged 6–12 months [28,29,50–61]. Similar findings were also revealed in our study. This could be related to feeding practices, contamination and poor quality complementary food, which is mainly cereal based [4]. Similarly, maternal illiteracy was found to be a predictor variable for underweight and wasting. The former result is consistent with previous studies in Ethiopia and other low and middle income countries [28,29,62–65]. Similarly, findings in Ethiopian DHS also showed that childhood wasting was associated with maternal illiteracy [28,29]. This indicates that educating mothers will be an important activity for improving the nutritional status of children in a given community. Likewise, compared to children whose mothers were housewives, children of farmer mothers were more exposed to being underweight.

This might be related to less time allocation for child care due to a high work load and absence from the home. Farm work also expends more energy, and mothers may be more frequently exhausted, which could result in poor appetite and might affect the quality and quantity of breast milk. Moreover, children who did not take the first milk (colostrum) after their birth were more stunted than those who did. This could be related to infection emanating from pre-lacteal foods, which is a predictor for child stunting in North-west Ethiopia [58]. Therefore, activities which improve the colostrum intake (e.g.,- antenatal care) should be done in the study area [66]. Furthermore, children who were living in households without toilets were more exposed to stunting compared to those from households owning toilets. This could be related to poverty in general, a well-known factor for malnutrition. Additionally, contamination of the soil due to open defecation increases the risk for frequent episodes of childhood diarrhea, which has a negative impact on the nutritional status of sub-Saharan African children [50].

However, this study has the following limitations: This study considered only the lent fasting period, among the seven official fasting periods in Ethiopian Orthodox religion. It also included only children aged between 6–23 months, out of those children below seven years of age, who are exempted from the official fasting in the religion.

### **3.5. Conclusions**

This study showed that there is high prevalence of stunting (31.6–33.7%), underweight (11.7–15.7%), and wasting (4.4–4.8%) in the study area; however, these values are lower than those reported earlier in the Tigray region (39.3%, 23.0% and 11.1%), and of national figures (38.4%, 23.6% and 9.9%). These are indications of improvement of child nutrition in rural Tigray. However, the prevalence's are still higher, both in children of fasting mothers and the study

population especially during Ethiopian Orthodox lent fasting period. Thus, religious fasting should not go against the existing nutritional intervention modalities. Mother's fasting during the lactation period of an indexed child was the common predictor for child stunting, underweight and wasting. Maternal illiteracy and age of children and maternal fasting during pregnancy of indexed children were also common variables affecting combined effects of underweight and wasting, and underweight and stunting, whereas, the type of job of mothers and lack of colostrum feeding of children after birth were associated with underweight and absence of a toilet in the household with the stunting of children. Hence, improving maternal education and reducing the work burden in farming activities or promotion and support of mothers in off-farm income generation nearby their households are recommended. Improving household latrine coverage and breastfeeding of colostrum should be encouraged and strengthened. 'Grains, roots, and tubers' was the most frequently eaten food group by 6–23-month-old children, followed by 'legumes and nuts' and then 'other fruits and vegetables'. Therefore, to increase nutrients and improve the palatability of grains and legumes, traditional food processing techniques like dehulling/dehusking, soaking and germination, and slightly roasting are crucial. Additionally, the proportion of children who consumed dairy products, flesh foods and eggs who met the MDD was small. The average DDS of the children was also low, being worst in children of fasting mothers especially during the lent fasting period. Thus, regardless of the fasting period, diet diversity should be properly addressed in preparation of complementary foods by incorporating animal products [dairy products; flesh foods; eggs], vitamin A-rich fruits and vegetables. In general, nutritional status is affected by maternal fasting status during pregnancy and lactation periods, and during the lent fasting period in rural Tigray, Ethiopia. Therefore, without strengthening and involvement of religious institutions and leaders in the existing nutrition



improvement activity by the Church and in the National Nutrition Program (NNP) of, in particular nursing mothers and growing children, reduction of undernutrition would not be successful and sustainable. As this study is focused on the food consumption and growth index pattern of 6–23-month-old children, further studies on physiological consequences of levels of micronutrients or bio-markers are recommended.

Author Contributions: B.B.D., C.L., S.R. and H.K.B. involved in research design; B.B.D. conducted the research; B.B.D. analyzed the data; B.B.D. wrote the paper; B.B.D., C.L., S.R., T.N., and H.K.B. edited draft manuscript; the final content was the primary responsibility of B.B.D. All authors read and approved the final manuscript.

Funding: This study was financially supported by German Federal Ministry for Economic Cooperation and Development (BMZ).

Acknowledgments: This study is part of the PhD work of the first author and was conducted with the financial support from the German Federal Ministry for Economic Cooperation and Development (BMZ). The authors would also like to thank the data collectors of this study, and Habteab and Muez, the staff of GIZ Nutrition Sensitive Agriculture project in the Tigray region and Gebretsadekane, Genta Afeshum Woreda Health Office Nutrition coordinator, in Ethiopia. They would also like to thank Andrea Flaccus for English language editing.

Conflicts of Interest: No conflict of interest among authors.

### **3.6. References**

1. UNICEF. Levels and Trend in Child Malnutrition: UNICEF/WHO/World Bank Group Joint Estimates, Key Findings of the 2017 Edition. 2017, p. 16. Available online: [http://www.who.int/nutgrowthdb/jme\\_brochure2017.pdf](http://www.who.int/nutgrowthdb/jme_brochure2017.pdf) [accessed on 15 May 2018].

2. Ethiopian Public Health Institute [EPHI]. The Cost of Hunger in Ethiopia; EPHI: Addis Ababa, Ethiopia, 2013.
3. Federal Democratic Republic of Ethiopia [FDRE]. Seqota Declaration Implementation Plan [2016–2030]: Summary Program Approach Document; FDRE: Addis Ababa, Ethiopia, 2016.
4. Mulugeta, A.; Hagos, F.; Kruseman, G.; Linderhof, B.; Stoecker, B.; Abraha, Z.; Yohannes, M.; Samuel, G.G. Factors Contributing to Child Malnutrition in Tigray, Northern Ethiopia. *East Afr. Med. J.* 2010, 87, 248–254. [PubMed]
5. Felisbino-mendes, M.; Villamor, E.; Velasquez-melendez, G. Association of maternal and child nutritional status in Brazil: A population based cross-sectional study. *PLoS ONE* 2014, 9, e87486. [CrossRef] [PubMed]
6. Negash, C.; Whiting, S.J.; Henry, C.J.; Belachew, T.; Hailemariam, T.G. Association between maternal and child nutritional status in Hula, rural southern Ethiopia: A cross sectional study. *PLoS ONE* 2015, 10, e0142301. [CrossRef] [PubMed]
7. Silva, D.E.; Valente, A.; Borges, A.; Almeida, F.; Neves, E.; Afonso, C. Relationship between the mothers' nutritional status with that of a child population from São Tomé Príncipe, “Africa”. *Rev. Bras. Saúde Matern. Infant. Recif.* 2017, 17, 327–335. [CrossRef]
8. Zhang, J.; Kang, H.; Wang, L.; Zhao, X.; Liu, R.; He, L. A Prevalence Analysis to Assess the Relationship between Maternal and Children's Nutritional Status in North-Western Rural China Clinics in Mother and Child Health. *Clin. Mother Child Health* 2016, 13, 100242. [CrossRef]
9. Desalegn, B.B.; Lambert, C.; Riedel, S.; Negese, T.; Biesalski, H.K. Ethiopian Orthodox Fasting and Lactating Mothers: Longitudinal Study on Dietary Pattern and Nutritional Status

- in Rural Tigray Tigray, Ethiopia. *Int. J. Environ. Res. Public Health* 2018, 15, 1767. [CrossRef] [PubMed]
10. Alive and Thrive. IYCF Practices, Beliefs, and Influences in Tigray Region, Ethiopia; Alive and Thrive: Addis Ababa, Ethiopia, 2010.
  11. Kumera, G.; Tsedal, E.; Ayana, M. Dietary diversity and associated factors among children of Orthodox Christian mothers/caregivers during the fasting season in Dejen district, north west Ethiopia. *Nutr. Metab.* 2018, 15, 16. [CrossRef]
  12. Tamiru, M.W.; Tolessa, B.E.; Abera, S.F. Under Nutrition and Associated Factors Among Under-Five Age Children of Kunama Ethnic Groups in Tahtay Adiyabo Woreda, Tigray Regional State, Ethiopia: Community Based Study. *Int. J. Nutr. Food Sci.* 2015, 4, 277–288. [CrossRef]
  13. Coates, J.F.; Swindale, A.; Bilinsky, P. Household Food Insecurity Access Scale [HFIAS] for Measurement of Household Food Access: Indicator Guide [v. 3]; Food and Nutrition Technical Assistance Project; Academy for Educational Development: Washington, DC, USA, 2007; p. 36.
  14. UNICEF. The Status of the World's Children; Oxford University Press: Geneva, Switzerland, 1998; p. 135.
  15. Baye, K.; Guyot, J.P.; Icard-Vernière, C.; Mouquet-Rivier, C. Nutrient intakes from complementary foods consumed by young children [aged 12–23 months] from North Wollo, northern Ethiopia: The need for agro-ecologically adapted interventions. *Public Health Nutr.* 2013, 16, 1741–1750. [CrossRef]

16. Gatahun, E.A. Dietary Diversity Feeding Practice and Determinants among Children Aged 6-23 Months in Kemba Woreda, Southern Ethiopia Implication for Public Health Intervention. *J. Nutr. Food Sci.* 2015. [CrossRef]
17. Gebremedhin, S.; Baye, K.; Bekele, T.; Tharaney, M.; Asrat, Y.; Abebe, Y.; Reta, N. Predictors of dietary diversity in children ages 6 to 23 months in largely food-insecure area of South Wollo, Ethiopia. *Nutrition* 2017, 33, 163–168. [CrossRef] [PubMed]
18. Hirvonen, K.; Hoddinott, J.; Minten, B.; Stifel, D. Children's Diets, Nutrition Knowledge, and Access to Markets. *World Dev.* 2017, 95, 303–315. [CrossRef]
19. Mekonnen, T.C.; Workie, S.B.; Yimer, T.M.; Mersha, W.F. Meal frequency and dietary diversity feeding practices among children 6–23 months of age in Wolaita Sodo town, Southern Ethiopia. *J. Health Popul. Nutr.* 2017, 36, 18. [CrossRef] [PubMed]
20. Solomon, D.; Aderaw, Z.; Tegegne, T.K. Minimum dietary diversity and associated factors among children aged 6–23 months in Addis Ababa, Ethiopia. *Int. J. Equit. Health* 2017, 16, 1–9. [CrossRef] [PubMed]
21. Nguyen, P.H.; Avula, R.; Ruel, M.T.; Saha, K.K.; Ali, D.; Tran, L.M.; Frongillo, E.A.; Menon, P.; Rawat, R. Maternal and Child Dietary Diversity Are Associated in Bangladesh, Vietnam, and Ethiopia. *J. Nutr.* 2013, 143, 1176–1183. [CrossRef] [PubMed]
22. Iqbal, S.; Zakar, R.; Zakar, M.Z.; Fischer, F. Factors associated with infants' and young children's [6–23 months] dietary diversity in Pakistan: Evidence from the demographic and health survey 2012–2013. *Nutr. J.* 2017, 16, 78. [CrossRef] [PubMed]
23. Hamner, H.C.; Perrine, C.G.; Gupta, P.M.; Herrick, K.A.; Cogswell, M.E. Food consumption patterns among U.S. children from birth to 23 months of age, 2009–2014. *Nutrients* 2017, 9, 942. [CrossRef] [PubMed]

24. Gladys, C. Dietary Assessment and Nutritional Status of Children [6 23 Months] Consuming Local Food Recipes in Vihiga County, Western Kenya; University of Nairobi Digital Repository: Nairobi, Kenya, 2017.
25. Divya, N.; Rajanish, K.V.; Malavika, J.; Sharma, A. The study of dietary diversity score in children between 6 months to 23 months: A hospital based study. *Int. J. Contemp. Pediatrics* 2018, 5, 1053–1057.
26. Demissie, T.; Ali, A.; Zerfu, D. Availability and consumption of fruits and vegetables in nine regions of Ethiopia with special emphasis to vitamin A deficiency. *Ethiop. J. Health Dev.* 2009, 23, 216–222. [CrossRef]
27. Schwei, R.J.; Tesfay, H.; Asfaw, F.; Jogo, W.; Busse, H. Household dietary diversity, vitamin A consumption and food security in rural Tigray, Ethiopia. *Public Health Nutr.* 2017, 20, 1540–1547. [CrossRef]
28. Central Statistical Agency [CSA]. Ethiopia Demographic and Health Survey 2011; Central Statistical Agency: Addis Ababa, Ethiopia; ICF International Calverton: Calverton, MD, USA, 2012; p. 452.
29. Central Statistical Agency [CSA]. Ethiopia Demographic and Health Survey 2016; Central Statistical Agency: Addis Ababa, Ethiopia; ICF International Calverton: Calverton, MD, USA, 2017; p. 551.
30. Neumann, C.; Harris, D.M.; Rogers, L.M. Contribution of animal source foods in improving diet quality and function in children in the developing world. *Nutr. Res.* 2002, 22, 193–220. [CrossRef]
31. Ethiopian Public Health Institute [EPHI]. Ethiopia National Food Consumption Survey; EPHI: Addis Ababa, Ethiopia, 2013.

32. Gamo, S.G. Dietary Diversity and Associated Factors among Infants and Young Children in Wolaita Zone, Southern Ethiopia. *Sci. J. Clin. Med.* 2017, 6, 53. [CrossRef]
33. Lander, R.; Enkhjargal, T.S.; Batjargal, J.; Bolormaa, N.; Enkhmyagmar, D.; Tserendolgor, U.; Tungalag, S.; Bailey, K.; Gibson, R.S. Poor dietary quality of complementary foods is associated with multiple micronutrient deficiencies during early childhood in Mongolia. *Public Health Nutr.* 2010, 13, 1304–1313. [CrossRef] [PubMed]
34. Kassa, T.; Meshesha, B.; Haji, Y.; Ebrahim, J. Appropriate complementary feeding practices and associated factors among mothers of children age 6–23 months in Southern Ethiopia, 2015. *BMC Pediatr.* 2016, 16, 1–10. [CrossRef] [PubMed]
35. Dangura, D.; Gebremedhin, S. Dietary diversity and associated factors among children 6–23 months of age in Gorche district, Southern Ethiopia: Cross-sectional study. *BMC Pediatr.* 2017, 17, 1–7. [CrossRef] [PubMed]
36. Belew, A.K.; Ali, B.M.; Abebe, Z.; Dachew, B.A. Dietary diversity and meal frequency among infant and young children: A community based study. *Ital. J. Pediatr.* 2017, 43, 6–15. [CrossRef] [PubMed]
37. Khan, G.N.; Ariff, S.; Khan, U.; Habib, A.; Umer, M.; Suhag, Z.; Hussain, I.; Bhatti, Z.; Ullah, A.; Turab, A.; et al. Determinants of infant and young child feeding practices by mothers in two rural districts of Sindh, Pakistan: A cross-sectional survey. *Int. Breastfeed J.* 2017, 12, 1–8. [CrossRef] [PubMed]
38. Ludwig, C.; Bau, A. Nutrition Baseline Survey—Ethiopia-for the Global Programme: Food and Nutrition Security and Enhanced Resilience; Food and Nutrition Security and Enhanced Resilience: Addis Ababa, Ethiopia, 2007.

39. Morseth, M.S.; Torheim, L.E.; Gebremariam, M.K.; Chandyo, R.K.; Ulak, M.; Shrestha, S.K.; Shrestha, B.; Henjum, S. Tracking of infant and young child feeding practices among 9- to 24-month-old children in Nepal: The MAL-ED Birth Cohort Study. *Public Health Nutr.* 2018, 21, 355–364. [CrossRef] [PubMed]
40. WHO. Indicators for Assessing Infant and Young Child Feeding Practices. Part 3 Country Profiles; WHO: Geneva, Switzerland, 2010; pp. 1–47.
41. Ali, D.; Saha, K.K.; Nguyen, P.H.; Diressie, M.T.; Ruel, M.T.; Menon, P.; Rawat, R. Household Food Insecurity Is Associated with Higher Child Undernutrition in Bangladesh, Ethiopia, and Vietnam, but the Effect Is Not Mediated by Child Dietary Diversity. *J. Nutr.* 2013, 143, 2015–2021. [CrossRef]
42. Tegegne, M.; Sileshi, S.; Benti, T.; Teshome, M.; Woldie, H. Factors associated with minimal meal frequency and dietary diversity practices among infants and young children in the predominantly agrarian society of Bale zone, Southeast Ethiopia: A community based cross sectional study. *Arch. Public Health* 2017, 75, 1–11.
43. Arimond, M.; Ruel, M.T. Dietary diversity is associated with child nutritional status: Evidence from 11 demographic and health surveys. *J. Nutr.* 2004, 134, 2579–2585. [CrossRef] [PubMed]
44. Ogechi, U.P.; Chilezie, O.V. Assessment of Dietary Diversity Score, Nutritional Status and Socio-demographic Characteristics of Under-5 Children in Some Rural Areas of Imo State, Nigeria. *Mal. J. Nutr.* 2017, 23, 425–435.
45. Steyn, N.P.; Nell, J.H.; Nantel, G.; Kennedy, G.; Labadarios, D. Food variety and dietary diversity scores in children: Are they good indicators of dietary adequacy? *Public Health Nutr.* 2006, 9, 644–650. [CrossRef] [PubMed]

46. Zhao, W.; Yu, K.; Tan, S.; Zheng, Y.; Zhao, A.; Wang, P.; Zhang, Y. Dietary diversity scores: An indicator of micronutrient inadequacy instead of obesity for Chinese children. *BMC Public Health* 2017, 17, 440. [CrossRef] [PubMed]
47. Senarath, U.; Godakandage, S.S.P.; Jayawickrama, H.; Siriwardena, I.; Dibley, M.J. Determinants of inappropriate complementary feeding practices in young children in Sri Lanka: Secondary data analysis of Demographic and Health Survey 2006–2007. *Matern. Child Nutr.* 2012, 8 [Suppl. 1], 60–77. [CrossRef] [PubMed]
48. Amugsi, D.A.; Mittelmark, M.B.; Oduro, A. Association between Maternal and Child Dietary Diversity: An Analysis of the Ghana Demographic and Health Survey. *PLoS ONE* 2015, 10, e0136748. [CrossRef] [PubMed]
49. EOTC/USAID. Patriarch of the Ethiopian Orthodox Tewahedo Church Endorses Nutrition Sermon Guide. Encouraging Better Nutrition for Mothers and Children; USAID: Addis Ababa, Ethiopia, 2019; pp. 1–2. Available online: <https://2012-2017.usaid.gov/ethiopia/news-information/press-releases/patriarchethiopian-orthodox-tewahedo-church-endorses> [accessed on 29 March 2018].
50. Akombi, B.J.; Agho, K.E.; Hall, J.J.; Wali, N.; Renzaho, A.M.N.; Merom, D. Stunting, wasting and underweight in Sub-Saharan Africa: A systematic review. *Int. J. Environ. Res. Public Health* 2017, 14, 863. [CrossRef]
51. Alemayehu, M.; Tinsae, F.; Hailelassie, K.; Seid, O.; Gebregziabher, G.; Yebyo, H. Undernutrition status and associated factors in under-5 children, in Tigray, Northern Ethiopia. *Nutrition* 2015, 31, 964–970. [CrossRef]



52. Altare, C.; Delbiso, T.D.; Mutwiri, G.M.; Kopplow, R.; Guha-Sapir, D. Factors associated with stunting among pre-school children in southern highlands of Tanzania. *J. Trop. Pediatr.* 2016, 62, 390–408. [CrossRef]
53. Birhanu, A.; Mekonen, S.; Atenafu, A. Stunting and associated factors among children aged 6-59 months in Lasta woreda, north east Ethiopia, 2015: A community based cross sectional study design. *J. Fam. Med.* 2017, 4, 1112.
54. Derso, T.; Tariku, A.; Biks, G.A.; Wassie, M.M. Stunting, wasting and associated factors among children aged 6–24 months in Dabat health and demographic surveillance system site: A community based cross-sectional study in Ethiopia. *BMC Pediatr.* 2017, 17, 1–9. [CrossRef] [PubMed]
55. Desalegn, B.B.; Kinfe, E.; Fikre, K.; Bosha, T. Stunting and It's Associated Factors in Under Five Years Old Children: The Case of Hawassa University Technology Villages, Southern Ethiopia. *IOSR. J. Environ. Sci. Toxicol. Food Technol.* 2016, 10, 25–31.
56. Endris, N.; Asefa, H.; Dube, L. Prevalence of Malnutrition and Associated Factors among Children in Rural Ethiopia. *Biomed. Res. Int.* 2017, 2017. [CrossRef] [PubMed]
57. Ettyang, G.A.K.; Sawe, C.J. Factors Associated with Stunting in Children under Age 2 in the Cambodia and Kenya 2014 Demographic and Health Surveys. DHS Work Pap No 126. 2016. Available online: <http://dhsprogram.com/pubs/pdf/WP126/WP126.pdf> [accessed on 11 March 2018].
58. Fentahun, W.; Wubshet, M.; Tariku, A. Undernutrition and associated factors among children aged 6–59 months in East Belesa District, northwest Ethiopia: A community based cross-sectional study. *BMC Public Health* 2016, 16, 1–10. [CrossRef] [PubMed]

59. Jayatissa, R.; Bekele, A.; Piyasena, C.L.; Mahamithawa, S. Assessment of nutritional status of children under five years of age, pregnant women, and lactating women living in relief camps after the tsunami in Sri Lanka. *Food Nutr. Bull.* 2006, 27, 144–152. [CrossRef] [PubMed]
60. Nkurunziza, S.; Meessen, B.; Van geertruyden, J.P.; Korachais, C.; Nkurunziza, S.; Meessen, B.; Van geertruyden, J.P.; Korachais, C. Determinants of stunting and severe stunting among Burundian children aged 6–23 months: Evidence from a national cross-sectional household survey, 2014. *BMC Pediatr.* 2017, 17. [CrossRef] [PubMed]
61. Ocampo-Guirindola, M.L.; Garcia-Malabad, C.J.; Valdeabella-Maniego, M.L.M.; Punzalan, S.L.M. Association between dietary diversity score and nutritional status of Filipino children aged 6–23 months. *Philipp. J. Sci.* 2016, 145, 57–69.
62. Bbaale, E. Maternal education and child nutritional status: Evidence from Uganda. *Afr. J. Econ. Manag. Stud.* 2014, 5, 52–74. [CrossRef]
63. Makoka, D.; Masibo, P.K. Is there a threshold level of maternal education sufficient to reduce child undernutrition? Evidence from Malawi, Tanzania and Zimbabwe. *BMC Pediatr.* 2015, 1–10. [CrossRef]
64. Nyaruhucha, C.N.M.; Msuya, J.M.; Mamiro, P.S.; Kerengi, A.J. Nutritional status and feeding practices of under-five children in Simanjiro District, Tanzania. *Tanzan. Health Res. Bull.* 2006, 8, 162–167. [CrossRef]
65. Vollmer, S.; Bommer, C.; Krishna, A.; Harttgen, K.; Subramanian, S.V. The association of parental education with childhood undernutrition in low- and middle-income countries: Comparing the role of paternal and maternal education. *Int. J. Epidemiol.* 2017, 46, 312–323. [CrossRef] [PubMed]
66. Tessema, M.; Belachew, T.; Ersino, G. Feeding patterns and stunting during early childhood in rural communities of Sidama, south Ethiopia. *Pan. Afr. Med J.* 2013, 14, 1–12.

# **Chapter 4**

## **Paper III**

#### **4. Calculator for Inadequate Micronutrient Intake (CIMI): Validation of the Software in 12-23-Months-old Children and Lactating Women of Rural Ethiopia**

Beruk Berhanu Desalegn<sup>1,2\*</sup>, Christine Lambert<sup>2</sup>, Ute Gola<sup>3</sup>, Simon Riedel<sup>2</sup>, Tegene Negese<sup>1</sup> and Hans Konrad Biesalski<sup>2</sup>

<sup>1</sup>College of Agriculture, Hawassa University, Postal code: 05, Hawassa, Ethiopia; [tegenengss38@gmail.com](mailto:tegenengss38@gmail.com)

<sup>2</sup>Institute of Biological Chemistry and Nutrition, University of Hohenheim, Garbenstr. 30, 70593 Stuttgart, Germany; [christine.lambert@uni-hohenheim.de](mailto:christine.lambert@uni-hohenheim.de) (C.L.); [riedel@sciencedataservices.com](mailto:riedel@sciencedataservices.com) (S.R.); [biesal@uni-hohenheim.de](mailto:biesal@uni-hohenheim.de) (H.K.B.)

<sup>3</sup>Day-med-concept GmbH, Garbenstr. 30, 70593 Berlin, Germany; [Ute.Gola@t-online.de](mailto:Ute.Gola@t-online.de) (U.G.)

\* Correspondence: [berhanuberuk@gmail.com](mailto:berhanuberuk@gmail.com) Tel.: +251-941-048-918

**A version of this chapter was submitted to *Nutrients* on 22/05/2019 as**  
Desalegn, B., Lambert, C., Widmer, C., Gola, U., Riedel, S., Negese, T., Biesalski, H.K.  
Calculator for Inadequate Micronutrient Intake (CIMI): Validation of the software in 12-23-  
months-old children and lactating women of rural Ethiopia.

# **Calculator for Inadequate Micronutrient Intake (CIMI): Validation of the Software in 12-23-Months-Old Children and Lactating Women of Rural Ethiopia**

## **Abstract**

Background: Religion and religious activities like fasting play a pivotal role in the dietary pattern of an individual in countries like Ethiopia. However, micronutrient intake and its effect during Ethiopian fasting vs. non-fasting periods have not been assessed. Many of the dietary assessment studies in Ethiopia use only diet diversity score (DDS), a proxy indicator for micronutrient adequacy because most of the quantitative dietary assessment techniques are time consuming and need experts to calculate dietary intake. Therefore, the objective of this study was to assess the dietary nutrient intakes of 12-23-months-old children and lactating women at fasting vs. non-fasting period in rural Ethiopia; to be able develop and validate a Calculator for Inadequate Micronutrient Intake (CIMI) program, which is a simple, easy-to-use app based on amounts of different food and beverage groups which calculates dietary micronutrient intake of an individual. Methods: A community-based longitudinal assessments were conducted using 12-23-months-old children and lactating women in lent fasting (n=377 and 568) and non-fasting (n=404 and 518) periods in rural Ethiopia. Statistical analyses were performed using SPSS and Excel. For validation, individual nutrient intake was analyzed by two different methods: CIMI and the established nutrition assessment software NutriSurvey (NS). Afterwards, average mean intake and percentage of participants identified with an inadequate intake of  $<2/3$  of recommended nutrient intake (RNI) were compared using the two methods. Results: Energy and almost all macro- and micronutrients intakes of children and women were lower in lent fasting period compared to non-fasting. Likewise, the prevalence of inadequate intake of energy, protein and

most micronutrients were higher in both children (1.7-9.1%) and lactating women (1.6-21.4%) during lent fasting than non-fasting period. Based on CIMI and NS calculations, average dietary nutrient intake for children ( $R=0.741-0.956$ ) and lactating mothers ( $R=0.779-0.920$ ) were comparable. Conclusion: More attention should be given to nutrition intervention related to diet improvement activities of young children and lactating women especially during religious fasting periods. CIMI developed for the rural Ethiopian setting estimates the average nutrient intake accurately; and identifies inadequate micronutrient intake of individuals using the RNI of FAO/WHO enabling enumerators to provide feedback and suggest improvements.

Keywords: lent fasting; children; lactating mothers; CIMI; micronutrient; dietary intake; Ethiopia

#### **4.1. Introduction**

Religious fasting plays a pivotal role for changing the dietary pattern of households, as a result the nutritional status of individuals living in the households, especially nutritionally vulnerable children and women [1–4]. In Ethiopia, more than two-third of the population are either a follower of Ethiopian Orthodox or Muslim, who practice a strict fasting during their religious fasting periods [5]. But, the fasting traditions and the number of fasting days in the calendar year are prominently different between these two religions [6]. In Ethiopian orthodox religion, lent fasting is the longest among the seven official fasting periods. In this term, abstaining from any animal source foods completely, and any foods and water for at least some hours daily, except Saturday and Sunday is religious obligatory. Accordingly, studies in the 6-23-months-old children and lactating mothers in Ethiopia indicated that maternal and child undernutrition and consumption of foods prepared from low diverse food groups were higher in lent fasting compared to non-fasting period in rural Ethiopia [1,2]. To the best of our knowledge, there is no

study which has assessed the dietary micronutrient intake of Ethiopian Orthodox women and their children.

For appropriate growth and development, micronutrients are crucial apart from the macronutrients (carbohydrates, proteins and fats) [7]. The inadequate intake of these micronutrients below the amount required leads to the invisible form of malnutrition called micronutrient deficiency or 'Hidden hunger' [8]. If this inadequate supply of micronutrient persists for a longer period, then it will be increasing the risk of disease, growth and cognitive development impairment at the early stage of childhood, and maternal illness and decrease their life expectancy. Therefore, early identification of micronutrient deficiencies will be an important step to design and initiate appropriate prevention and intervention activities. But these are lacking, because quantitative nutrient intake data are often not available. As many of the existing dietary nutrient intake assessment techniques (food frequency questionnaires (FFQs), food weight record and 24-h recall) are time consuming, need well trained field workers during the surveys and an expert to calculate dietary intake using a nutrition software and identify undernutrition. Alternatively, many researches use different types of diet diversity scores as proxy indicator for dietary adequacy. These might be one of the reason why hidden hunger is not addressed properly in Ethiopia and other African countries, where this phenomenon remains high and persistent at large [9]. Thus, developing a tool that can evaluate the nutrient intake and detect the gap between the nutrient intake and the recommended intake, but easy-to-use, fast and applicable in rural poor setting is important to early identify the problem, set an appropriate prevention and intervention activities and monitor the progress.

The Calculator for Inadequate Micronutrient Intake (CIMI) is a simple, easy-to-use, informative, web-based application method, which was first developed using commonly consumed

Indonesian foods for Indonesian population. CIMI calculates energy and nutrient intake accurately, and identifies nutrient inadequacy according to FAO/WHO recommended nutrient intake (RNI) regarding age, sex and physiological stage [10]. Therefore, the purpose of this study was

1) to assess the dietary nutrient intakes of 12-23-months-old children and lactating women during Ethiopian Orthodox fasting and non-fasting periods, in rural Tigray, North Ethiopia

2) to adapt and validate CIMI, a simple, easy-to-use, informative, web-based application method which can calculate the micronutrient intake of Ethiopian population based on their traditional dietary patterns.

## **4.2. Materials and Methods**

### **4.2.1. Study Area, Design, Participants, Sampling and Ethical Clearance**

The survey was conducted during lent fasting and non-fasting period in Rural Genta Afeshum woreda of North Ethiopia, between February–April and May 2017, respectively. Description of the whole study areas, sample size calculations, sampling methods, inclusion and exclusion criteria and ethical issues were published, elsewhere [1,2]. However, in this study, the children between 6-11-month and > 23 months of age in either of the surveys, and those lost for follow-up study were excluded and thus the final total sample size was reduced to 377 and 404 in lent fasting and non-fasting assessments, respectively. Those children who were < 12-month in the lent fasting but aged  $\geq$  12-month in the non-fasting period were included in the study. Lactating mothers also participated in the present study in lent fasting (n=568) and non-fasting periods (n=518). The lactating mothers participated in the lent fasting period, but their children > 23 months of age were excluded from the survey during non-fasting period.



#### **4.2.2. Data Collection**

Before conducting the quantitative dietary survey, the principal investigator identified food items, recipes, cooking and serving materials and how they were served commonly for the children and women in the study area. Food items were measured using a laboratory weight balance (2 kg maximum weight: Model CS 2000, Ohaus Corporation, Newark, NJ, USA). Then, common food pictures and utensils were also used to support the mothers for estimating the amount of food consumed by them and their children. Following this, trainings were given for the 10 experienced data collectors who participated in the surveys. Details about the data collectors and the questionnaires used are found in previous published articles [1,2].

#### **4.2.3. Development of Calculator for Inadequate Micronutrient Intake (CIMI) for Ethiopia**

Before structuring the food groups for the CIMI program, the type of food items commonly consumed in Ethiopian population were identified from Socio-economic survey dataset (2016) based on the amount consumed in kg per capita per day. Beside this, we also reviewed literatures and published articles which provide information related to food consumed in Ethiopia. Accordingly, 93 food items were identified and categorized into 31 food groups (Table 4.1). Then, the energy, macro-, and micro-nutrients composition of each food item in the 31 food groups was adopted from Ethiopian composition table III and IV, published articles elsewhere in Ethiopia or Kenyan food composition tables [11]. Additionally, missing nutrients were taken from US Department of Agriculture database (USDA) 2018. Based on this, the nutrient profile of each food group was identified, using the share contribution of each food item contributing to the specific food group and calculated accordingly. Following this, the amount of nutrient intake was calculated by CIMI out of the sum of nutrient intakes of all food groups. An algorithm for

classifying the bioavailability of iron and zinc was adapted from CIMI developed for Indonesia [10]. Then, CIMI automatically calculated the absolute intake and percent of RNI fulfillment of energy and each of the 14 nutrients included.

#### **4.2.4. Data Management and Analyses**

The 93 food items considered for the development of CIMI for Ethiopia with their nutrient compositions were entered in NutriSurvey (NS) software version 2007 as database for analysis. Beside for the development of CIMI software, CIMI server was created. The purpose of this server is to store the dietary data entered in the CIMI software after the data automatically synced when the tablet gets Wi-Fi network. Then, the estimated amount of food items consumed in the last 24 hours by the children and women involved in this study were entered in NS, which calculated the nutrient intake. However, the estimated amount of food items was sorted into the 31 food groups categorized and summed up together to get the amount consumed as a food group and then entered into CIMI software. NS is an internationally established nutritional assessment software. In this software, each food item eaten has to be entered with the estimated amount individually. Data entries on NS were carried out by the principal investigator, but for CIMI, five experienced data entrant were trained and entered the data. Then, the tablets with CIMI data were synced in the Wi-Fi network to the server daily after the data entry and finally the results were downloaded from the server. Following these, variables coding, data transfer, cleaning and completeness of data were carried out in SPSS for window version 20 (IBM Corporation, Armonk, NY, USA), consecutively. The average nutrient intakes and the number/percentage of participants below the inadequate intake cutoff of  $<2/3$  RNI were calculated both for NS and CIMI software's. Specifically, for iron and zinc, the RNI could be different because only CIMI considers different bioavailability levels, as a result the cutoff differs for individuals. The

average mean differences (Z) of each nutrient intake was calculated as an average of the mean difference results obtained by subtracting the results produced by CIMI (Y) from NS results (X) of the participants included in the study (**Equation a**), and the average mean difference % (R) for each nutrient was calculated as the average mean difference (Z) divided by the average nutrient result produced by NS and then multiplying the dividend by 100 (**Equation b**).

$$Z = \frac{\sum_{i=1}^n [X_i - Y_i]}{n} \quad [\text{Equation a}]$$

$$R = \left[ \frac{Z}{\frac{\sum_{i=1}^n X_i}{n}} \right] * 100 \quad [\text{Equation b}]$$

where Z is the average mean difference of each nutrient and energy, X is the nutrient value of each participant calculated by NS, Y is the nutrient value of each participant calculated by CIMI, n is the number of participants, R is the average mean difference % of each nutrient and energy.

Based on this, the mean difference % expressed in terms of NS was used to categorize the nutrient intake result produced by CIMI as very high accurate (+/-0<5%), good accuracy (+/-5-15%) moderate accuracy (+/-15-30%) and low accuracy (+/->30%) [12]. Additionally, the correlation on each nutrient intake results between CIMI and NS for 12-23-months-old children and lactating women were analyzed using Pearson correlation test at the significance level of p=0.001. Microsoft Excel 2010 was also used to compute dietary intake data.

**Table 4.1** Description of the food groups in CIMI Ethiopia

Food group	Foods contributed to the calculation of the average nutrient content of the food group
White maize	White maize
Yellow maize	Yellow maize
Wheat	Wheat
Sorghum and millet	Sorghum and Finger millet
Red teff	Red Teff
White teff	White teff
Barley and related	Barley, Rice and Oats
Beans	Haricot bean, Broad bean, Chickpea, Pea, Lentil, Kidney bean, Soybean and Vetch
Oil seeds	Linseed, Niger seed, Peanut, Sesame seed, Sunflower seed, Safflower seed, Amaranth seed and white Fenugreek
Kocho and bulla	<i>Bulla and Kocho</i>
Potato and related	Irish potato, Sweet potato, Yam, Cassava, Taro, Anchote and <i>Amicho</i>
Meat	Goat meat, Sheep meat, Beef meat and Chicken meat
Fish	Fish
Egg	Chicken egg
Milk	Milk, cow
Cheese	Cheese, cow
Butter and oils	Butter, Vegetable oil and Palm oil
Sugar and honey	Sugar and Honey
Sugarcane	Sugarcane
Moringa	Moringa
Carrot and related	Pumpkin and Carrot
Green leaves	Chard, Lettuce and Ethiopian kale
Green pepper and related	Green pepper and Garlic
Pepper powder	Pepper powder
Other vegetables	Tomato, Onion, Cabbage, Beets and Snap bean ( <i>Fossolia</i> )
Mango and related	Mango, Papaya, Guava and Apricot
Banana	Banana
Orange and related	Orange, Mandarin, Avocado, Pineapple, Apple, Strawberry, Lemon, Cactus, Passion fruit, Custard apple, Watermelon, Peach and Pomegranate
Coffee and tea	Tea and Coffee
Coca Cola and Mirinda	Berez, Coca Cola, Pepsi Cola, Fanta, Mirinda and Sprite
Mild alcohol	Beer, <i>Tella</i> , <i>Tej</i> and <i>Borde</i>

### 4.3. Results

Among the 12-23-months-old children, 166 (44.0%) in fasting and 182 (45.0%) in the non-fasting period assessed were females. The mean age of the children and mothers during fasting and non-fasting periods were  $16.8 \pm 3.3$  and  $17.6 \pm 3.5$  months; and  $29.8 \pm 6.34$  and  $29.8 \pm 6.4$  years, respectively. The median LAZ, WAZ and WLZ for the children were -1.70, -1.07 and -0.42, respectively and the BMI for the mothers was 19.27 during the lent fasting period. But the WAZ and BMI were improved in the non-fasting period to -0.91 for the children and to 19.3 for mothers (Table 4.2).

**Table 4.2** Socio-demographic characteristics of the participants involved in the study in rural Southern and Northern Ethiopia

Variables		Northern Ethiopia	
		Fasting season*	Non-fasting season**
		N (%)	N (%)
Sex of children (n=377*, 404**)	Female	166 (44.0)	182 (45.0)
	Male	211 (56.0)	222 (55.0)
Mean age of women (years) (n=568*, 518**)		29.8 (6.3)	29.76 (6.4)
Mean age of children (months) (n=377*, 404**)		16.8 (3.3)	17.60 (3.5)
Nutritional status of children (n=373*, 386**) Median (SD)	LAZ	-1.70 (-2.45, -0.96)	-
	WAZ	-1.07 (-1.78, -0.46)	-0.91 (-1.50, -0.28)
	WLZ	-0.42 (-1.11, 0.22)	-
Nutritional status of women Median (SD) (kg/m <sup>2</sup> ) (n=566*, 517**)	BMI	19.3 (17.9, 21.0)	19.3 (17.9, 21.2)

The median intake of energy, calcium, iron, zinc, vitamin B1, niacin and vitamin C 24-hr prior to the start of the survey were below the standard estimated for the 12-23-months-old breastfeeding children considering the low bioavailability of iron and zinc, in both fasting and non-fasting periods, according to NS and CIMI software's. But, the estimated need for protein and vitamin A were fulfilled in both study periods, except for vitamin A calculated by CIMI. Likewise, the median intake of energy, calcium, zinc, vitamin C, vitamin A and niacin (except for non-fasting using NS) for the lactating mothers were below the RNI standard in both the study periods. However, the dietary protein, iron and vitamin B1 intakes satisfied the RNI standard for the same group, regardless of the study periods. Furthermore, except vitamin A, the other macro- and micro-nutrients and energy intakes calculated by the CIMI and NS software's were comparable. Moreover, the energy, protein and all other micronutrients intakes were improved during non-fasting than fasting period in both study groups (Table 4.3).

**Table 4.3** Average intake of energy and nutrients in 12-23 months old children and lactating mothers calculated by NS and CIMI

Nutrients		Children (12-23months)				Lactating women			
		Fasting season (n=377)		Non-fasting season (n=404)		Fasting season (n=568)		Non-fasting season (n=518)	
		NS	CIMI	NS	CIMI	NS	CIMI	NS	CIMI
Energy (kcal)	Mean (SD)	482 (267)	477(325)	532 (331)	529 (356)	1891(717)	1838 (706)	2150(530)	2079(502)
	Median (25th, 75th)	457(302, 637)	431 (270, 625)	489 (330, 653)	475 (315, 630)	1930(1313, 2358)	1871(1283, 2299)	2123 (1812, 2277)	2092(1750, 2398)
Protein (g)	Mean (SD)	13.9 (8.8)	14.8 (11.6)	14.7(9.6)	15.3(11.0)	57.0 (25.4)	58.7(27.9)	61.2 (16.3)	60.6 (15.9)
	Median (25th, 75th)	12.2 (8.0, 18.5)	12.4 (7.1, 18.9)	13.0 (8.9, 18.4)	12.6 (8.9, 19.1)	55.9 (36.1, 74.5)	57.1 (36.2, 77.9)	59.9 (50.7, 70.6)	60.8 (50.1, 70.5)
Iron (mg)	Mean (SD)	8.5 (5.0)	9.5 (7.5)	9.3(6.3)	10.0 (7.3)	36.7 (16.1)	38.4 (17.3)	41.5 (12.8)	41.5 (13.4)
	Median (25th, 75th)	7.7 (5.0, 11.6)	7.9 (4.96, 12.0)	8.1(5.6, 11.5)	8.2 (5.8, 12.1)	36.9 (24.34 45.6)	38 (24.5, 48.8)	40.2 (34.6, 47.7)	41.6(34.5, 47.5)
Zinc (mg)	Mean (SD)	3.50 (2.45)	3.44 (2.63)	3.57(2.42)	3.62(2.72)	15.1(6.6)	14.2 (6.5)	16.5 (4.3)	15.6 (3.95)
	Median (25th, 75th)	3.04 (1.85, 4.69)	3.03 (1.77, 4.43)	3.08 (2.14, 4.48)	3.05 (2.19, 4.43)	15.1(9.7, 19.4)	14.4 (9.2, 17.9)	16.2 (13.7, 19.2)	16(13.1, 17.9)
Vitamin A (µg RE)	Mean (SD)	298 (255)	99.2(98.3)	380 (395)	147 (187)	486 (413)	228 (327)	675 (676)	331 (475)
	Median (25th, 75th)	266 (130, 395)	81.2 (30.7, 152)	255(76.6, 563)	95.7(30.8, 185)	377 (234, 571)	154 (87, 238)	525(185, 887)	195 (77.8, 348)
Calcium (mg)	Mean (SD)	81.0 (61.0)	88.1 (74.5)	84.5 (64.2)	91.1 (72.1)	322(148)	355 (191)	330 (105)	341(118)
	Median (25th, 75th)	66.0 (41.0, 110.0)	66.8 (43.9, 117)	69.1 (47.3, 101.7)	72.3 (47.3, 110)	312(200, 412)	336 (204, 459)	323 (270, 380)	324 (270, 387)
Mg (mg)	Mean (SD)	132 (83.3)	154 (122)	144.5(104.6)	161.8(125)	605(226)	660(270)	708(175)	724 (178)
	Median (25th, 75th)	120 (74.1, 175)	132 (83.5, 209.5)	121.5(86.4, 178.8)	134(90.0, 197)	612 (431, 748)	670 (443, 837)	705 (608, 827)	743 (618, 838)
Vitamin B1 (mg)	Mean (SD)	0.39 (0.26)	0.42 (0.33)	0.42(0.28)	0.44(0.33)	1.75(0.73)	1.77 (0.77)	1.89 (0.47)	1.88 (0.47)
	Median (25th, 75th)	0.36 (0.21, 0.54)	0.36 (0.21, 0.55)	0.37(0.25, 0.52)	0.37(0.26, 0.52)	1.75(1.14, 2.23)	1.77 (1.14, 2.28)	1.89(1.61, 2.23)	1.92(1.58, 2.19)
Niacin (mg)	Mean (SD)	3.12 (1.97)	3.21 (2.44)	3.32(2.34)	3.45(2.79)	14.8 (5.3)	13.9 (5.3)	17.3 (4.64)	16.1 (4.2)
	Median (25th, 75th)	2.90 (1.72, 4.11)	2.81 (1.78, 4.33)	2.80(1.99, 4.14)	2.93(2.00, 4.14)	14.9 (11.0, 18.3)	13.9 (9.9, 17.6)	16.8(14.5, 19.6)	16.2 (13.5, 18.3)
Vitamin B6 (mg)	Mean (SD)	0.15 (0.14)	0.19 (0.22)	0.17(0.19)	0.21(0.23)	0.75(0.54)	0.89 (0.60)	0.82 (0.64)	0.87(0.62)
	Median (25th, 75th)	0.11 (0.05, 0.18)	0.13 (0.06, 0.26)	0.12(0.07, 0.20)	0.14(0.08, 0.27)	0.62(0.40, 0.93)	0.75 (0.43, 1.19)	0.62(0.37, 1.02)	0.69(0.45, 1.07)
Vitamin B12 (µg)	Mean (SD)	0.19 (1.71)	0.10 (0.20)	0.12(0.24)	0.11(0.23)	0.01(0.50)	0.01(0.48)	0.09 (0.41)	0.07(0.26)
	Median (25th, 75th)	0.00(0.00, 0.00)	0.00(0.00, 0.00)	0.00(0.00, 0.19)	0.00(0.00, 0.03)	0.00(0.00, 0.00)	0.00(0.00, 0.00)	0.00(0.00, 0.00)	0.00(0.00, 0.00)
Pentatonic acid (mg)	Mean (SD)	1.16 (1.81)	0.91 (0.67)	1.12(0.73)	0.98(0.70)	4.05 (2.00)	3.34 (1.52)	3.91 (1.12)	3.49 (0.91)
	Median (25th, 75th)	0.90 (0.57, 1.42)	0.79 (0.47, 1.22)	1.01(0.63, 1.44)	0.85(0.54, 1.24)	3.86 (2.40, 5.41)	3.33 (2.16, 4.23)	3.87 (3.18, 4.62)	3.49 (2.91, 4.01)
Vitamin C (mg)	Mean (SD)	11.9 (12.5)	11.2 (11.9)	13.9(13.2)	12.98(12.81)	47.7 (28.4)	65.7 (34.4)	48.2 (23.5)	64.0 (29.1)
	Median (25th, 75th)	6.1 (2.87, 20.8)	6.9 (2.87, 16.8)	8.4(3.58, 23.1)	9.9(3.35, 17.7)	41.73(29.64, 58.64)	60.5 (44.0, 83.8)	41.3(32.7, 57.7)	61.3 (42.2, 81.5)
Vitamin D (µg)	Mean (SD)	0.34 (1.48)	0.20 (0.39)	0.24 (0.47)	0.25 (0.48)	0.01(0.11)	0.03 (0.10)	0.03(0.19)	0.06(0.19)
	Median (25th, 75th)	0.00 (0.00, 0.00)	0.02 (0.00, 0.04)	0.00 (0.00, 0.19)	0.02(0.01, 0.09)	0.00(0.00, 0.00)	0.02(0.01, 0.03)	0.00(0.00, 0.00)	0.02(0.01, 0.05)

The estimated need for the 12-23-months old breastfed children from the complementary foods considering the average breast milk intake was adapted [13–15]. The estimated need of energy, protein, calcium, iron, zinc, vitamin B1, vitamin C and vitamin A were 548kcal, 5g, 346mg, 11.4mg (low bioavailability), 7.6mg (low bioavailability), 0.38mg, 5.18mg, 8mg and 126 retinol equivalent ( $\mu\text{g RE}$ ), respectively for the 12-23-months old breastfeeding children. For the lactating women, the recommended nutrient intake (RNI): energy (2650kcal), protein (55g), calcium (100mg), iron (30mg for low bioavailability), zinc (17mg low for bioavailability), vitamin B1 (1.5mg), niacin (17mg), vitamin C (70mg) and vitamin A (850  $\mu\text{g RE}$ ) [16–18].

Inadequate intakes of many nutrients and energy ( $n=10$ ) were prevalent ( $>50\%$ ) for the 12-23-months-old children in fasting and non-fasting periods, except for protein iron, magnesium and vitamin B1. Inadequate intake of vitamin A, calcium, vitamin D, vitamin C (only for results calculated by NS) vitamin B12, vitamin B6 and pantoic acid was also prevalent ( $>50\%$ ) in lactating mothers. Beside these, the prevalence of inadequate intakes of vitamin B1, protein and iron for the 12-23-months-old children and energy, zinc and niacin intake for lactating mothers during fasting were between 30-69%, but these decreased during the non-fasting period, except the vitamin C for lactating mothers. The prevalence of low intake of magnesium for the 12-23-months-old children and protein, iron, magnesium and vitamin B1 for the lactating mothers were less than 30%. These also declined in non-fasting compared to fasting periods, regardless of the software used for identification of the inadequacy (Table 4.4).

**Table 4.4** Number and percentage of 12-23 months old children and lactating women below the threshold of inadequate intake (<2/3 RNI)

	Children (12-23months)				Lactating women			
	Fasting period (n=377)		Non-fasting period (n=404)		Fasting period (n=568)		Non-fasting period (n=518)	
Nutrient	CIMI n (%)	NS n (%)	CIMI n (%)	NS n (%)	CIMI n (%)	NS n (%)	CIMI n (%)	NS n (%)
Energy (kcal)	273 (72.4)	272(72.1)	289 (71.5)	270 (66.8)	248 (43.7)	242 (42.6)	134 (25.9)	114 (22.0)
Protein (g)	120 (31.8)	114(30.2)	90 (22.3)	92(22.8)	146 (25.7)	151 (26.6)	32 (6.2)	27 (5.2)
Iron (mg)	180 (47.7)	191 (50.7)	180 (44.6)	193 (47.8)	96 (16.9)	93 (16.4)	0 (0.00)	0 (0.00)
Zinc (mg)	324 (85.9)	323(85.7)	352 (87.1)	348 (86.1)	197 (34.7)	179 (31.5)	71 (13.7)	53(10.2)
Vitamin A (µg RE)	363 (96.3)	188 (49.9)	340 (84.2)	207(51.2)	544 (95.8)	424 (74.6)	445 (85.9)	277 (53.7)
Calcium (mg)	368 (97.6)	375 (99.5)	399 (98.8)	402(99.5)	526 (92.6)	562 (98.9)	509 (98.3)	515 (99.4)
Magnesium (mg)	41 (10.9)	39 (10.3)	24 (5.9)	33(8.2)	9 (1.60)	9(1.60)	1 (0.20)	0 (0.00)
Vitamin B1 (mg)	170 (45.1)	165 (43.8 )	163 (40.3)	177(43.8)	120 (21.1)	105 (18.5)	1 (0.20)	15 (2.9)
Niacin (mg)	274 (72.7)	274 (72.7)	252 (62.4)	257 (63.6)	187 (32.9)	152 (26.8)	63 (12.2)	35(6.8)
Vitamin B6 (mg)	309 (82.0)	338 (89.7)	374 (92.6)	381 (94.3)	460 (81.0)	493 (86.8)	432 (83.4)	435 (84.0)
Vitamin B12 (µg)	365 (96.8)	360 (95.5)	381 (94.3)	379 (93.8)	568 (100.0)	568 (100)	511 (98.6)	507 (97.9)
Pantatonic acid (mg)	295 (78.2)	271 (71.9)	340 (84.2)	308 (76.2)	467 (82.2)	364 (64.1)	477 (92.1)	398 (76.8)
Vitamin C (mg)	318 (84.4)	277 (73.5)	317 (78.5)	268 (66.3)	160 (28.2)	331 (58.3)	153 (29.5)	309 (59.7)
Vitamin D (µg)	377 (100)	372 (98.7)	404 (100)	404 (100)	568 (100)	568 (100)	518 (100)	518 (100)

NB: The inadequacy of the 12-23-months old children and lactating women was declared as the energy, protein and other micronutrients intake was below two-third of the recommended nutrient intake (RNI). For the zinc and iron, the RNI for CIMI was set based on the individual consumption pattern and category of bioavailability, otherwise for the NS, the low and 5% bioavailability of zinc and iron was used in this study, respectively. The RNI for the 12-23-months old breastfed children was energy (950kcal (male), 850kcal (female)), protein (13.5g), calcium (500mg), iron (11.6mg), zinc (8.3mg), vitamin B1 (0.5mg), niacin (6mg), vitamin B6 (0.5mg), vitamin C (30mg), pantothenic acid (2mg), vitamin B12 (0.9 µg), vitamin A (400 µg), vitamin D (5 µg) and magnesium (60mg). For the lactating mothers, the recommended nutrient intake (RNI): energy (2650kcal), protein (55g), calcium (1000mg), iron (30mg and 15mg for very low and low bioavailability, respectively), zinc (17mg low for bioavailability), vitamin B1 (1.5mg), niacin (17mg), vitamin C (70mg), vitamin A (850RAE), vitamin B6 (2mg), pentatonic acid (7mg), vitamin B12 (2.8 µg), vitamin D (5 µg) and magnesium (270mg) [16–18].



**Table 4.5** Average mean differences of calculated nutrient intake by CIMI and NS. Categorization of accuracy was performed according percentage of mean difference in relation to mean NS result


Nutrient	Children (12-23months) (n=781)			Lactating women (n=1086)		
	Mean difference between CIMI and NS result (+SD)	Mean difference in % of mean nutrient intake calculated by NS	Accuracy category	Mean difference between CIMI and NS result (+SD)	Mean difference in % of mean nutrient intake calculated by NS	Accuracy category
Energy (kcal)	4.15 (186.23)	1.06	Very high accurate	59.05 (273.33)	1.91	Very high accurate
Protein (g)	-0.71 (5.68)	-3.49	Very high accurate	-0.60 (8.99)	-1.71	Very high accurate
Iron (mg)	-0.82 (4.31)	-9.48	Good accurate	-0.85 (8.05)	-3.42	Very high accurate
Zinc (mg)	0.01 (1.71)	-4.09	Very high accurate	0.92 (2.91)	4.44	Very high accurate
Vitamin A (µg RE)	213.31 (202.21)	-22.90	Moderate accurate	301.01 (353.64)	24.51	Moderate accurate
Calcium (mg)	-6.75 (43.06)	-8.74	Good accurate	-22.46 (83.27)	-7.08	Good accurate
Magnesium (mg)	-19.66 (73.37)	-16.16	Moderate accurate	-36.02 (117.26)	-6.50	Good accurate
Vitamin B1 (mg)	-0.02 (0.17)	-6.34	Good accurate	0.00 (0.27)	-0.96	Very high accurate
Niacin(mg)	-0.12 (1.57)	-5.31	Good accurate	1.05 (2.48)	5.74	Good accurate
Vitamin B6 (mg)	-0.04 (0.14)	-36.08	Low accurate	-0.10 (0.31)	-21.28	Moderate accurate
Vitamin B12 (µg)	0.01 (0.17)	-	-	0.01 (0.15)	-	-
Pentatonic acid (mg)	0.15 (0.40)	10.77	Good accurate	0.57 (0.85)	11.23	Good accurate
Vitamin C (mg)	0.82 (8.03)	-8.04	Good accurate	-16.95 (19.09)	-42.06	Low accurate
Vitamin D (µg)	0.00 (0.13)	-	-	-0.03 (0.09)	-	-


NB: Nutrients with a very high accuracy (difference expressed as % of NS result: +/-0<5%), good accuracy (+/-5-15%), moderate accuracy (+/-15-30%) and low accuracy (+/->30%), this is adopted from Lambert et al., 2018. For vitamin B12 and vitamin D, the mean difference % was not calculated, as most of these values were zero in NS, which is a denominator for computation.


Among the mean differences of CIMI from NS calculated for each nutrient intake, the highest variation was observed in vitamin A, both for children and mothers, whereas the lowest was observed in vitamin D and vitamin B1 for children and mothers, respectively. However, the % of mean difference using the actual average nutrient intake in terms of NS was highest in vitamin B6 for children and vitamin C for mothers. The lowest was observed in the energy and vitamin B1 for children and mothers, respectively. According to Lambert and colleagues' criteria for categorizing the precision of CIMI in terms of NS, the results of energy, protein, iron, zinc, calcium, vitamin B1, niacin and pentatonic acid were either very high or accurate. Whereas, low precision was observed in vitamin B6 for the 12-23-months-old children and vitamin C for the lactating mothers (Table 4.5).

Beside these, the highest correlation coefficient ( $R$ ) between CIMI and NS was observed in vitamin D (0.956) and protein (0.920) for the 12-23-month-old children's and lactating women, respectively. But, the lowest was observed in vitamin B12 (0.741) for the 12-23-month-old children and vitamin A (0.779) for the lactating mothers (Figures 4.1 and 4.2). All correlations were highly significant ( $P < 0.001$ ).

**Fig. 4.1** Correlation of energy and nutrient intake of Ethiopian women calculated by NS and CIMI










NB: The red line in these scatter plots is the reference line ( $y=x$ ;  $R=1$ ), which means the results produced from the two programs are identical. The black line represents the regression line of the data pairs calculated by CIMI and NS.

**Fig. 4.2** Correlation of energy and nutrient intake in between NS and CIMI in 12-23 months old children, Ethiopia







NB: The red line in these scatter plots is the reference line ( $y=x$ ;  $R=1$ ), which means the results produced from the two programs are identical. The black line represents the regression line of the data pairs calculated by CIMI and NS.

#### 4.4. Discussion

To the best of our knowledge, this is the first study in Ethiopia, which focuses on a) adaptation and validation of a simple and rapid web-based dietary nutrient intake assessment tool for Ethiopian population, b) assessment of dietary nutrient and energy intakes of the 12-23-months-



old children and their lactating mothers during religious fasting and non-fasting periods in Northern Ethiopia.

#### **4.4.1. Validation of Calculator for Inadequate Micronutrient Intake (CIMI)**

For the first time, CIMI was developed in Indonesia for the Indonesian population [10]. The CIMI software presented in this study is the second next to Indonesia, which is developed for Ethiopian population, using commonly consumed Ethiopian foods. Like CIMI developed in Indonesia, we used the food group approach which considered the amount of food consumed per capita per day, the nutrient composition of the food items specially the micronutrients that are of public health importance in Ethiopia (vitamin A, Fe, and Zn) and standard food grouping (e.g.:- vegetable to vegetable) [9,19–22]. Shape and volume of food items, local utensils or food size or volume used in local market as standardized local units and varietal differences were included to create the 31 food groups. For example, chard, lettuce and Ethiopian kale were under ‘green leaves’ food group based on their micronutrient contents (Fe, Zn and vitamin A), shape and because of being vegetables. Likewise, sweet potato, potato, taro, cassava, yam, *amicho* (boiled decorticated root part of false banana) and anchote (*Coccinia abyssinica*) are aggregated based on their shape and natural group into ‘potato and related foods’ group. Maize and teff are grouped into two groups each as white and yellow, and white and red, because these grains are highly consumed by majority of Ethiopians and based on their pro-vitamin A and Fe contents, respectively [11]. Beside this, assessment of food amounts a slider with 15 intervals, which considered the minimum and maximum amount estimated to be eaten by the young children and adult was also considered.

Accordingly, the average intake of most nutrients and energy calculated by CIMI and NS were nearly similar, except for vitamin A, in both the children and lactating women, and vitamin C in the lactating women, regardless of the periods. Our vitamin A result is in line with previous CIMI study in Indonesia [10]. The possible explanation for variable vitamin A intake estimation by CIMI in our case was, the average vitamin A content of the butter and oil food group was lower than the palm oil, which is one of the item created this group, but was available by the cooperative enterprises in the present study area and consumed during the fasting and non-fasting surveys. The availability of different oil types at different time in the country depends on the massive import oil types by the subsidy of the Ethiopian government. Therefore, deselecting the unconsumed oil type from this food group will be an important activity for future CIMI use. The reason for the high vitamin C values of CIMI for the lactating women was due to the great share (43.8%) of the tomato for the 'other vegetables' group, but less consumed compared to the cabbage, which has relatively lower vitamin C increased the average vitamin C content of the 'other vegetables' group.

Protein, iron, calcium, magnesium, vitamin B1, and B6 intake estimated by CIMI were slightly higher than NS, in both the study groups. The reason for the high CIMI of protein, iron and calcium could be due to the consumption of high amount of peas, although peas' contribution to the relevant food group is only 12.4%, which does not represent the real share of peas in this group. Peas have got a lower content of these nutrients compared to most food items in the group. Beside this, the intervals to set up the amount with the slider get larger if the consumption is increasing in most of the food groups in CIMI. Therefore, the selection of the slider either below or above the estimated amount of the food group will depend on the nearest amount set up

in the sliders. So if the amount approaches to the upper slider, then automatically the upper slider will be selected and these will lead to overestimation of intake of specific nutrients when compared to NS. The later could also work for magnesium and vitamin B1 in pepper and egg food groups, respectively. Ethiopian kale, which has got a higher vitamin B6 content than other vegetables in the 'green leaves' food group, contributed 58.4% for this group. But Ethiopian kale was less consumed in the study area and this leads to an overestimation of vitamin B6 intake in CIMI.

The number (percentage) of participants identified as inadequate in nutrients ( $<2/3$  RNI) by CIMI and NS were similar in most nutrients considered, except for vitamin A for 12-23-months-old children and lactating women, and vitamin C only for the lactating women, regardless of the study period. These could be due to the high and low average values of the vitamin A and vitamin C in the 'butter and oil', and 'other vegetable groups' for individuals than the values in NS, respectively.

According to Lambert et al. (2018), vitamin A was moderately calculated by CIMI for the 12-23-months-old children and lactating women. The reason mentioned earlier for variation in the average values of vitamin A produced by CIMI and NS, leads to higher mean % difference, which is the criteria for this classification. Likewise, CIMI calculated the vitamin B6 intake with low and moderate precisions for the 12-23-months-old children and lactating women, respectively. The possible reason could be the small mean vitamin B6 differences between CIMI and NS in few individuals that increased their mean difference (%) dramatically (Table 4.5). The moderate precision of magnesium in the 12-23-months-old children might be related with the average CIMI of the group which was higher than the commonly consumed pea entered in NS.

The low precision in the vitamin C variation could be due to the average vitamin C value of the ‘other vegetables’ group, which was relatively high because of tomato. Strong correlations between CIMI and NS were found in all the nutrients and energy for both study groups (Figure 4.1 and 4.2). The correlation coefficients between CIMI and NS of protein, iron, zinc and vitamin A for the 12-23-months-old children and energy, protein, iron and zinc for lactating mothers were higher compared with the previous study in Indonesian population [10]. This could be attributed to the smaller sample size in the Indonesian study. Furthermore, the correlation coefficients of most nutrients in our study were higher than validation results of CIMI in Tanzania women, except for energy [12]. In general, our validation study confirmed that CIMI developed for rural Ethiopian population with commonly consumed Ethiopian foods calculates the average nutrient intake accurately. Additionally, it identifies the inadequate micronutrient intake of individuals using FAO/WHO recommendation for nutrient intake (RNI). Unlike other dietary assessment techniques, CIMI provides result immediately after the interview is completed, which enables the enumerator to provide immediate feedback to the interviewee and suggest dietary improvements considering the socio-economic status of the household, culture of the society, availability of potential and alternative food items in local farms and/or market. Currently, most of the dietary assessment and intervention studies use diet diversity scores, a proxy indicator for assessing the dietary quality at household and individual levels [23]. However, this technique only focus on the assessment of diversity of food groups consumed, because the smaller amount of intake from the food groups will challenge the validity of the diet diversity score result, as a proxy indicator for micronutrient adequacy of an individual. To overcome this problem, many studies have used a cutoff of  $\geq 15$  g to count the food group as

consumed in women in reproductive age [23]. But, still the controversy remains on how quantification is practical in real qualitative diet diversity assessment technique, unless the diet diversity data is extracted from other quantitative dietary intake data. Thus, CIMI can replace the qualitative diet diversity, by producing much more results on the diet quality of an individual, and overcoming the challenges encountered by DDS. CIMI can also be used in cross-sectional and/or longitudinal/ intervention studies, in situation where the conventional diet diversity score is not appropriate or requires the quantitative dietary assessment methods. These indicate that CIMI developed for Ethiopian population is applicable for dietary micronutrient assessment at individual level in rural setting. The accumulation of this individual data can provide information on dietary nutrient intake at population level for designing targeted micronutrient prevention and intervention activities. In general, CIMI can be an important tool for nutrition and related researchers, practitioners, policy makers and planners in Ethiopia.

#### **4.4.2. Dietary Nutrient Intakes of 12-23-months-old Children and Lactating Women in Rural Ethiopia**

The median intake of energy, iron, zinc, thiamin, niacin and calcium in this study were below the estimated need from complementary food for the 12-23-months-old breastfeeding children, regardless of the study periods. But, the median intake of dietary protein, and vitamin C (only during non-fasting period) met their needs. These results are in line with previous studies in different parts of Ethiopia [13,24–26]. The vitamin A intake of the 12-23-months children in the present study was also above the estimated need and higher than its median intake in the 6-35-months-old children of Tigray region and other findings, elsewhere in Ethiopia [13,24–27]. This could be due to the consumption of palm oil fortified with vitamin A for preparation of

complementary food as noticed during the survey periods. Furthermore, the prevalence of inadequate dietary intakes ( $<2/3$  RNI) of zinc was very high ( $>75\%$ ) in the 12-23-months-old children, followed by vitamin C, energy and niacin (51-75%), and protein, iron, vitamin A and vitamin B1 (20-50%), respectively. But, the prevalence of low intake of energy, protein, niacin and vitamin C were decreased during non-fasting compared to fasting period. The finding of recent investigation in the same study area evidenced that the prevalence of 6-23-months-old children who fulfilled the minimum acceptable diet (MAD) criteria was 2.3% during fasting and 6.7% during non-fasting period. A study in *Libo Kemkem* and *Fogera* districts in Ethiopia showed that zinc and vitamin A deficiencies were prevalent in the school-age children [28]. Furthermore, a recent national study in Ethiopia revealed that the average dietary zinc intake of 6-35-months-old children from Tigray region and at national level were low [22]. A study conducted in four African countries including Ethiopia also revealed that anemia, vitamin A and zinc deficiencies are of public health concern in these countries [29].

In the present study, the median intake of energy, protein, vitamin A, calcium, vitamin C and niacin (fasting period only) were below the recommended nutrient intake (RNI) for lactating mothers, but not for iron, zinc and vitamin B1. Except for iron, the intake of energy, protein, zinc and vitamin A were very close to the values for Tigray region and national average [27]. Our findings on iron and calcium intake were higher than earlier studies on lactating and pregnant women in Northern part of Ethiopia [30,31]. These might be due to the fact that food taboos may be related to a reduction of food intake by mothers during pregnancy. Main reasons for food taboos are fear of obstetric complications associated with the delivery of a bigger baby, plastering on the fetal head, fear of abortion, evil eye and fetal abnormality [32–34]. The

prevalence of inadequate dietary intake of vitamin A and vitamin C was higher, followed by energy, zinc, niacin, vitamin B1 and iron. Previous studies in Ethiopian children and women confirmed that the prevalence of iron, zinc and vitamin A deficiencies was high using blood biomarkers [35–40]. Except for calcium and vitamin C, the inadequacies were decreased after the two months lent fasting periods. Therefore, considering the continuously high prevalence of inadequate intake of vitamin C in both study groups lead to the assumption that consumption of vitamin C rich foods are chronically low in the study area. Therefore, the deficiency of these micronutrients is expected to be higher in our study district and further studies using biomarkers should be done. The possible explanation for this might be related with the low consumption of fruits, animal source foods including dairy products, root crops like potato, anchote in the study area [1,2]. Surprisingly, almost all the 12-23-months-old children and lactating mothers did not take vitamin B12, vitamin D and calcium. Previous national study in Ethiopia found that only 15.1% of non-pregnant women were deficient in vitamin B12 using blood biomarker [35]. The big difference from our finding could be related to the method of assessment where the serum total vitamin B12 was used as biomarker, which gives less information in identifying vitamin B12 deficiency [41,42]. Therefore, urgent investigation on the deficiency of vitamin B12 should consider the biomarker like methyl malonic acid or Holotranscobalamin in urine and blood samples, to understand the situation in the study area more precisely[42,43]. Likewise, earlier study in Ethiopian non-pregnant women showed that there was no participant that ever consumed vitamin D-rich foods, fortified foods, or dietary supplements, and as a result Vitamin D insufficiency is a serious problem, even though Ethiopia is located near the Equator [44]. Highest prevalence of inadequate intake of calcium was also found in studies conducted in

Gondar (90.4%), Butajira (99%) districts, and at national (89-96%) level [30,45,46]. Therefore, the proposed universal calcium supplementation in Ethiopia should be implemented [46]. Therefore, nutrition sensitive agriculture in both plant and livestock production should be promoted and implemented in sustainable manner.

Previous studies in the same study population and study area evidenced that the dietary diversity score were low in general, lower in the lent fasting compared to non-fasting period, both in the lactating mothers and their 6-23-months-old children [1,2]. In line with this, our present study confirmed that dietary nutrient intake (DNI) is lower in lent fasting compared to non-fasting period, both in the young children and lactating women. These indicate that, dietary nutrient intake in the district was sub-optimal, which might be one of the multi-faceted cause for the highest prevalence of maternal and child undernutrition in the study district, Tigray region and Ethiopia at large (Table 4.2).

Our present study has many strengths. Among these, the average DNI of the young children and women and the prevalence of inadequate intake of most important nutrients were addressed with a large sample size in rural Ethiopia. The longitudinal study design considered the longest religious fasting period among the seven official Ethiopian Orthodox fasting periods, whereas the non-fasting period was the fifty days after the fasting period. For the first time in Ethiopia, dietary assessment software is developed using commonly consumed foods in rural Ethiopia. Additionally, the validity of this software was assessed using groups (children and women) vulnerable for malnutrition in religious fasting/non-fasting periods. However, only one 24-hr dietary recall was used for this assessment, and the study was conducted in only one administrative region in Ethiopia, out of the eleven, are limitations of this study.



#### **4. 5. Conclusions**

Our study indicated that except protein and magnesium, the prevalence of inadequate intake ( $<2/3$  RNI) of energy and most micronutrients was high in the 12-23-months-old children. Likewise, the prevalence of inadequate intake of vitamin A, calcium, vitamin B6, vitamin B12, pentatonic acid and vitamin C were also high in the lactating mothers. However, these high prevalences of inadequacy in energy and macro-and micronutrients in the 12-23-months-old children and lactating mothers were more affected during the lent fasting period. Therefore, the present nutrition intervention should give more attention on diet improvement activities of the children and women especially during religious fasting periods. Alarming, almost all the 12-23-months-old children and lactating mother's in the study area obtained below two-third of the RNI for vitamin B12, vitamin D and calcium from their diet. Therefore, urgent further investigation should be done for immediate public health nutrition intervention in the district, in addition to the implementation of the proposed universal calcium supplementation in Ethiopia. Our result also confirmed that CIMI developed for rural Ethiopian population estimates the average nutrient intake accurately. It also identifies inadequate micronutrient intake of individuals using FAO/WHO recommended nutrient intake (RNI) directly after the interview. This enables the enumerator to provide immediate nutrition education and suggest context based dietary improvements, unlike other dietary assessment techniques. CIMI can also replace the DDS, a qualitative dietary assessment tool, by overcoming the drawback, which is related with counting insignificant intake in the score and by producing more results. CIMI can also be used in cross-sectional and/or longitudinal/ intervention studies, in situation where the conventional

diet diversity is not appropriate or requires the quantitative dietary assessment methods. Therefore, CIMI can be used for dietary assessment survey in Ethiopia.

Funding: German Federal Ministry for Economic Cooperation and Development (BMZ).

Acknowledgments: We, the authors would like to thank the mothers and children who are participated in this study. We would like also to forward our felt thank to the data collectors, community facilitators and data encoders involved during the course of this study. Last, but not least, the first authors would like to extend the appreciation the generous financial support of German Federal Ministry for Economic Cooperation and Development (BMZ) for his PhD study at University of Hohenheim, Germany.

Conflicts of Interest: No conflict of interest among authors.

#### **4.6. References**

1. Desalegn, B.B.; Lambert, C.; Riedel, S.; Negese, T.; Biesalski, H.K. Ethiopian orthodox fasting and lactating mothers: Longitudinal study on dietary pattern and nutritional status in rural tigray, Ethiopia. *Int. J. Environ. Res. Public Health* 2018, 15 (8), 1767.
2. Desalegn, B.; Lambert, C.; Riedel, S.; Negese, T.; Biesalski, H.K. Feeding Practices and Undernutrition in 6–23-Month-Old Children of Orthodox Christian Mothers in Rural Tigray, Ethiopia: Longitudinal Study. *Nutrients* 2019, 11(1), 138.
3. Hirvonen, K.; Taffesse, A.S.; Worku, H.I. Seasonality and household diets in Ethiopia. *Public Health Nutr.* 2016,19(10), 1723–30.
4. Kumera, G.; Tsedal, E.; Ayana, M. Dietary diversity and associated factors among

- children of Orthodox Christian mothers/caregivers during the fasting season in Dejen District, North West Ethiopia. *Nutr Metab.* 2018, 15, 16.
5. CSA. Population and Housing Census of Ethiopia. 2007.
  6. Bazzano, A. N.; Potts, K. S.; Mulugeta, A. How do pregnant and lactating women , and young children , experience religious food restriction at the community level ? A qualitative study of fasting traditions and feeding behaviors in four regions of Ethiopia. *PLoS ONE* 2018, 13(12), e0208408. <https://doi.org/10.1371/journal.pone.0208408>
  7. Bhandari. S.; Banjara, M.R. Micronutrients Deficiency, a Hidden Hunger in Nepal: Prevalence, Causes, Consequences, and Solutions. *Int. Sch. Res. Not.* 2015, Volume 2015, Article ID 276469. Available from: <http://www.hindawi.com/journals/isrn/2015/276469/>
  8. Biesalski, H.K.; Jana, T. Micronutrients in the life cycle: Requirements and sufficient supply. *NFS J.* 2018, 1, 1–11.
  9. Central Statistical Agency (CSA). Ethiopia Demographic and Health Survey 2016; Central Statistical Agency: Addis Ababa, Ethiopia; ICF International Calverton: Calverton, MD, USA, 2017; p. 551.
  10. Jati, I.R.A.P.; Widmer, C.; Purwestri, R.C.; Wirawan, N.N.; Gola, U.; Lambert, C.; Biesalski, H.K. Design and validation of a program to identify inadequate intake of iron, zinc, and vitamin A. *Nutrition* 2014, 30, 1310–1317. Available from: <http://dx.doi.org/10.1016/j.nut.2014.03.015>
  11. Abebe, Y.; Bogale, A.; Hambidge, K.; Stoecker, J.; Bailey, K.; Gibson, R. Phytate, zinc ,

- iron and calcium content of selected raw and prepared foods consumed in rural Sidama , Southern Ethiopia, and implications for bioavailability. *J. Food Compos. Anal.* 2007, 20, 161–168.
12. Lambert, C.; Eleraky, L.; Mbwana, H.; Kinabo, J.; Biesalski, H.K.; Riedel, S, et al. Validation of a food group based nutrition software to assess nutrient intake in Tanzania. In: *Global food security and food safety: The role of universities*. Ghent, Belgium: Ghent University; 2018. p. 2018.
  13. Gibson, R.S.; Abebe, Y.; Hambidge, K.M.; Arbide. I.; Teshome, A.; Stoecker, B.J. Original Article Inadequate feeding practices and impaired growth among children from subsistence farming households in. *Matern Child Nutr.* 2009, 5, 260–75.
  14. Dewey, K.G.; Brown, K.H. Update on technical issues concerning complementary feeding of young children in developing countries and implications for intervention programs. *Food Nutr. Bull.* 2003, 24 (1), 5–28.
  15. WHO. Complementary feeding of young children in developing countries: a review of current scientific knowledge. Geneva, Switzerland, 1998; p. 248.
  16. FAO/WHO. Vitamin and mineral requirements in human nutrition : report of a joint FAO/WHO expert consultation, Bangkok, Thailand, 21–30 September 1998, 2004; p.362.
  17. FAO/WHO/UNU. Human energy requirements: report of a joint FAO/WHO/UNU expert consultation. Rome, Italy, 2001; p.103.
  18. FAO/WHO/UNU. Protein and amino acid requirements in human nutrition : report of a

- joint FAO/WHO/UNU expert consultation. Geneva, Switzerland, 2002, 2007; p.284.
19. Tessema, M.; Groote, H. De.; Brouwer.,I.D.; Feskens, E.J.M.; Belachew, T.; Zerfu, D.; et al. Soil Zinc Is Associated with Serum Zinc But Not with Linear Growth of Children in Ethiopia. *Nutrients* 2019, 11, 221.
  20. Tezera, R.; Sahile, Z.; Yilma, D.; Misganaw, E.; Mulu, E. Prevalence of anemia among school-age children in Ethiopia : a systematic review and meta-analysis. *Systematic Reviews* 2018, 7, 80.
  21. Demissie, T.; Ali, A.; Mekonen, Y.; Haider, J.; Umeta, M. Magnitude and distribution of vitamin A deficiency in Ethiopia. *Food and Nutrition Bulletin* 2010, 31(2), 234–41.
  22. Ayana, G.; Moges, T.; Samuel, A.; Asefa, T.; Eshetu, S.; Kebede, A. Dietary zinc intake and its determinants among Ethiopian children 6–35 months of age. *BMC Nutrition* 2018, 4(1), 30.
  23. FAO and FHI 360. Minimum Dietary Diversity for Women A Guide to Measurement. Rome, 2016; p.82.
  24. Mesfin, A.; Henry, C.; Girma, M.; Whiting, S.J. Use of pulse crops in complementary feeding of 6-23-month-old infants and young children in Taba Kebele, Damot Gale District, Southern Ethiopia. *J. Public Health Africa* 2016, 6(1), 1–6.
  25. Baye, K.; Guyot, J.P.; Icard-Vernière, C.; Mouquet-Rivier, C. Nutrient intakes from complementary foods consumed by young children (aged 12-23 months) from North Wollo, northern Ethiopia: the need for agro-ecologically adapted interventions. *Public*

- Health Nutr. 2013, 16(10), 1741–50.
26. Mengistu, G.; Moges, T.; Samuel, A.; Baye, K. Energy and nutrient intake of infants and young children in pastoralist communities of Ethiopia. *Nutrition* 2017, 41, 1–6.
  27. Ethiopian Public Health Institute (EPHI). Ethiopia National Food Consumption Survey, 2013; p.95.
  28. Herrador, Z.; Sordo, L.; Gadisa, E.; Moreno, J.; Nieto, J.; Benito, A.; et al. Cross-sectional study of malnutrition and associated factors among school aged children in rural and urban settings of fogera and libo kemkem districts, ethiopia. *PLoS One*. 2014, 9(9), e105880.
  29. Harika, R.; Faber, M.; Samuel, F.; Mulugeta, A.; Kimiywe, J.; Eilander, A. Are Low Intakes and Deficiencies in Iron, Vitamin A, Zinc, and Iodine of Public Health Concern in Ethiopian, Kenyan, Nigerian, and South African Children and Adolescents? *Food Nutr Bull*. 2017, 38 (3), 405–27.
  30. Amare, B.; Moges, B.; Moges, F.; Fantahun, B.; Admassu, M. Nutritional status and dietary intake of urban residents in Gondar , Northwest Ethiopia. *BMC Public Health* 2012, 12, 752.
  31. Hailelassie, K.; Mulugeta, A.; Girma, M. Feeding practices, nutritional status and associated factors of lactating women in Samre Woreda, South Eastern Zone of Tigray, Ethiopia. *Nutr. J*. 2013, 12(1), 1.
  32. Biza, Z.N. Food Taboos and Misconceptions Among Pregnant Women of Shashemene

- District, Ethiopia, 2012. *Sci.J. Public Heal.* 2015, 3(3), 410.
33. Zerfu, T.A.; Umeta, M.; Baye, K. Dietary habits, food taboos, and perceptions towards weight gain during pregnancy in Arsi, rural central Ethiopia: a qualitative cross-sectional study. *J. Health Popul. Nutr.* 2016, 35 (1), 22.
  34. Getnet, W, Aycheh W, Tessema T. Determinants of Food Taboos in the Pregnant Women of the Awabel District , East Gojjam Zone , Amhara Regional State in Ethiopia. *Adv Public Heal.* 2018;2018:6.
  35. Ethiopian Public Health Institute (EPHI). Ethiopian National Micronutrient Survey Report. Addis Ababa, Ethiopia, 2016; p.126.
  36. Herrador, Z.; Sordo, L.; Gadisa E, Buño A, Gómez-Rioja R, Iturzaeta, J.M.; et al. Micronutrient deficiencies and related factors in school-aged children in ethiopia: A cross-sectional study in Libo Kemkem and Fogera Districts, Amhara Regional State. *PLoS One* 2014, 9(12), 1–20.
  37. Lebso, M.; Anato, A.; Loha, E. Prevalence of anemia and associated factors among pregnant women in Southern Ethiopia: A community based cross-sectional study. *PLoS One* 2017, 12(12), 1–11.
  38. Gebremedhin, S.; Enquselassie, F.; Umeta, M. Prevalence of prenatal zinc deficiency and its association with socio-demographic, dietary and health care related factors in Rural Sidama, Southern Ethiopia: A cross-sectional study. *BMC Public Health* 2011, 11, 898.
  39. Gebreselassie, S.G.; Gase, F.E.; Deressa, M.U. Prevalence and Correlates of Prenatal

- Vitamin A Deficiency in Rural Sidama , Southern Ethiopia. *J. Heal. Popul. Nutr.* 2013, 31(2), 185–94.
40. Gebremedhin, S; Enquselassie, F.; Umeta, M. Prevalence and Correlates of Maternal Anemia in Rural Sidama, Southern Ethiopia. *Afr. J. Reprod. Health.* 2014, 18, 44–53.
  41. Herrmann, W.; Obeid, R. Utility and limitations of biochemical markers of vitamin B12 deficiency. *Eur. J. Clin. Invest.* 2013, 43, 3, 231–7.
  42. Hannibal, L.; Lysne, V.; Behringer, S. Biomarkers and Algorithms for the Diagnosis of Vitamin B 12 Deficiency. *Front. Mol. Biosci.* 2016, 3, 27.
  43. Kwok, T.; Cheng, G.; Lai, W.K.; Poon, P.; Woo, J.; Pang, C. P. Use of Fasting Urinary Methylmalonic Acid to Screen for Metabolic Vitamin B12 Deficiency in Older Persons. *Nutrition* 2004, 20, 764-768.
  44. Gebreegziabher T, Stoecker BJ. Vitamin d insufficiency in a sunshine-sufficient area: Southern ethiopia. *Food Nutr Bull.* 2013;34[4]:429–33
  45. Asayehu, T.T.; Lachat, C.; Henauf, S. D.; Gebreyesus, S.H. Dietary behaviour, food and nutrient intake of women do not change during pregnancy in Southern Ethiopia. *Matern Child Nutr.* 2017, 13, e12343.
  46. Tesfaye, B.; Sinclair, K.; Wuehler, S.E.; Moges, T.; De-regil, L.M.; Dickin, K.L. Applying international guidelines for calcium supplementation to prevent pre-eclampsia : simulation of recommended dosages suggests risk of excess intake in Ethiopia. *Public Health Nutr.* 2018, 22(3), 531–41.



# **Chapter 5**

## **General discussion**

## **5. General discussion**

In this chapter, the core findings on the dietary pattern and undernutrition of lactating women (underweight) and 6-23-months-old children (underweight, wasting and stunting) in relation with the Ethiopian Orthodox lent fasting and non-fasting periods, considering the fasting status of lactating mothers are discussed. The factors associated with undernutrition of the lactating mothers and their breastfeeding 6-23-months-old children are also dealt. Furthermore, the trend on estimated dietary energy and macro- and micronutrients intake and the prevalence of inadequacy of the lactating mothers and their 12-23-months-old children between/during fasting and non-fasting periods are discussed. Moreover, the development of CIMI app and the validation of the results in comparison with an established nutrition assessment software called NutriSurvey are discussed. Finally, the conclusion of the whole study is also conferred.

### **5.1. Undernutrition in Lactating Women and their 6-23-Months-Old Children**

#### **5.1.1. Prevalence of undernutrition in lactating women and their 6-23-months-old children**

In low and middle income countries, undernutrition in mothers and their children persists as an inescapable and devastating condition [1]. It is also the principal cause among the total morbidity, mortality and disability-adjusted-life-years. Maternal undernutrition is an independent factor of the nutritional status of their children. Earlier studies in different countries have shown that mothers who were short (<145 cm) and had low body mass index were linked with low birth weight (LBW) of their children, increased offspring mortality, underweight, stunting during the childhood, and these undernutrition problems appear later at the adulthood [2–5]. Likewise, studies elsewhere, in Ethiopia also confirmed that maternal undernutrition is associated with

child birth outcome, stunting and underweight [6–9]. The reason for some of the cases could be, as the woman is shorter, then the storing of protein and energy is expected to reduce and have smaller size of reproductive organs and pelvis diameter. These, could limit the intra-uterine growth of the fetus, and later the growth of an infant by reducing the quantity and quality of breast milk, as a result a stunted child [10]. In the present study, the prevalence of short maternal height (< 145cm) was 1.4% (Chapter 2, Table 2.4), which was lower than previous findings in Tigray region and national average, in Ethiopia [11–13]. Similarly, the prevalence of underweight in the lactating mothers was 32.6-32.8% in the study area, which was lower than results reported in earlier studies in Ethiopia [13–15]. But, alarmingly, this prevalence was very high (51%) in fasting lactating mothers compared to the 25% prevalence in non-fasting mothers (Chapter 2, Table 2.4). The later finding was comparable with results of earlier studies in lactating mothers living in mid-land agro-ecology of North Ethiopia [15]. At the same time, the average body mass index (BMI) values of the fasting mothers was lower compared to non-fasting group, which is consistent with the findings of studies conducted on Greek Orthodox Christians [16]. This could be related to the decreased number of meals, resulting from lower calorie intake in fasting people compared to non-fasting during fasting days [10,17].

In addition to the very high prevalence of underweight in fasting mothers found in our study, the prevalence of wasting, underweight and stunting were 4.0-4.8%, 11.7-15.7% and 31.6-33.7% in the 6-23-months-old children of the study area (Chapter 3, Table 3.2). Our findings were lower than earlier findings for the Tigray region and national prevalence's of the last Demographic and Health Survey (DHS) in Ethiopia. According to Ethiopian DHS (2016), the prevalence of stunting and underweight decreased from 58% and 41% in 2000 to 38% and 24%,

with only 2% decrease in wasting in 16 years' interval, at national level, which was the aggregate for the regions including Tigray [13]. These trends in reduction of undernutrition indicated that nutritional status of Ethiopia children is improving, although the existing prevalence is unacceptably high. The present study also showed that the average z-scores of stunting (LAZ), underweight (WAZ) and wasting (WLZ) of the children of the fasting mothers were significantly lower compared to the non-fasting group (Chapter 3, Table 3.5). Additionally, the average z-scores of WAZ of children of both fasting and non-fasting mothers showed improvement in non-fasting period after the lent fasting period (Chapter 3, Table 3.6). These indicated that fasting affect the nutritional status of women at lactation, and it also impaired the nutritional status of their breastfeeding children. Furthermore, during the long fasting period, the nutritional status of the children was also affected, regardless of their mother fasting status.

#### **5.1.2. Factors associated with undernutrition of lactating women and their 6-23-months-old children**

Maternal and child undernutrition is the problem in the community caused by several factors of multi-dimensional nature. According to UNICEF conceptual framework of undernutrition, the immediate causes are inadequate intake and disease condition of an individual, whereas, the long term causes are household food insecurity, inadequate care, unhealthy household environment and lack of health services and shortage of income leading to poverty at the household and community level, and the basic causes for undernutrition are closely linked with social, political and economic basis of the society [1,18]. Accordingly, lots of efforts have been exerted in separate ways for long time, but the outcome was far below the global targets by most of the countries. As a result, fighting undernutrition shifted its focus from the narrow approach of

undernutrition to multi-sectoral approach using multi-sectoral forces to synergize the efforts and accelerate the reduction of undernutrition in different countries and worldwide [19]. Therefore, nutrition is the core of the sustainable development goal (SDG), and it is a vital way for achieving the 12 SDGs out of the 17. Even, the remaining 5 also contribute for the improvement of nutrition worldwide [19]. In this regard, the government of Ethiopia has been doing a lot of activities using a multi-sectoral approach by integrating different national sector ministries. But, undernutrition as a public health problem still remains, due to inadequate commitment and lack of conducive structure in government system. As a result, the SEQOTA declaration, which needs a strong and high level commitment of government, with a goal of ending undernutrition by 2030, under the National Nutrition Program II was launched [19,20]. This goal is planned to be achieved through integrating community development in agriculture, health, nutrition, education, water, sanitation and hygiene and social protection activities [19]. However, our findings urged the involvement of religious institution in the implementation of multi-sectoral approach to accelerate reduction of undernutrition and achieve the goal set in 2030. For example, about one-third of lactating mothers included in our study fasted during their pregnancy and lactation periods (Chapter 2, Table 2.2), which is slightly lower than earlier finding (38%) in pregnant mothers of Shashemene district in Ethiopia [21]. Mothers who were fasting during their pregnancy and lactation periods in this study had 1.7 and 2.9 times more odds to be underweight than respective none fasting groups during the same periods, respectively (Chapter 2, Table 2.3). Likewise, children of fasted mothers during the pregnancy and lactation period were also more exposed to undernutrition (stunting, underweight, and wasting (only during lactation)) than those children whose mothers did not fast in the same periods (Chapter 3, Table 3.3). These show that

fasting during pregnancy and lactation periods of the mothers affects, not only the nutritional status of mothers, but also the health and nutritional status of their fetus and/or breastfed child [10,22]. Earlier studies in sub-Saharan African including Ethiopia and Asian countries confirmed that younger mothers and children between 13-23-months-old had a higher probability of being underweight, and underweight and stunted compared to older and children between 6-12 months of age, respectively [13,23–26]. The later (underweight and stunted) might be related with the contamination of complementary food coupled with poor nutritionally quality, which is mainly cereal based food in Tigray region, in Ethiopia [27]. Similarly, mothers whose children were 13-18 months old were more likely to be underweight compared to those who had children between 6-12 months old, which is in line with the result found in a study conducted by Hailselassie and his colleagues [12]. The possible reasons could be, as the child becomes older his/her nutritional requirements increases, which is also supported by the extra energy expenditure related with child care, and no increase in the food intake, which is even aggravated by the decreased intake of the mother [10]. Mothers who experienced any illness in the last four weeks preceding our lent fasting survey had higher odds of being underweight compared to healthy lactating mothers, which is in line with another study conducted in Limu district of Southern Ethiopia [28]. Related with this, about one-third of the lactating mothers were living in households with non-improved water source. This was also associated with increased odds of underweight compared to mothers living in households with an improved water source for the household consumption. This is basically due to frequent illness related to water borne diseases and contamination. Illness usually affects the appetite of an individual and decreases food intake and also absorption of nutrients. It may also change the metabolism, and as a result increase the nutrient requirements

[10]. Mothers who were living with relatives' /parents grandfather was the decision maker for the income, and received aid in the form of food and/or cash and/or in kind before our fasting survey and were more likely to be underweight than mothers living in households, the income decision was her husband and not had any aid experience. The former could be due to the fact that sharing a household income to more family members or due to decreased working power and resulted in lower income, which is related with loss of father or divorce or it might be related to living in households headed by the grandfathers. This affects usually young mothers, which is associated with underweight in this study. Furthermore, it could be related to an extended family, and given the allocation of food priority given to the children, which could lead to reducing of the serving size of the meal or skipping the meal by mothers [29]. Households experiencing aid are most of the time food insecure and don't have farming land or sufficient household income, which affect quality of food of household members, especially children and women, could be the reason for the presence of more underweight women in this households. Mothers who were living in households who did not own chicken during the survey were more exposed to underweight compared to their counterparts. Earlier study in three African countries indicated that households owning livestock had lower prevalence of child stunting than those without livestock [30]. Therefore, in general promotion of animal husbandry could improve the low consumption of animal source food in the study area.

In addition to these, studies in other low and mid-income countries and Ethiopia showed that maternal educational status was a predictor for childhood wasting [13,25,31–34]. In line with this, our study confirmed that maternal illiteracy was associated with childhood wasting and

stunting. Therefore, improving the education status of the mother is one of the important interventions for reduction of child undernutrition in the study district.

Maternal occupation was also one of the predictor variables of child underweight in our study. Compared to children whose mothers were housewives, children born of farmer mothers were more exposed to underweight. This could be due to high work load and absence from home, which leads to provision of less time for child care. Additionally, expending more energy on farming activity might cause poor appetite of mothers reduce the quantity and quality of breast milk produced. Beside these, children who were living in households without toilet and who were not fed colostrum after their birth were more likely to be stunted compared to those who were living in households owning toilet and did not take colostrum. The potential reason for the first case could be the infection related with pre-lacteal foods, which was associated with child stunting in a study conducted in Ethiopia [35]. Due to absence of toilet in households, open defecation increases the risk of frequent diarrhea episodes, this affects the nutritional status of children [23]. For these, promoting health education by relevant institutions on the importance of colostrum feeding during the antenatal care should be provided. Strengthening the construction of low cost latrine through the existing health extension package should be continued.

## **5.2. Feeding Practices and Nutrient Intake of Lactating Women and their 6-23-Months-Old Children**

Ensuring the recommended diversification of diet and number of meals in a day to day life of an individual is an important component for supplying energy, macro- and micronutrients in required amount to fight against maternal and child undernutrition [1,36]. In this regard, mothers are recommended to take at least two additional meals during their lactation period, from at least



five defined food groups out of ten, in a day [37,38]. Similarly, if a 6-23-months-old child eats from at least four food groups in his/her daily meals, then he/she fulfills the minimum diet diversity (MDD). This should be obtained from two or more meals (MMF) if he/she is 6–8 months old and three or more for 9–23 months for the breastfeeding children. Therefore, the minimum acceptable diet (MAD), a combination of both the MDD and MMF criterion, leads to appropriate growth and development of infant and young children [39].

In this survey, two-thirds of the mothers (65.4%) did not change the food intake during their lactation period. Even worse, one-third of lactating mothers ate less than three times during fasting period, which is lower than the expected daily three meals of a normal adult in real context. Comparable finding was observed in a study conducted in lactating women of Samre district in North Ethiopia [12]. However, the prevalence of those mothers who had less than three meals in a day was reduced to 8.4% in the non-fasting period in this survey. But, the number of meals eaten both by the fasting and non-fasting mothers were improved in the non-fasting period after the two months of lent fasting. Therefore, further study should be done for understanding how the non-fasting mothers` were affected in this regard. Likewise, three out of four children from fasting and non-fasting mothers ate the minimum number of meals recommended, during the fasting and non-fasting periods, which was in line with studies conducted in the northern and southern part of Ethiopia [40,41].

In this study, most of the mothers had a diet from a maximum of three food groups during the fasting period, regardless of their fasting status. However, the consumption of diet from more diversified food groups were increased during the non-fasting period (Chapter 2, Table 2.4). This is because the consumption of animal source foods (ASFs) and the number of meals eaten during

the non-fasting period was increased. As a result, the energy, macro- and micronutrients intake were increased, although the prevalence of inadequacy of these energy and micronutrients remain higher, except iron, magnesium, vitamin B1 and niacin. However, almost all the lactating mothers and their 12-23-months-old children included in the study did not take vitamin B12, vitamin D and calcium, in both the study periods (Chapter 4, Table 4.3 and 4.4). Earlier national study in Ethiopia using blood biomarkers found that about 20% women at reproductive age were deficient of vitamin B12 [42]. However, this high discrepancy could be due to the fact that serum total vitamin B12 used for assessment can give less information to identify vitamin B12 deficiency [43,44]. Further study should be done using methyl malonic acid in urine or Holotranscobalamin in blood samples, to get more reliable information to be able to design appropriate preventive or intervention measures [43,45]. In line with our finding in the present study, there were no pregnant women ever consumed vitamin D-rich foods, fortified foods or dietary supplements, which might have resulted in vitamin D deficiency, as a serious problem, in spite of the fact that Ethiopia is located near the Equator [46]. Beside these, the highest prevalence of dietary calcium inadequacy were found in studies conducted in Gondar and Butajira districts, and at national (89-96%) level in Ethiopia [47–49]. Therefore, nutrition sensitive agriculture in both plant and livestock production should be promoted and implemented in sustainable manner, in addition to the implementation of proposed universal calcium supplementation [49].

In the present study, only 2.5% and 6.9% of the 6-23-months-old children consumed complementary foods prepared from four and more diversified food groups, in the fasting and non-fasting periods, respectively, which was lower than those found in a study conducted in the

same age group of children of Orthodox mothers during the fasting period, in Gojam district of Ethiopia [50]. It was also lower than the results of studies conducted in India, Bangladesh and Vietnam, which were 38%, 48.4% and 83.2%, respectively. Furthermore, only 2.3-6.7% of the 6–23-month-old children met the minimum acceptable diet criteria in the study population, which was in line with findings for the Tigray region and national figure, in the DHS 2016 report for Ethiopia [13]. However, it was lower than that found on the 6-23-months-old children of Orthodox religion follower (8.6%) in North west Ethiopia [51]. Furthermore, the proportion of 6–23-month-old children of fasting mothers who fulfilled the criteria for MAD was lower compared to children of non-fasting mothers. It was also lower during the fasting compared to non-fasting period, which implied that the inadequate dietary micronutrients intakes were prevalent in the fasting period. In support of these, the average dietary intake of energy, protein and all micronutrients included in this study were higher in non-fasting compared to fasting period, despite the average intake of these were below the estimated need of the breastfeeding children from the complementary food and high prevalence of energy and nutrient inadequacies in the study population. For example, the prevalence of inadequate dietary intakes ( $<2/3$  RNI) of zinc, vitamin C, energy and niacin was very high ( $>50\%$ ) in the 12-23-months-old children, followed by protein, iron, vitamin A and vitamin B1 (20-50%), respectively. Studies conducted in Amhara and Tigray regions, and at national level in Ethiopia confirmed that zinc and vitamin A deficiencies are high in children [52,53]. Another study conducted in three other African countries and Ethiopia also revealed that anemia, vitamin A and zinc deficiencies are public health problems in these countries [54]. The prevalence of dietary inadequacy ( $2/3$  RNI) of Vitamin C was high in both the children and lactating mothers and almost steady in both study

periods. This can indicate that low consumption of vitamin C rich foods is the chronic problem in the study area, which might be related with the low consumption of fruits, animal source foods including dairy products, root crops like potato, anchote in the study area [10,22]. Therefore, consumption of vitamin C rich foods should be promoted to improve the bioavailability of minerals. This alone can't be enough, unless a traditional processing techniques is used to improve the bioavailability of iron, zinc and calcium, in a population, in which cereals and legumes are the main staple foods and a source of energy and most of the nutrients needed for an individual (Chapter 2, Table 2.4). Beside these, the proportion of mothers who consumed more cups of coffee was higher in the fasting than non-fasting period, in the whole study population, and these were due to higher proportion of the non-fasting mothers included in this study who took significantly more coffee in the fasting than non-fasting periods. However, the reason for less consumption for the fasting mothers could be related to some Ethiopian Orthodox religion monarchist, who preached to followers not to consume coffee during fasting periods. Therefore, moderate intake of coffee, coupled with discouraging intake close to meals to be eaten should be done, otherwise, it could also reduce their bioavailability, leading to increased risk of micronutrient deficiencies, especially of iron, calcium, and zinc [55].

### **5.3. CIMI: A Promising Dietary Assessment Application for Rural Ethiopian Population**

Assessing the dietary pattern of an individual, and a community at large, is the fundamental for setting an appropriate prevention and intervention activities in a given country. It can also help to identify, both poor and good dietary habits, and identify the extent of risk and deficiency for specific nutrients in a certain community, by the practitioners, researchers and development agencies [56]. As a result, qualitative and quantitative dietary intake data have been collected

using dietary assessment techniques, which have been developed and implemented globally. However, most of the quantitative dietary assessment methods are time consuming, need an experienced and well trained data collectors, and an expert to calculate the energy and nutrient intakes using software before identifying extent of the nutritional problem in the study population. Thus, quantitative dietary assessment methods are less applicable in developing countries like Ethiopia, where more than three-fourth of the population are living in rural poor settings, which most often leads to less availability of quantitative dietary data [57]. In this study, a web-based nutritional assessment application called Calculator for Inadequate Micronutrient Intake (CIMI), which is simple, easy-to-use and informative and consider the 93 commonly consumed Ethiopian foods was adapted for Ethiopian population and validated against internationally established nutrition assessment software called NutriSurvey (NS). Accordingly, the average intake values of energy, protein and most nutrients produced by CIMI and NS were almost similar, for the lactating women and their 12-23-months-old children, regardless of the two study periods. Exceptions were observed in vitamin A in both women and children, and vitamin C only in the lactating women, which were higher in NS and CIMI, compared to the results produced by CIMI and NS, respectively. The findings on the vitamin A was in line with the result found in the validation study of Indonesian CIMI [58]. The reason for the vitamin A in our case was, the butter and oil food group in CIMI has got a lower average vitamin A content compared to the fortified palm oil consumed by almost all of the study households surveyed during the fasting and non-fasting periods. The consumption of different oil types in the district depends on the availability of oil types through the cooperatives provided under the massive import by the subsidy of the Ethiopian government and the distribution through the aid in the

district. To solve the problem related with vitamin A, deselection of the unconsumed oil type in the oil and butter food will be important in future CIMI use. The increased result produced in the vitamin C was due to the greater share of the tomato (43.8%) for the 'other vegetables' food group in the CIMI and its relatively high vitamin C content, but less consumed compared to the highly consumed cabbage by the adults in the study area. The average intakes of protein, iron and calcium produced by CIMI were also higher than NS in both the study groups, which was related with the consumption of high amount of peas, despite its share in the 'beans' food group is small (12.4%). The other reason was the higher contents of protein, iron and calcium in the food items compared to peas, which created the 'beans' food group. Furthermore, if the amount to be set up between the two sliders gets larger, then the consumption is increasing in most of the food groups created the CIMI. The selection of the lower or upper, where the amount consumed in the group fails depends on the nearest amount in the setup of the sliders, which also affects the beans group. Therefore, if the amount approaches to the upper slider, then automatically the upper slider will be selected and this will lead to overestimation of intake of specific nutrients compared to NS, otherwise underestimate for lower slider selection. Similarly, the average intake results produced for magnesium in the pepper and vitamin B1 and egg food groups were overestimated by selecting the upper slider during the data entry of CIMI in the higher proportion of an individual. The vitamin B6 was also slightly higher in results produced by CIMI, which was due to the fact that Ethiopian kale, which was less consumed in the study area, but contributed much (58.4%) to the 'green leaves' food group overestimated the average intake in the study groups, regardless of the study periods.

The results in the present study also showed that the proportion of lactating women and the 12-23-months old children who were identified as having inadequate intake of energy and most nutrients ( $<2/3$  RNI) were similar by CIMI and NS, in the study periods. Like the average intake for the vitamin A of the lactating women and the 12-23-months-old children, and vitamin C for the lactating women group, the prevalence identified by the CIMI and NS was also variable, in both the study periods. The reasons for these discrepancies are also similar with those earlier stated for the average differences for vitamin A and vitamin C intakes in our study. Additionally, according to criteria set by Lambert and her colleagues (2018), the results of vitamin A for both of the study groups, magnesium for the 12-23-months-old children and vitamin B6 for the lactating women, were produced by CIMI had moderate accuracy compared to NS. Similarly, the accuracy of results produced by CIMI of the vitamin B6 in the 12-23-months-old children and vitamin C for the lactating women were low. The reason for the vitamin B6 was due to the actual small average values produced, so that the small mean differences in few individuals between CIMI and NS dramatically increased the mean difference %, which is the drawback of the accuracy classification method we used. Whereas, the vitamin C was due to the reasons mentioned for the discrepancies observed in the average results produced. Despite these, the intake of energy, protein and most micronutrients, for both of the study groups, in both study periods were calculated either in good or very high accuracy by CIMI. Beside these, strong correlations were found between CIMI and NS results in all the nutrients and energy for the 12-23-months-old children ( $R=0.741-0.956$ ) and lactating women ( $R=0.779-0.920$ ) (Chapter 4, Figure 4.1 and 4.2). Our results found in the correlation analysis of results produced by NS and CIMI for energy, protein, iron, zinc and vitamin A in both the study groups were higher than

previous study in Indonesian children and women, except that of energy for lactating women. This discrepancy could be related to the smaller sample size of the study participants compared to this study. Additionally, the correlation coefficients of all nutrients, except energy in our study were higher than those found in a CIMI trial conducted on Tanzanian women [59]. Thus, CIMI developed for rural Ethiopian population calculates the average nutrient intake accurately, and identifies the inadequate micronutrient intake of individuals using FAO/WHO recommendation for nutrient intake (RNI).

Unlike other dietary assessment methods, CIMI provides result immediately after the interview is completed, which allows the enumerator to provide direct feedback to the interviewee to improve the dietary habits considering the socio-cultural and economic status of the community, and the potential crops available in the farm and market of the district. Currently, most of the dietary quality information have been collected using the qualitative method called diet diversity score, which is a proxy indicator at the individual and household levels, in most of the developing countries, as most of the quantitative assessment methods are not feasible in these countries [38]. In this technique, the principal focus is assessing the diversity of the standardized food groups consumed in the diet and quantifying in to the score to indirectly predict the micronutrients adequacy in general, using an established cutoff points mentioned earlier. Therefore, the information that can be produced from this method is limited unless an alternative method is used to identify specific nutrient deficiency in the given community. Considering these, CIMI can replace the diet diversity, by producing much more results on the diet quality of an individual, and solving the problems that can be encountered by diet diversity assessment method. In situations where the diet diversity and other quantitative dietary assessment methods



are not appropriate, CIMI can be also used in cross-sectional and/or longitudinal/intervention studies. Cumulative analysis of individual results, can be used in mapping nutrition situation of a given district or country using the GPS information to be produced during the survey carried out by CIMI. Therefore, appropriate and specific prevention or intervention activities can be planned and done timely. Therefore, our results indicate that, the CIMI developed for Ethiopian population using the commonly consumed Ethiopian foods are applicable in rural setting, where 85% of the population is living and suffering from multiple malnutrition problems. In general, the use of CIMI is wider, which covers from the simple cross-sectional study in a certain area to the complex longitudinal study at the national level in Ethiopia.

#### **5.4. Conclusion**

Based on our findings on the anthropometric measurements of lactating women and their 6-23-months-old children, maternal and child undernutrition is a serious public health problem in the study district. But, it was lower than the prevalences of undernutrition in children and women at the reproductive age group (15-49 years) reported in the Ethiopian Demographic and Health Survey (2016) for Tigray region, where this study was conducted. Therefore, these indicated that the nutritional status of women and their children in the Tigray region was improving, despite the unacceptable higher prevalences of undernutrition in fasting mothers (underweight) and children (stunting, underweight and wasting) born of these fasting mothers and the study population especially during the Ethiopian Orthodox lent fasting period. Mother fasting status during pregnancy and lactation periods of the indexed child was amongst the predictor variables of the undernutrition for the lactating women and their 6-23-months-old children in the study population. Thus, the existing maternal and child nutrition intervention activities should consider

religious fasting periods. Mothers who were young, had sickness within four weeks preceding the fasting period, and children between 13 and 18 months of age were more exposed to maternal underweight. Mothers who were living in the households whose decision makers were grandfathers, received aid, not owning chickens and had no access to water were also associated with maternal underweight. Therefore, households with younger mothers and non-improved water sources should get priority in nutritional intervention. Furthermore, promoting chicken husbandry, and improving the WASH in the community may also improve the low consumption of animal source foods and reduce maternal undernutrition in the study district. Likewise, maternal education and occupational status, child age, colostrum intake status, in addition to toilet presence in the household where the child was living associated with child undernutrition in the study district. Therefore, improving the maternal education, reducing the work burden in farming activities and/or supporting of mothers in non-farm income generating nearby by their living residence, household latrine coverage and breastfeeding of colostrum should be promoted and strengthened. The number of meals eaten, both by the lactating women and their 6-23-months-old children are below the recommendation, from the less diversified food groups. These food groups are mainly from the cereals and legumes, which are known to have high amount of phytic acid and tannin which limit the absorption of minerals and protein by chelating with it. In addition to the low consumption of all other food groups, by both the study groups in the study population, the proportion of fasting lactating women and their 6-23-months-old children who consumed animal source foods (dairy, meat, poultry and fish) were very low compared to the lactating women and their 6-23-months-old in the non-fasting group. Similarly, the consumption of animal source foods was affected during the lent fasting than non-fasting period at the study

population level. This sub-optimal dietary pattern was synergized with more cup of coffee consumption by the lactating women, and even worst in the lent fasting period. Furthermore, we also concluded that the prevalence of inadequate intakes ( $<2/3$  RNI) of dietary energy and most micronutrients, except protein and magnesium are high in the 12-23-months-old children. Similarly, the proportion of lactating women who did not take adequate amount of vitamin A, calcium, vitamin B6, vitamin B12, pentatonic acid and vitamin C were also high. These high prevalences of inadequacy was more affected during the lent fasting period, in both the young children and lactating women. Thus, activities which will improve the diet diversity by incorporating the nuts and seeds, dark green leafy vegetables, vitamin A and C rich fruits and animal source products, and increasing the meal frequency, for both the lactating women and children should be done urgently. Besides these, use of traditional processing techniques like dehulling/dehusking, soaking and germination, and slightly roasting are crucial during preparation of food for the nutrient needy groups like lactating and pregnant women, and their young children. Additionally, moderate intake of coffee after some hour interval of meal to be eaten should be promoted to increase the absorption of non-heme iron from plant source foods. Shockingly, the intake of the vitamin B12, vitamin D and calcium by almost all the lactating women and their 12-23-months-old children were below the recommended intake ( $2/3$  RNI) from their diet in the 24-hr preceding the fasting and non-fasting surveys. Thus, implementation of the proposed universal calcium supplementation in Ethiopia, in addition to urgent investigation and appropriate other public health nutrition activities should be done in the district. In general, the activities which will focus on the dietary improvement (quality and quantity) should get more priority, and special attention should be given for fasting periods.

We also concluded that, the Calculator for Inadequate Micronutrient Intake (CIMI) adapted using commonly consumed Ethiopian foods for rural Ethiopian population estimates the average nutrient intake accurately. CIMI also identifies the individual inadequate intake of energy, macro- and micronutrients using the FAO/WHO recommended nutrient intake (RNI) immediately after the interview. Therefore, this feature enables the interviewers to give immediate feedback and suggest context based dietary improvements for the interviewee. CIMI can also replace the DDS by solving the problem encounter by this technique and producing more quantitative results to design specific nutrition intervention. It can be also used in cross-sectional and/or longitudinal and/or intervention studies, where the DDS and other quantitative dietary assessment techniques are not appropriate. In general, CIMI can be used in Ethiopia, as a simple dietary assessment tool for nutrition and related researchers, policy makers, implementers and evaluators.

## **5.5. References**

1. Black RE, Allen LH, Bhutta ZA, Caulfield LE, de Onis M, Ezzati M, et al. Maternal and child undernutrition: global and regional exposures and health consequences. *Lancet*. 2008;371[9608]:243–60.
2. Addo OY, Stein AD, Fall CH, Gigante DP, Guntupalli AM, Horta BL, et al. Maternal height and child growth patterns. *J Pediatr* [Internet]. 2013;163[2]:549–554.e1. Available from: <http://dx.doi.org/10.1016/j.jpeds.2013.02.002>
3. Özaltın E, Hill K, Subramanian S V. Association of Maternal Stature With Offspring Mortality, Underweight, and Stunting in Low- to Middle-Income Countries. *JAMA*. 2011;303[15]:1507–16.

4. Silva D e, Valente A, Borges A, Almeida F, Neves E, Afonso C. Relationship between the mothers ' nutritional status with that of a child population from São Tomé Príncipe , “ Africa .” *Rev Bras Saúde Matern Infant*, Recife. 2017;17[2]:327–35.
5. Felisbino-mendes M, Villamor E, Velasquez-melendez G. Association of maternal and child nutritional status in Brazil : A population based cross-sectional study. *PLoS One*. 2014;9[1]:e87486.
6. Negash C, Whiting SJ, Henry CJ, Belachew T, Hailemariam TG. Association between maternal and child nutritional status in Hula, rural Southern Ethiopia: A cross sectional study. *PLoS One*. 2015;10[11].
7. Endalamaw A, Engeda EH, Ekubagewargies DT, Belay GM. Low birth weight and its associated factors in Ethiopia : a systematic review and meta- analysis. 2018;6:1–12.
8. Bililign N, Legesse M, Akibu M. *iMedPub Journals A Review of Low Birth Weight in Ethiopia : Socio-Demographic and Obstetric Risk Factors Abstract Magnitude of low birth weight*. 2018;1–5.
9. Gebremedhin M, Ambaw F, Admassu E, Berhane H. Maternal associated factors of low birth weight : a hospital based cross-sectional mixed study in Tigray , Northern Ethiopia. *BMC Pregnancy Childbirth* [Internet]. 2015;15[22]:1–8. Available from: <http://dx.doi.org/10.1186/s12884-015-0658-1>
10. Desalegn BB, Lambert C, Riedel S, Negese T, Biesalski HK. Ethiopian Orthodox Fasting and Lactating Mothers : Longitudinal Study on Dietary Pattern and Nutritional Status in

- Rural Tigray , Ethiopia. *Int J Environ Res Public Heal*. 2018;15[1767]:1–20.
11. Sitotaw IK, Kiday H, Adama Y. Comparison of nutritional status and associated factors of lactating women between lowland and highland communities of District Raya , Alamata ,. *BMC Nutr*. 2017;3[61]:1–12.
  12. Hailelassie K, Mulugeta A, Girma M. Feeding practices, nutritional status and associated factors of lactating women in Samre Woreda, South Eastern Zone of Tigray, Ethiopia. *Nutr J [Internet]*. 2013;12[1]:1.
  13. CSA. Ethiopia Demographic and Health Survey 2016. Addis Ababa; 2017.
  14. Alemayehu M, Argaw A, Geberemariam A. Factors Associated with Malnutrition among Lactating Women in Subsistence Farming Households from Dedo and Seqa-Chekorsa Districts, Jimma Zone, 2014. *Dev Ctry Stud [Internet]*. 2015;5[21]:114–22. Available from: <http://www.iiste.org/Journals/index.php/DCS/article/view/27469>
  15. Roba KT, O’Connor TP, Belachew T, O’Brien NM. Seasonal variation in nutritional status and anemia among lactating mothers in two agro-ecological zones of rural Ethiopia: A longitudinal study. *Nutrition [Internet]*. 2015;31[10]:1213–8. Available from: <http://dx.doi.org/10.1016/j.nut.2015.03.007>
  16. Sarri KO, Tzanakis NE, Linardakis MK, Mamalakis GD, Kafatos AG. Effects of Greek orthodox christian church fasting on serum lipids and obesity. *BMC Public Health*. 2003;8:1–8.
  17. Sarri KO, Tzanakis NE, Linardakis MK, Mamalakis GD, Kafatos AG. Effects of Greek

- orthodox christian church fasting on serum lipids and obesity. BMC Public Health. 2003;3[16].
18. UNICEF. The status of the world's children. Geneva, Switzerland: Oxford University Press; 1998. 135 p.
  19. FMOH. National Nutrition Program 2016-2020. Addis Ababa, Ethiopia; 2016.
  20. Federal Democratic Republic of Ethiopia [FDRE]. SEQOTA declaration implementation plan [ 2016 – 2030]: Summary programme approach document. Addis Ababa, Ethiopia; 2016.
  21. Biza Zepro N. Food Taboos and Misconceptions Among Pregnant Women of Shashemene District, Ethiopia, 2012. Sci J Public Heal [Internet]. 2015;3[3]:410. Available from: <http://www.sciencepublishinggroup.com/journal/paperinfo.aspx?journalid=251&doi=10.11648/j.sjph.20150303.27>
  22. Desalegn B, Lambert C, Riedel S, Negese T, Biesalski H. Feeding Practices and Undernutrition in 6–23-Month-Old Children of Orthodox Christian Mothers in Rural Tigray, Ethiopia: Longitudinal Study. Nutrients [Internet]. 2019;11[1]:138. Available from: <http://www.mdpi.com/2072-6643/11/1/138>
  23. Akombi BJ, Agho KE, Hall JJ, Wali N, Renzaho AMN, Merom D. Stunting, wasting and underweight in Sub-Saharan Africa: A systematic review. Int J Environ Res Public Health. 2017;14[8]:1–18.
  24. Alemayehu M, Tinsae F, Hailelassie K, Seid O, Gebregziabher G, Yebyo H.

- Undernutrition status and associated factors in under-5 children, in Tigray, Northern Ethiopia. *Nutrition* [Internet]. 2015;31[7–8]:964–70. Available from: <http://dx.doi.org/10.1016/j.nut.2015.01.013>
25. CSA. Ethiopia Demographic and Health Survey 2011. Addis Ababa; 2012.
  26. Bhandari S, Sayami JT, Thapa P, Sayami M, Kandel BP, Banjara MR. Dietary intake patterns and nutritional status of women of reproductive age in Nepal: Findings from a health survey. *Arch Public Heal* [Internet]. 2016;74[1]:1–11. Available from: <http://dx.doi.org/10.1186/s13690-016-0114-3>
  27. Mulugeta A, Hagos F, Kruseman G, Linderhof B, Stoecker B, Abraha Z, et al. Factors Contributing to Child Malnutrition in Tigray , Northern Ethiopia. *East Afr Med J*. 2010;87[6]:248–54.
  28. Demissie T, Mekonen Y, Haider J. Agroecological comparison of leveles and correlates of nutritional status of women. *Ethiop J Heal Dev*. 2003;17[3]:189–96.
  29. Wu X, Li L. Family size and maternal health: Evidence from the One-Child policy in China. *J Popul Econ*. 2012;25[4]:1341–64.
  30. Mosites EM, Rabinowitz PM, Thumbi SM, Montgomery JM. The Relationship between Livestock Ownership and Child Stunting in Three Countries in Eastern Africa Using National Survey Data. *PLoS One*. 2015;10[9]:1–10.
  31. Bbaale E. Maternal education and child nutritional status : evidence from Uganda. *African J Econ Manag Stud*. 2014;5[1]:52–74.



32. Makoka D, Masibo PK. Is there a threshold level of maternal education sufficient to reduce child undernutrition ? Evidence from Malawi , Tanzania and Zimbabwe. *BMC Pediatr* [Internet]. 2015;15:1–10. Available from: <http://dx.doi.org/10.1186/s12887-015-0406-8>
33. Nyaruhucha CNM, Msuya JM, Mamiro PS, Kerengi AJ. Nutritional status and feeding practices of under-five children in Simanjiro District , Tanzania. *Tanzan Health Res Bull*. 2006;8[3]:162–7.
34. Vollmer S, Bommer C, Krishna A, Harttgen K, Subramanian S V. The association of parental education with childhood undernutrition in low- and middle-income countries : comparing the role of paternal and maternal education. *Int J Epidemiol*. 2017;46[August 2016]:312–23.
35. Fentahun N, Belachew T, Lachat C. Determinants and morbidities of multiple anthropometric deficits in southwest rural Ethiopia. *Nutrition* [Internet]. 2016;32[11–12]:1243–9. Available from: <http://dx.doi.org/10.1016/j.nut.2016.03.023>
36. WHO. Complementary feeding of young children in developing countries: a review of current scientific knowledge. Geneva, Switzerland; 1998. p. 248.
37. Essential Nutrition Action [ENA]. Key behaviours for optimal breastfeeding , complementary feeding and maternal nutrition at critical stages in the life cycle of women and children. 2008.
38. FAO and FHI 360. Minimum Dietary Diversity for Women A Guide to Measurement.

Rome; 2016. 82 p.

39. WHO. Indicators for assessing infant and young child feeding practices. Part 3 Ctry Profiles. 2010;1–47.
40. Belew AK, Ali BM, Abebe Z, Dachew BA. Dietary diversity and meal frequency among infant and young children: A community based study. *Ital J Pediatr*. 2017;43[1]:6–15.
41. Dangura D, Gebremedhin S. Dietary diversity and associated factors among children 6-23 months of age in Gorche district, Southern Ethiopia: Cross-sectional study. *BMC Pediatr* [Internet]. 2017;17[1]:1–7. Available from: <http://dx.doi.org/10.1186/s12887-016-0764-x>
42. Ethiopian National Micronutrient Survey Report. Addis Ababa, Ethiopia; 2016.
43. Hannibal L, Lysne V, Behringer S. Biomarkers and Algorithms for the Diagnosis of Vitamin B 12 Deficiency. *Front. Mol. Biosci*. 2016;3(27). doi: 10.3389/fmolb.2016.00027
44. Herrmann W, Obeid R. Utility and limitations of biochemical markers of vitamin B12 deficiency. *Eur J Clin Invest*. 2013;43[3]:231–7.
45. Kwok T, Cheng G, Lai WK, Poon P, Woo J, Pang CP. Use of Fasting Urinary Methylmalonic Acid to Screen for Metabolic Vitamin B12 Deficiency in Older Persons. *Nutrition*. 2004;20:1–4.
46. Gebreegziabher T, Stoecker BJ. Vitamin d insufficiency in a sunshine-sufficient area: Southern ethiopia. *Food Nutr Bull*. 2013;34[4]:429–33.
47. Amare B, Moges B, Moges F, Fantahun B, Admassu M. Nutritional status and dietary intake of urban residents in Gondar , Northwest Ethiopia. *BMC Public Health*.

2012;12[752].

48. Asayehu TT, Lachat C, Henauw S De, Gebreyesus SH. Dietary behaviour, food and nutrient intake of women do not change during pregnancy in Southern Ethiopia. *Matern Child Nutr.* 2017; 13, e12343.
49. Tesfaye B, Sinclair K, Wuehler SE, Moges T, De-regil LM, Dickin KL. Applying international guidelines for calcium supplementation to prevent pre-eclampsia : simulation of recommended dosages suggests risk of excess intake in Ethiopia. *Public Health Nutr.* 2018;22[3]:531–41.
50. Kumera G, Tsedal E, Ayana M. Dietary diversity and associated factors among children of Orthodox Christian mothers/caregivers during the fasting season in Dejen district, north west Ethiopia. *Nutr Metab [Lond].* 2018;15[16].
51. Mulat E, Alem G, Woyraw W, Temesgen H. Uptake of minimum acceptable diet among children aged 6 – 23 months in orthodox religion followers during fasting season in rural area , DEMBECHA , north West Ethiopia. *BMC Nutr.* 2019;5[18]:1–10.
52. Herrador Z, Sordo L, Gadisa E, Buño A, Gómez-Rioja R, Iturzaeta JM, et al. Micronutrient deficiencies and related factors in school-aged children in ethiopia: A cross-sectional study in Libo Kemkem and Fogera Districts, Amhara Regional State. *PLoS One.* 2014;9[12]:1–20.
53. Ayana G, Moges T, Samuel A, Asefa T, Eshetu S, Kebede A. Dietary zinc intake and its determinants among Ethiopian children 6–35 months of age. *BMC Nutr [Internet].*

- 2018;4[1]:30. Available from:  
<https://bmcnutr.biomedcentral.com/articles/10.1186/s40795-018-0237-8>
54. Harika R, Faber M, Samuel F, Mulugeta A, Kimiywe J, Eilander A. Are Low Intakes and Deficiencies in Iron, Vitamin A, Zinc, and Iodine of Public Health Concern in Ethiopian, Kenyan, Nigerian, and South African Children and Adolescents? *Food Nutr Bull*. 2017;38[3]:405–27.
  55. Gebremedhin S. Prevalence of prenatal zinc deficiency and its association with socio-demographic, dietary and health care related factors in Rural Sidama, Southern Ethiopia: A cross-sectional study. *BMC Public Health* [Internet]. 2011;11[898]. Available from: <http://bmcpublichealth.biomedcentral.com/articles/10.1186/1471-2458-11-898>
  56. Lyu L, Hsu Y, Chen H, Lo C, Lin C. Taiwanese Journal of Obstetrics & Gynecology Comparisons of four dietary assessment methods during pregnancy in Taiwanese women. *Taiwanese Journal of Obstetrics & Gynecology* [Internet]. 2014;53[2]:162–9. Available from: <http://dx.doi.org/10.1016/j.tjog.2014.04.007>
  57. Development initiatives. 2018 Global nutrition report: Shining a light to spur action on nutrition [Internet]. Bristol, UK; 2018. 261 p. Available from: <http://ebrary.ifpri.org/cdm/ref/collection/p15738coll2/id/130399%5Cnhttps://www.ifpri.org/publication/hidden-hunger-approaches-tackling-micronutrient-deficiencies>
  58. Jati IRAP, Widmer C, Purwestri RC, Wirawan NN, Gola U, Lambert C, et al. Design and validation of a program to identify inadequate intake of iron, zinc, and vitamin A. *Nutrition* [Internet]. 2014;30[11–12]:1310–7. Available from:

<http://dx.doi.org/10.1016/j.nut.2014.03.015>

59. Lambert C, Eleraky L, Mbwana H, Kinabo J, Biesalski HK, Riedel S, et al. Validation of a food group based nutrition software to assess nutrient intake in Tanzania. In: Global food security and food safety: The role of universities. Ghent, Belgium: Ghent University; 2018. p. 2018.

## 6. Summary

Religious fasting is one of the categories of dietary or food taboos, which may affect the dietary intake and nutritional status of individuals. In Ethiopia, about half of the population are followers of Ethiopian Orthodox Tewahedo religion, and approximately 250 days per annum are fasting days. In these fasting days, lactating and pregnant women and children are exempted from fasting. However, lactating and pregnant women fast and are not also happy to prepare non-fasting foods for their children during the fasting days due to fear of contamination of family food.

Early identification of micronutrient deficiencies in Ethiopia are flouted, as most often the quantitative dietary data are not available. As a result, the hidden hunger might have not been addressed properly, where it remains high and persistent. Therefore, easy to use, less costly and applicable assessment tool which can estimate the quantitative dietary intake of an individual or a community is urgently needed to achieve the national and international goals set for eradicating malnutrition. The Calculator for Inadequate Micronutrient Intake (CIM) is a simple, easy-to-use, informative, web-based application of quantitative dietary assessment method, which was first developed in Indonesia for Indonesian population. It estimates energy and nutrient intake correctly, and identifies nutrient inadequacy according to FAO/WHO recommended nutrient intake (RNI) regarding age, sex and physiological stage. Thus, the present study was conducted with the aim of assessing and comparing the nutritional status and dietary intake of lactating women and their 6-23-months-old children in fasting and non-fasting periods, and to adapt and validate the CIMI program for Ethiopian population.

This study was conducted in rural Genta Afeshum district, in Tigray, Ethiopia, where almost all people in the woreda are followers of Ethiopian Orthodox Christianity. A longitudinal study was conducted using 575 and 522 lactating women and their 6-23-months-old children in the lent fasting and non-fasting.

In the present study it was found out that the prevalence of underweight ( $\text{BMI} < 18.5 \text{ kg/m}^2$ ) in fasting lactating women was high (50.6%) which is associated with maternal age, maternal illness within four weeks preceding the fasting survey, fasting status during their pregnancy and lactation period of their children included in this study. Additional predictor variables for maternal underweight were grandfathers' as household decision maker, use of non-improved water source, household aid experience and the absence of chicken in the household. The average number of meals, diet diversity, and animal source foods consumption scores were significantly higher in non-fasting compared to fasting periods, regardless of the fasting status ( $p < 0.001$ ,  $p < 0.05$  and  $p < 0.001$ , respectively). Whereas, 31.6–33.7%, 11.7–15.7% and 4.4–4.8% of the 6-23-months-old children in the study population were stunted, underweight and wasted, respectively. In the fasting period, the weight-for-length (WLZ) and length-for-age (LAZ) values for the 6-23-months-old children of non-fasting mothers were significantly higher ( $p < 0.05$ ) than the children of the fasting mothers' group. Similarly, the median weight-for-age (WAZ) and diet diversity score (DDS) of children of fasting mothers were also significantly lower in fasting compared to non-fasting period. The proportion of the 6-23-months-old children who met the minimum acceptable diet (MAD) was small (2.3-6.7%) in the study population; however, this proportion was significantly ( $p < 0.001$ ) higher in the non-fasting than fasting period in the children of fasting mothers. Age of the child, maternal fasting status during pregnancy and

lactation periods, maternal education and occupation were associated with child underweight. Likewise, age of the child, colostrum intake status, maternal fasting status during pregnancy and lactation period and toilet presence in the household were associated with child stunting. But, maternal fasting status during lactation period and maternal education predicted wasting in the children precisely. The average energy, protein and almost all micronutrients intakes of children and women were lower in fasting compared to non-fasting period. At the same time, the prevalence of inadequate intakes of energy, protein and most micronutrients were higher in both the children and lactating women during fasting than non-fasting period. The result of this study revealed that the correlation coefficients for the average dietary nutrient intake calculated by CIMI and the reference software NutriSurvey (NS) were between 0.741-0.956 for the children and between 0.779-0.920 for the lactating women groups. As a conclusion, the dietary pattern and nutritional status of lactating women and their breastfed children are affected during the fasting period. Therefore, the existing multi-sectoral nutrition intervention strategies in Ethiopia should include religious institutions in a sustainable manner. CIMI adapted for the rural Ethiopian setting estimates the average nutrient intake accurately; and identifies inadequate micronutrient intake of individuals enabling enumerators to provide feedback and suggest improvements. Thus, CIMI can be used in Ethiopia, as a simple dietary assessment tool by nutrition and related researchers, policy makers, implementers and evaluators.



## **Zusammenfassung**

Beim religiösen Fasten werden Lebensmitteltabus praktiziert, die die Nährstoffaufnahme und den Ernährungszustand eines Individuums beeinflussen können. In Äthiopien besteht etwa die Hälfte der Bevölkerung aus äthiopisch-orthodoxen Anhängern der Tewahedo-Religion. In dieser Religion wird an ungefähr 250 Tagen im Jahr gefastet, wobei stillende und schwangere Frauen und Kinder vom Fasten ausgeschlossen sind. In der Regel wird das Fasten jedoch auch von laktierenden und schwangeren Frauen praktiziert, und diese Frauen bereiten ihren Kindern während der Fastentage nicht gerne tierische Lebensmittel zu, aus Angst, das Familienessen zu „verunreinigen“.

Die frühzeitige Erkennung von Mikronährstoffmängeln in Äthiopien scheitert meist an der mangelnden Verfügbarkeit von quantitativen Daten zur Nährstoffaufnahme. Infolgedessen wird der verborgene Hunger nicht richtig wahrgenommen und bekämpft, wodurch er dauerhaft und in einer hohen Prävalenz auftritt. Um die nationalen und internationalen Ziele zur Beseitigung der Unterernährung zu erreichen, ist ein einfaches, kostengünstiges und praktisch anzuwendendes Erhebungsinstrument mit dem die quantitative Nährstoffaufnahme einer Einzelperson oder einer Gemeinschaft geschätzt werden kann, dringend erforderlich. Der Calculator for Inadequate Micronutrient Intake (CIMI) ist eine simple, benutzerfreundliche, informative App zur quantitativen Erhebung der Micro- und Makronährstoffzufuhr, deren Vorläuferversion für die indonesische Bevölkerung entwickelt wurde. CIMI schätzt die Energie- und Nährstoffaufnahme korrekt ein und identifiziert gemäß der FAO / WHO empfohlenen Nährstoffzufuhr (RNI) eine unangemessen niedrige Nährstoffaufnahme in Bezug auf Alter, Geschlecht und Schwangerschaft/Stillzeit.

Ziel der vorliegenden Studie war die Erfassung und Beurteilung des Ernährungsstatus und der Nahrungsaufnahme laktierender Frauen und ihrer 6-23 Monate alten Kinder in Fasten- und Nicht-Fastenzeiten. Desweiteren sollte das CIMI-Programm an die äthiopische Bevölkerung angepasst und dessen Anwendbarkeit untersucht werden.

Diese Studie wurde im ländlichen Bezirk Genta Afeshum in Tigray, Äthiopien durchgeführt, in dem fast alle Menschen Anhänger des äthiopisch-orthodoxen Christentums sind. Es handelte sich um eine longitudinale Studie an der 575 und 522 stillende Frauen und ihre 6-23 Monate alten Kinder während einer Fasten- und einer Nicht-Fastenperiode teilnahmen.

Die vorliegende Studie ergab, dass die Prävalenz von Untergewicht ( $\text{BMI} < 18,5 \text{ kg} / \text{m}^2$ ) bei fastenden laktierenden Frauen hoch war (50,6%). Diese hohe Prävalenz an Untergewicht war mit dem mütterlichen Alter, einer Erkrankung der Mutter innerhalb von vier Wochen vor der Befragung, des Fastens während der Schwangerschaft und der Stillzeit ihrer in diese Studie eingeschlossenen Kinder assoziiert. Weitere Prädiktoren für ein mütterliches Untergewicht waren Großväter als Entscheidungsträger im Haushalt, die Verwendung einer unsicheren Wasserquelle, der Erhalt von Haushaltshilfen und das Fehlen von Hühnern im Haushalt. Die durchschnittliche Mahlzeitenanzahl, die Nahrungsvielfalt und der Konsum von tierischen Lebensmitteln war in der Nicht-Fastenperiode signifikant höher ( $p < 0,001$ ,  $p < 0,05$  bzw.  $p < 0,001$ ), unabhängig vom Fastenstatus. Dem gegenüber waren 31,6–33,7%, 11,7–15,7% und 4,4–4,8% der 6-23 Monate alten Kinder in der Studienpopulation zu klein für ihr Alter, untergewichtig und zu leicht für ihr Alter. In der Fastenperiode waren die Werte für das Gewicht für die Körpergröße (WLZ) und für die Größe des Alters (LAZ) für die 6-23 Monate alten Kinder nicht fastenden Müttern signifikant höher ( $p < 0,05$ ) als die Kinder der Gruppe der

fastenden Müttern. In ähnlicher Weise waren auch der Median des Altersgewichts (WAZ) und des Diet Diversity Scores (DDS) der Kinder von fastenden Müttern während der Fastenperiode signifikant niedriger als im fastenfreien Zeitraum. Der Anteil der 6-23 Monate alten Kinder, die die akzeptable Mindestdiät (MAD) verzehrten, war in der Studienpopulation gering (2,3-6,7%). Bei Kindern von fastenden Müttern war dieser Anteil jedoch in der Nicht-Fastenphase ( $n < p < 0,001$ ) signifikant höher als in der Fastenzeit. Das Alter des Kindes, der Fastenstatus der Mutter während der Schwangerschaft und Stillzeit, die Bildung und Beschäftigung der Mutter waren mit kindlichem Untergewicht assoziiert. Ebenso waren das Alter des Kindes, die Kolostrumgabe, der Fastenstatus der Mutter während der Schwangerschaft und Stillzeit und die Verfügbarkeit einer Toilette im Haushalt mit Stunting verbunden. Der Fastenstatus der Mutter während der Stillzeit und die Bildung der Mutter sagten das Risiko für ein zu geringes altersspezifisches Körpergewicht des Kindes präzise voraus. Die durchschnittliche Zufuhr an Energie-, Eiweiß- und fast aller Mikronährstoffe von Kindern und Frauen waren im Vergleich zu Nicht-Fastenzeiten niedriger. Gleichzeitig war die Prävalenz einer unzureichenden Zufuhr von Energie, Eiweiß und den meisten Mikronährstoffen sowohl bei Kindern als auch bei stillenden Frauen während des Fastens höher als bei Nicht-Fasten. Das Ergebnis dieser Studie zeigt, dass die von CIMI und der Referenzsoftware Nutrisurvey (NS) berechneten Korrelationskoeffizienten für die durchschnittliche Nährstoffaufnahme zwischen 0,741-0,956 bei Kinder und 0,779-0,920 bei laktierenden Frauen lagen.

Zusammenfassen lässt sich sagen, dass durch das Fasten das Ernährungsmuster und der Ernährungszustand von stillenden Frauen und deren Kindern während der Fastenzeit betroffen ist. Daher sollten in Äthiopien religiöse Institutionen in multisektorale Ernährungsinterventionen

einbezogen werden. CIMI, angepasst an die ländliche äthiopische Ernährungsweise, schätzt die durchschnittliche Nährstoffaufnahme präzise ein und identifiziert eine unzureichende Mikronährstoffaufnahme von Individuen, die es Enumeratoren ermöglicht Feedback zu geben und Verbesserungen vorzuschlagen. Daher kann CIMI in Äthiopien als einfaches Instrument zur Beurteilung der Ernährung von Wissenschaftlern, politischen Entscheidungsträgern und Mitarbeitern von Ernährungserhebungen und -interventionen verwendet werden.

## List of all references

- Abate, K. H., & Belachew, T. (2017). Women's autonomy and men's involvement in child care and feeding as predictors of infant and young child anthropometric indices in coffee farming households of Jimma Zone, South West of Ethiopia. *PLoS ONE*, 12(3), 1–16. <https://doi.org/10.1371/journal.pone.0172885>
- Abebe, Y., Bogale, A., Hambidge, K., Stoecker, J., Bailey, K., & Gibson, R. (2007). Phytate , zinc , iron and calcium content of selected raw and prepared foods consumed in rural Sidama , Southern Ethi ... foods consumed in rural Sidama , Southern Ethiopia , and impl. *Journal of Food Composition and Analysis*, 20(July), 161–168. <https://doi.org/10.1016/j.jfca.2006.09.003>
- Abrha, S., Shiferaw, S., & Ahmed, K. Y. (2016). Overweight and obesity and its socio-demographic correlates among urban Ethiopian women: Evidence from the 2011 EDHS. *BMC Public Health*, 16(1), 1–7. <https://doi.org/10.1186/s12889-016-3315-3>
- Addo, O. Y., Stein, A. D., Fall, C. H., Gigante, D. P., Guntupalli, A. M., Horta, B. L., ... Martorell, R. (2013). Maternal height and child growth patterns. *Journal of Pediatrics*, 163(2), 549–554.e1. <https://doi.org/10.1016/j.jpeds.2013.02.002>
- Ahmed, A., Abdulahi, A., Shab-bidar, S., and Rezaei, S. (2017). Nutritional Status of Under Five Children in Ethiopia : A Systematic Review and Meta-Analysis. *Ethiopian Journal of Health Science*, 27(2), 175-188
- Akombi, B. J., Agho, K. E., Hall, J. J., Wali, N., Renzaho, A. M. N., and Merom, D. (2017). Stunting, wasting and underweight in Sub-Saharan Africa: A systematic review.

- International Journal of Environmental Research and Public Health*, 14(8), 1–18.  
<https://doi.org/10.3390/ijerph14080863>
- Alemayehu, M., Tinsae, F., Hailelassie, K., Seid, O., Gebregziabher, G., & Yebyo, H. (2015). Undernutrition status and associated factors in under-5 children, in Tigray, Northern Ethiopia. *Nutrition*, 31(7–8), 964–970. <https://doi.org/10.1016/j.nut.2015.01.013>
- Alemayehu, M., Argaw, A., & Geberemariam, A. (2015). Factors Associated with Malnutrition among Lactating Women in Subsistence Farming Households from Dedo and Seqa-Chekorsa Districts, Jimma Zone, 2014. *Developing Country Studies*, 5(21), 114–122.  
 Retrieved from <http://www.iiste.org/Journals/index.php/DCS/article/view/27469>
- Alemayehu, M., Tinsae, F., Hailelassie, K., Seid, O., Gebregziabher, G., & Yebyo, H. (2015). Undernutrition status and associated factors in under-5 children, in Tigray, Northern Ethiopia. *Nutrition*, 31(7–8), 964–970. <https://doi.org/10.1016/j.nut.2015.01.013>
- Alemu, Z. A., Ahmed, A. A., Yalew, A. W., and Birhanu, B. S. (2016). Non random distribution of child undernutrition in Ethiopia: Spatial analysis from the 2011 Ethiopia demographic and health survey. *International Journal for Equity in Health*.  
<https://doi.org/10.1186/s12939-016-0480-z>.
- Ali, D., Saha, K. K., Nguyen, P. H., Diressie, M. T., Ruel, M. T., Menon, P., & Rawat, R. (2013). Household Food Insecurity Is Associated with Higher Child Undernutrition in Bangladesh, Ethiopia, and Vietnam, but the Effect Is Not Mediated by Child Dietary Diversity. *Journal of Nutrition*, 143(12), 2015–2021. <https://doi.org/10.3945/jn.113.175182>

- Ali Naser, I., Jalil, R., Wan Muda, W. M., Wan Nik, W. S., Mohd Shariff, Z., & Abdullah, M. R. (2014). Association between household food insecurity and nutritional outcomes among children in Northeastern of peninsular Malaysia. *Nutrition Research and Practice*, 8(3), 304–311. <https://doi.org/10.4162/nrp.2014.8.3.304>
- Ali, Z., and Abizari, A. (2018). Ramadan fasting alters food patterns , dietary diversity and body weight among Ghanaian adolescents. *Nutrition Journal*, 17(75), 1–14.
- Alive & Thrive. (2010). IYCF Practices , Beliefs , and Influences in Tigray Region , Ethiopia. Alive & Thrive.
- Altare, C., Delbiso, T. D., Mutwiri, G. M., Kopplow, R., & Guha-Sapir, D. (2016). Factors associated with stunting among pre-school children in southern highlands of Tanzania. *Journal of Tropical Pediatrics*, 62(5), 390–408. <https://doi.org/10.1093/tropej/fmw024>
- Altuntas, Y. E., Gezen, F. C., Sahoniz, T., Kement, M., Aydin, H., Sahin, F., ... Oncel, M. (2013). Ramadan fasting in patients with a stoma: a prospective study of quality of life and nutritional status. *Ostomy Wound Management*, 59(5), 26–32. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/23669258>
- Amare, B., Moges, B., Moges, F., Fantahun, B., Admassu, M., Mulu, A., & Kassu, A. (2012). Nutritional status and dietary intake of urban residents in Gondar, Northwest Ethiopia. *BMC Public Health*, 12(1). <https://doi.org/10.1186/1471-2458-12-752>
- Amagloh, F., Atuna, R., McBride, R., Carey, E., & Christides, T. (2017). Nutrient and Total Polyphenol Contents of Dark Green Leafy Vegetables, and Estimation of Their Iron Bioaccessibility Using the In Vitro Digestion/Caco-2 Cell Model. *Foods*, 6(7), 54.

<https://doi.org/10.3390/foods6070054>

- Amugsi, D. A., Mittelmark, M. B., & Oduro, A. (2015). Association between Maternal and Child Dietary Diversity : An Analysis of the Ghana Demographic and Health Survey. *PLoS ONE*, *10*(8), e0136748. <https://doi.org/10.1371/journal.pone.0136748>
- Anteneh, Z. A., Gedefaw, M., Tekletsadek, K. N., Tsegaye, M., & Alemu, D. (2015). Risk Factors of Overweight and Obesity among High School Students in Bahir Dar City, North West Ethiopia: School Based Cross-Sectional Study. *Advances in Preventive Medicine*, *2015*, 1–9. <https://doi.org/10.1155/2015/294902>
- Arimond, M., & Ruel, M. T. (2004). Dietary diversity is associated with child nutritional status : Evidence from 11 demographic and health surveys. *J.Nutr.*, *134*, 2579–2585.
- Alemu, E., & Atnafu, A. (2014). Prevalence of Overweight and/or Obesity and Associated Factors among High School Adolescents in Arada Sub city, Addis Ababa, Ethiopia. *Journal of Nutrition & Food Sciences*, *4*(2), 10–14. <https://doi.org/10.4172/2155-9600.1000261>
- Almond, D., & Mazumder, B. Federal Reserve Bank of Chicago Health Capital and the Prenatal Fasting During Pregnancy Health Capital and the Prenatal Environment : The Effect of Maternal Fasting During Pregnancy \*, No. 2007-2 FRB of Chicago Working Paper 74 (2011).
- Altuntas, Y. E., Gezen, F. C., Sahoniz, T., Kement, M., Aydin, H., Sahin, F., ... Oncel, M. (2013). Ramadan fasting in patients with a stoma: a prospective study of quality of life and nutritional status. *Ostomy Wound Management*, *59*(5), 26–32. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/23669258>



- Arimond, M., and Ruel, M. T. (2004). Dietary diversity is associated with child nutritional status : Evidence from 11 demographic and health surveys. *J.Nutr.*, *134*, 2579–2585.
- Arzoaquoi, S. K., Essuman, E. E., Gbagbo, F. Y., Tenkorang, E. Y., Soyiri, I., and Laar, A. K. (2015). Motivations for food prohibitions during pregnancy and their enforcement mechanisms in a rural Ghanaian district. *Journal of Ethnobiology and Ethnomedicine*, *11*(1). <https://doi.org/10.1186/s13002-015-0044-0>
- Asayehu, T. T., Lachat, C., Henauw, S. D., & Gebreyesus, S. H. (2017). Dietary behaviour, food and nutrient intake of women do not change during pregnancy in Southern Ethiopia. *Maternal and Child Nutrition*, *13*(2). <https://doi.org/10.1111/mcn.12343>
- Avery, A. (2004). Red Meat and Poultry Production and Consumption in Ethiopia and Distribution in Addis Ababa, Ethiopia (August).
- Awoke, A., Awoke, T., Alemu, S., & Megabiaw, B. (2012). Prevalence and associated factors of hypertension among adults in Gondar, Northwest Ethiopia: A community based cross-sectional study. *BMC Cardiovascular Disorders*, *12*(1), 1. <https://doi.org/10.1186/1471-2261-12-113>
- Awwad, J., Usta, I. M., Succar, J., Musallam, K. M., Ghazeeri, G., and Nassar, A. H. (2012). The effect of maternal fasting during Ramadan on preterm delivery: A prospective cohort study. *BJOG: An International Journal of Obstetrics and Gynaecology*, *119*(11), 1379–1386. <https://doi.org/10.1111/j.1471-0528.2012.03438.x>
- Ayana, G., Moges, T., Samuel, A., Asefa, T., Eshetu, S., & Kebede, A. (2018). Dietary zinc

- intake and its determinants among Ethiopian children 6–35 months of age. *BMC Nutrition*, 4(1), 30. <https://doi.org/10.1186/s40795-018-0237-8>
- Ayana, G., Moges, T., Samuel, A., Asefa, T., Eshetu, S., & Kebede, A. (2018). Dietary zinc intake and its determinants among Ethiopian children 6–35 months of age. *BMC Nutrition*, 4(1), 30. <https://doi.org/10.1186/s40795-018-0237-8>
- Baye, K., Guyot, J. P., Icard-Vernière, C., & Mouquet-Rivier, C. (2013). Nutrient intakes from complementary foods consumed by young children (aged 12-23 months) from North Wollo, northern Ethiopia: the need for agro-ecologically adapted interventions. *Public Health Nutrition*, 16(10), 1741–1750. <https://doi.org/10.1017/S1368980012005277>.
- Bbaale, E. (2014). Maternal education and child nutritional status : evidence from Uganda. *African Journal of Economic and Management Studies*, 5(1), 52–74. <https://doi.org/10.1108/AJEMS-06-2012-0040>.
- Belew, A. K., Ali, B. M., Abebe, Z., & Dachew, B. A. (2017). Dietary diversity and meal frequency among infant and young children: A community based study. *Italian Journal of Pediatrics*, 43(1), 6–15. <https://doi.org/10.1186/s13052-017-0384-6>.
- Betebo, B., Ejajo, T., Alemseged, F., & Massa, D. (2017). Household Food Insecurity and Its Association with Nutritional Status of Children 6 – 59 Months of Age in East Badawacho District, South Ethiopia. *Journal of Environmental and PublicHealth*, 2017. <https://doi.org/10.1155/2017/6373595>.
- Bhandari, S., and Banjara, M. R. (2015). Micronutrients Deficiency, a Hidden Hunger in Nepal: Prevalence, Causes, Consequences, and Solutions. *International Scholarly Research*

*Notices*, 2015, 1–9. <https://doi.org/10.1155/2015/276469>.

Bhandari, S., Sayami, J. T., Thapa, P., Sayami, M., Kandel, B. P., & Banjara, M. R. (2016).

Dietary intake patterns and nutritional status of women of reproductive age in Nepal:

Findings from a health survey. *Archives of Public Health*, 74(1), 1–11.

<https://doi.org/10.1186/s13690-016-0114-3>.

Biesalski Hans, K., and Jana, T. (2018). Micronutrients in the life cycle: Requirements and

sufficient supply. *NFS Journal*, 11(March), 1–11. <https://doi.org/10.1016/j.nfs.2018.03.001>.

Bililign, N., Legesse, M., & Akibu, M. (2018). A Review of Low Birth Weight in Ethiopia :

Socio-Demographic and Obstetric Risk Factors. *Global Journal of Research and Review*.

5(1:4). <https://doi.org/10.21767/2393-8854.100033>.

Birhanu, A., Mekonen, S., & Atenafu, A. (2017). Stunting and associated factors among children

aged 6-59 months in Lasta woreda , north east Ethiopia , 2015 : A community based cross

sectional study design. *Journal of Family Medicine*, 4(3), 1112.

Belew, A. K., Ali, B. M., Abebe, Z., & Dachew, B. A. (2017). Dietary diversity and meal

frequency among infant and young children: A community based study. *Italian Journal of*

*Pediatrics*, 43(1), 6–15. <https://doi.org/10.1186/s13052-017-0384-6>.

Biza Zepro, N. (2015). Food Taboos and Misconceptions Among Pregnant Women of

Shashemene District, Ethiopia, 2012. *Science Journal of Public Health*, 3(3), 410.

<https://doi.org/10.11648/j.sjph.20150303.27>.

Black, R. E., Allen, L. H., Bhutta, Z. A., Caulfield, L. E., de Onis, M., Ezzati, M., ... Rivera, J.

- (2008). Maternal and child undernutrition: global and regional exposures and health consequences. *The Lancet*, 371(9608), 243–260. [https://doi.org/10.1016/S0140-6736\(07\)61690-0](https://doi.org/10.1016/S0140-6736(07)61690-0).
- Biru, K., Jima, A., & Abeya, S. (2017). Prevalence of Chronic Energy Malnutrition and Maternal Health Service Utilizations among Lactating Mothers in Adama District, Oromia Region, Eastern Ethiopia. *Journal of Food Processing & Technology*, 8(1), 1–5.  
<https://doi.org/10.4172/2157-7110.1000650>
- Bitew, F. H., & Telake, D. S. (2010). *Undernutrition among Women in Ethiopia. B Undernutrition among Women in Ethiopia: Rural-Urban Disparity. DHS Working Papers*, (77).
- Caulfield, L. E., Onis, M. De, Blössner, M., & Black, R. E. (2004). Undernutrition as an underlying cause of child deaths associated. *Am J Clin Nutr*, 80, 193–198.
- Cogill, B. (2003). Anthropometric Indicators Measurement Guide Anthropometric Indicators Measurement Guide. *Food and Nutritional Technical Assistance, 2003 Revis*, 1–93.  
Retrieved from <http://www.fantaproject.org/sites/default/files/resources/anthropometry-2003-ENG.pdf>.
- CSA Central Statistical Agency of Ethiopia. (2007). Population and Housing Census 2007 Report. Central Statistical Agency, Addis Ababa, 385. Retrieved from [http://www.csa.gov.et/surveys/Population and Housing census/ETH-pop-2007/survey0/data/Doc/Reports/National\\_Statistical.pdf](http://www.csa.gov.et/surveys/Population%20and%20Housing%20census/ETH-pop-2007/survey0/data/Doc/Reports/National_Statistical.pdf)

CSA. (2012). *Ethiopia Demographic and Health Survey 2011*. Addis Ababa.

CSA. (2017). *Ethiopia Demographic and Health Survey 2016*. Addis Ababa.

Cunningham, K., Headey, D., Singh, A., Karmacharya, C., & Rana, P. (2017). Maternal and Child Nutrition in Nepal : Examining drivers of progress from the mid-1990s to 2010s.

*Global Food Security*, 13(February), 30–37. <https://doi.org/10.1016/j.gfs.2017.02.001>

Custodio, E., Perez-Formigo, J., Gadisa, E., Sordo, L., Cruz, I., Canavate, C., ... Moreno, J. (2013). Low dietary diversity is associated with child stunting in rural Ethiopia. *Annals of Nutrition and Metabolism*. Conference paper.

Dangura, D., & Gebremedhin, S. (2017). Dietary diversity and associated factors among children 6-23 months of age in Gorche district, Southern Ethiopia: Cross-sectional study. *BMC Pediatrics*, 17(1), 1–7. <https://doi.org/10.1186/s12887-016-0764-x>.

Demissie, T., Muroki, N. M., and Kogi-Makau, W. (1998). Food aversions and cravings during pregnancy: prevalence and significance for maternal nutrition in Ethiopia. *Food and Nutrition Bulletin*, 19(1), 20–26. <https://doi.org/10.1177/156482659801900104>.

Demissie, T., Mekonen, Y., and Haider, J. (2003). Agroecological comparison of leveles and correlates of nutritional status of women. *Ethiopian Journal of Health Development*, 17(3), 189–196.

Demissie, T., Ali, A., & Zerfu, D. (2009). Availability and consumption of fruits and vegetables in nine regions of Ethiopia with special emphasis to vitamin A deficiency. *Ethiopian Journal of Health Development*, 23(3), 216–222.

- Demissie, T., Ali, A., Mekonen, Y., Haider, J., & Umeta, M. (2010). Magnitude and distribution of vitamin A deficiency in Ethiopia. *Food and Nutrition Bulletin*, 31(2), 234–241. <https://doi.org/10.1177/156482651003100206>
- Derso, T., Tariku, A., Biks, G. A., & Wassie, M. M. (2017). Stunting, wasting and associated factors among children aged 6-24 months in Dabat health and demographic surveillance system site: A community based cross-sectional study in Ethiopia. *BMC Pediatrics*, 17(1), 1–9. <https://doi.org/10.1186/s12887-017-0848-2>
- Desalegn, B. B., Kinf, E., Fikre, K., & Bosha, T. (2016). Stunting and Its Associated Factors in Under Five Years Old Children : The Case of Hawassa University Technology Villages , Southern Ethiopia. *IOSR Journal of Environmental Science, Toxicology and Food Technology*, 10(11), 25–31. <https://doi.org/10.9790/2402-1011022531>
- Divya, N., Rajanish, K. V, Malavika, J., & Sharma, A. (2018). The study of dietary diversity score in children between 6 months to 23 months : a hospital based study. *International Journal of Contemporary Pediatrics*, 5(3), 1053–1057. DOI: <http://dx.doi.org/10.18203/2349-3291.ijcp20181541>
- Desalegn, B. B., Lambert, C., Riedel, S., Negese, T., & Biesalski, H. K. (2018). Ethiopian orthodox fasting and lactating mothers: Longitudinal study on dietary pattern and nutritional status in rural tigray, Ethiopia. *International Journal of Environmental Research and Public Health*, 15(8). <https://doi.org/10.3390/ijerph15081767>
- Desalegn, B., Lambert, C., Riedel, S., Negese, T., & Biesalski, H. (2019). Feeding Practices and Undernutrition in 6–23-Month-Old Children of Orthodox Christian Mothers in Rural Tigray, Ethiopia: Longitudinal Study. *Nutrients*, 11(1), 138.

<https://doi.org/10.3390/nu11010138>

Development Initiatives. (2017). *Global Nutrition Report 2017: Nourishing the SDGs*. Bristol, UK: Development Initiative, 115. <https://doi.org/10.1891/1058-1243.21.1.9>

Development initiatives. (2018). *2018 Global nutrition report: Shining a light to spur action on nutrition*. Bristol, UK. <https://doi.org/10.2499/9780896295643>

Dewey, K. G., & Brown, K. H. (2003). Update on technical issues concerning complementary feeding of young children in developing countries and implications for intervention programs. *Food and Nutrition Bulletin*, 24(1), 5–28.

DSM. (2015). Nutrition during the first 1 , 000 days. In *Vitamins in Motion for a brighter world* (pp. 1–24).

Ekwochi, U., Osuorah, C. D. I., Ndu, I. K., Ifediora, C., Asinobi, I. N., and Eke, C. B. (2016). Food taboos and myths in South Eastern Nigeria: The belief and practice of mothers in the region. *Journal of Ethnobiology and Ethnomedicine*, 12(1). <https://doi.org/10.1186/s13002-016-0079-x>

Endalamaw, A., Engeda, E. H., Ekubagewargies, D. T., & Belay, G. M. (2018). Low birth weight and its associated factors in Ethiopia : a systematic review and meta- analysis. *Italian Journal of Pediatrics*, 44 (141). <https://dx.doi.org/10.1186%2Fs13052-018-0586-6>

Endalew, B., Muche, M., and Tadesse, S. (2015). Assesment of food security situation in Ethiopia: A review. *Asian Journal of Agricultural Research*, 9(2), 55–68. <https://doi.org/10.3923/ajar.2015.55.68>

- Endris, N., Asefa, H., & Dube, L. (2017). Prevalence of Malnutrition and Associated Factors among Children in Rural Ethiopia. *BioMed Research International*, 2017, 8–10. <https://doi.org/10.1155/2017/658785>
- EOTC/USAID. (2016, January 12). Patriarch of the Ethiopian Orthodox Tewahedo Church Endorses Nutrition Sermon Guide Encouraging Better Nutrition for Mothers and Children ., pp. 1–2.
- Ersino, G., Henry, C. J., and Zello, G. A. (2016). Suboptimal Feeding Practices and High Levels of Undernutrition among Infants and Young Children in the Rural Communities of Halaba and Zeway, Ethiopia. *Food and Nutrition Bulletin*, 37(3), 409–424. <https://doi.org/10.1177/0379572116658371>
- Essential Nutrition Action (ENA). (2008). *Key behaviours for optimal breastfeeding, complementary feeding and maternal nutrition at critical stages in the life cycle of women and children*.
- Ethiopian Public Health Institute (EPHI). (2013). *Ethiopia National Food Consumption Survey*. Addis Ababa, Ethiopia.
- Ethiopian National Micronutrient Survey Report. (2016). Addis Ababa, Ethiopia.
- Ettyang, G. A. K., & Sawe, C. J. (2016). Factors associated with stunting in children under age 2 in the Cambodia and Kenya 2014 Demographic and Health Surveys. *DHS Working Papers No. 126* , (August). Retrieved from <http://dhsprogram.com/pubs/pdf/WP126/WP126.pdf>



FAO. (2016). FAO in Ethiopia *El Nino response plan 2016*. Addis Ababa, Ethiopia.

FAO. (2017). *Ethiopia Situation Report - May 2017*. Addis Ababa, Ethiopia.

FAO, & FHI360. (2016). Minimum Dietary Diversity for Women- A Guide to Measurement. Minimum Dietary Diversity for Women: A Guide for Measurement (p. 82).

FAO, IFAD, UNICEF, WFP, & WHO. (2018). The State of Food Security and Nutrition in the World 2018. Building climate resilience for food security and nutrition. Rome, FAO (pp. 3157–3166). <https://doi.org/10.1109/JSTARS.2014.2300145>

FAO and, & WHO. (2001). Human Vitamin and Mineral Requirements. Food and Nutrition Division, 181–94.

FAO, WHO, & UNU. (2001). *Human Energy Requirements. FAO Food and nutrition technical report* (pp. 1–105).

Protein and amino acid requirements in human nutrition. (2007). *World Health Organization Technical Report Series*.

Faris, M. A.-I. E., and Al-Holy, M. A. (2014). Implications of Ramadan intermittent fasting on maternal and fetal health and nutritional status: A review. *Mediterranean Journal of Nutrition and Metabolism*, 7, 107–118. <https://doi.org/10.3233/MNM-140011>

Fayet-Moore, F., Kim, J., Sritharan, N., & Petocz, P. (2016). Impact of breakfast skipping and breakfast choice on the nutrient intake and body mass index of Australian children. *Nutrients*, 8(8). <https://doi.org/10.3390/nu8080487>

Fasts, F. W., and Fasts, T. P. (2019). Fasting and abstinence of the Coptic Orthodox Church of

Alexandria.

[https://en.wikipedia.org/wiki/Fasting\\_and\\_abstinence\\_of\\_the\\_Coptic\\_Orthodox\\_Church\\_of\\_Alexandria](https://en.wikipedia.org/wiki/Fasting_and_abstinence_of_the_Coptic_Orthodox_Church_of_Alexandria). Accessed on March 19, 2019

Federal Republic of Ethiopia, D. R. M. and F. S. (2015). *Disaster Risk Management and Food Security Sector: Wereda Disaster Risk Profile: Genta Ahfeshum, Eastern Tigray. February 2015 Addis Ababa.*

Federal Democratic Republic of Ethiopia (FDRE). (2016). *SEQOTA declaration implementation plan ( 2016 – 2030): Summary programme approach document*. Addis Ababa, Ethiopia.

Felisbino-mendes, M., Villamor, E., & Velasquez-melendez, G. (2014). Association of maternal and child nutritional status in Brazil : A population based cross-sectional study. *PLoS ONE*, 9(1), e87486. <https://doi.org/10.1371/journal.pone.0087486>

Fentahun, W., Wubshet, M., & Tariku, A. (2016). Undernutrition and associated factors among children aged 6-59 months in East Belesa District, northwest Ethiopia: A community based cross-sectional study. *BMC Public Health*, 16(1), 1–10. <https://doi.org/10.1186/s12889-016-3180-0>

Fentahun, N., Belachew, T., & Lachat, C. (2016). Determinants and morbidities of multiple anthropometric deficits in southwest rural Ethiopia. *Nutrition*, 32(11–12), 1243–1249. <https://doi.org/10.1016/j.nut.2016.03.023>

FMOH. (2008). *The Federal Democratic Republic of Ethiopia: National Nutrition Strategy*. Addis Ababa, Ethiopia.

FMOH. (2013). *National Nutrition Program Republic of Ethiopia*. Addis Ababa, Ethiopia.

FMOH. (2016). *National Nutrition Program 2016-2020*. Addis Ababa, Ethiopia.

Gali, N., Tamiru, D., & Tamrat, M. (2017). The Emerging Nutritional Problems of School Adolescents : Overweight / Obesity and Associated Factors in Jimma Town , Ethiopia. *Journal of Pediatric Nursing*, 35, 98–104. <https://doi.org/10.1016/j.pedn.2017.03.002>

Gamo Sagaro, G. (2017). Dietary Diversity and Associated Factors Among Infants and Young Children in Wolaita Zone, Southern Ethiopia. *Science Journal of Clinical Medicine*, 6(4), 53. <https://doi.org/10.11648/j.sjcm.20170604.12>

Gebre, B., Biadgilign, S., Taddese, Z., Legesse, T., and Letebo, M. (2018). Determinants of malnutrition among pregnant and lactating women under humanitarian setting in Ethiopia. *BMC Nutrition*, 4(11), 1–8. <https://doi.org/https://doi.org/10.1186/s40795-018-0222-2>

Gebreegziabher, T., & Stoecker, B. J. (2013). Vitamin D insufficiency in a sunshine-sufficient area: Southern ethiopia. *Food and Nutrition Bulletin*, 34(4), 429–433. <https://doi.org/10.1177/156482651303400408>

Gebremedhin, S., Enquselassie, F., & Umata, M. (2011). Prevalence of prenatal zinc deficiency and its association with socio-demographic, dietary and health care related factors in Rural Sidama, Southern Ethiopia: A cross-sectional study. *BMC Public Health*, 11. <https://doi.org/10.1186/1471-2458-11-898>

Gebremedhin, M., Ambaw, F., Admassu, E., & Berhane, H. (2015). Maternal associated factors of low birth weight : a hospital based cross-sectional mixed study in Tigray , Northern

- Ethiopia. *BMC Pregnancy and Childbirth*, 15(22), 1–8. <https://doi.org/10.1186/s12884-015-0658-1>
- Gebremedhin, S., Enquselassie, F., & Umeta, M. (2014). Prevalence and Correlates of Maternal Anemia in Rural Sidama, . Southern Ethiopia. *African Journal of Reproductive Health*, 18(1), 44–53.
- Gebremedhin, S., Baye, K., Bekele, T., Tharaney, M., Asrat, Y., Abebe, Y., & Reta, N. (2017). Predictors of dietary diversity in children ages 6 to 23 mo in largely food-insecure area of South Wollo, Ethiopia. *Nutrition*, 33, 163–168. <https://doi.org/10.1016/j.nut.2016.06.002>
- Gebreselassie, S. G., Gase, F. E., & Deressa, M. U. (2013). Prevalence and Correlates of Prenatal Vitamin A Deficiency in Rural Sidama , Southern Ethiopia. *Journal of Health, Population and Nutrition*, 31(2), 185–194.
- Getnet, W., Aycheh, W., and Tessema, T. (2018). Determinants of Food Taboos in the Pregnant Women of the Awabel District , East Gojjam Zone , Amhara Regional State in Ethiopia. *Advances in Public Health*, 2018, 6. <https://doi.org/https://doi.org/10.1155/2018/9198076>
- Gewa, C. A., Murphy, S. P., Weiss, R. E., & Neumann, C. G. (2013). Determining minimum food intake amounts for diet diversity scores to maximize associations with nutrient adequacy: An analysis of schoolchildren’s diets in rural Kenya. *Public Health Nutrition*, 17(12), 2667–2673. <https://doi.org/10.1017/S1368980014000469>
- Gibson, R. S., Abebe, Y., Hambidge, K. M., Arbide, I., Teshome, A., & Stoecker, B. J. (2009). Inadequate feeding practices and impaired growth among children from subsistence farming households in Sidama, Southern Ethiopia. *Maternal and Child Nutrition*, 5, 260–275.

<https://doi.org/10.1111/j.1740-8709.2008.00179.x>

- Gladys, C. (2017). Dietary Assessment and Nutritional Status of Children (6-23 Months) Consuming Local Food Recipes in Vihiga County, Western Kenya. MSc Thesis. University of Nairobi, 2015(December).Pp, 137.
- Hadush, Z., Birhanu, Z., Chaka, M., & Gebreyesus, H. (2017). Foods tabooed for pregnant women in Abala district of Afar region, Ethiopia: an inductive qualitative study. *BMC Nutrition*, 3(1), 40. <https://doi.org/10.1186/s40795-017-0159-x>
- Haile, D., Azage, M., Mola, T., and Rainey, R. (2016). Exploring spatial variations and factors associated with childhood stunting in Ethiopia: Spatial and multilevel analysis. *BMC Pediatrics*, 16(1), 1–14. <https://doi.org/10.1186/s12887-016-0587-9>
- Harika, R., Faber, M., Samuel, F., Mulugeta, A., Kimiywe, J., & Eilander, A. (2017). Are Low Intakes and Deficiencies in Iron, Vitamin A, Zinc, and Iodine of Public Health Concern in Ethiopian, Kenyan, Nigerian, and South African Children and Adolescents? *Food and Nutrition Bulletin*, 38(3), 405–427. <https://doi.org/10.1177/0379572117715818>
- Hagos, S., Hailemariam, D., WoldeHanna, T., and Lindtjørn, B. (2017). Spatial heterogeneity and risk factors for stunting among children under age five in Ethiopia: A Bayesian geo-statistical model. *PLoS ONE*, 12(2). <https://doi.org/10.1371/journal.pone.0170785>
- Hailelassie, K., Mulugeta, A., & Girma, M. (2013). Feeding practices, nutritional status and associated factors of lactating women in Samre Woreda, South Eastern Zone of Tigray, Ethiopia. *Nutrition Journal*, 12(1), 1. <https://doi.org/10.1186/1475-2891-12-28>

- Hannibal, L., Lysne, V., Bjørke-Monsen, A.-L., Behringer, S., Grünert, S. C., Spiekerkoetter, U., ... Blom, H. J. (2016). Biomarkers and algorithms for the diagnosis of vitamin B12 deficiency. *Frontiers in Molecular Biosciences*, 3, 27.  
<https://doi.org/10.3389/fmolb.2016.00027>
- Hailu, C., Dejene, T., & Tebeje, B. (2015). Prevalence and Socio-Demographic/Economic Risk Factors of Low and High Body Mass Index of Urban versus Rural Pregnant Women at Booking for Antenatal Care in Governmental Health Centers of Jimma Zone, Southwest Ethiopia. *OALib*, 2(7), 1–17.
- Hamner, H. C., Perrine, C. G., Gupta, P. M., Herrick, K. A., & Cogswell, M. E. (2017). Food consumption patterns among U.S. children from birth to 23 months of age, 2009–2014. *Nutrients*, 9(9), 2009–2014. <https://doi.org/10.3390/nu9090942>
- Hasanain, F. G., Jamsiah, M., Zaleha, M. I., Tamil, A. M., & Mohammed, A. A. (2012). Association between drinking water sources and diarrhea with malnutrition among kindergarten's children in Baghdad city, Iraq. *Malaysian Journal of Public Health Medicine*, 12(1), 45–48.
- Herrador, Z., Sordo, L., Gadisa, E., Buño, A., Gómez-Rioja, R., Iturzaeta, J. M., ... Custodio, E. (2014). Micronutrient deficiencies and related factors in school-aged children in ethiopia: A cross-sectional study in Libo Kemkem and Fogera Districts, Amhara Regional State. *PLoS ONE*, 9(12), 1–20. <https://doi.org/10.1371/journal.pone.0112858>
- Herrador, Z., Sordo, L., Gadisa, E., Moreno, J., Nieto, J., Benito, A., ... Custodio, E. (2014). Cross-sectional study of malnutrition and associated factors among school aged children in

- rural and urban settings of fogera and libo kemkem districts, ethiopia. *PLoS ONE*, 9(9): e105880. doi:10.1371/journal.pone.0105880
- Herrmann, W., & Obeid, R. (2013). Utility and limitations of biochemical markers of vitamin B12 deficiency. *Eur J Clin Invest*, 43(3), 231–237. <https://doi.org/10.1111/eci.12034>
- Hirvonen, K., Taffesse, A. S., & Worku Hassen, I. (2016). Seasonality and household diets in Ethiopia. *Public Health Nutrition*, 19(10), 1723–1730. <https://doi.org/10.1017/S1368980015003237>
- Hirvonen, K., Hoddinott, J., Minten, B., & Stifel, D. (2017). Children’s Diets, Nutrition Knowledge, and Access to Markets. *World Development*, 95, 303–315. <https://doi.org/10.1016/j.worlddev.2017.02.031>
- Hm, H. T., Ahmed, E. M., and N, A. A. M. (2018). Food Taboos among Pregnant Women in Health Centers, Khartoum StateSudan, 2016. *International Journal of Science and Healthcare Research*, 3(1), 1-13.
- Hundera, T. D., Gemedo, H. F., Wirtu, D., & Kenie, D. N. (2015). Nutritional Status and Associated Factors among Lactating Mothers in Nekemte Referral Hospital and Health Centers, Ethiopia. *Food Science and Quality Management*, 35, 64–72. Retrieved from <http://iiste.org/Journals/index.php/FSQM/article/view/19192>
- Bazzano, A. N., Potts, K. S., & Mulugeta, A. (2018). How do pregnant and lactating women, and young children, experience religious food restriction at the community level? A qualitative study of fasting traditions and feeding behaviors in four regions of Ethiopia. *PLoS ONE*, 13(12). <https://doi.org/10.1371/journal.pone.0208408>

- IMF. (2011). *The Federal Democratic Republic of Ethiopia : Poverty Reduction Strategy Paper : Growth and transformation Plan 2010/11-2014/15-Volume I*. Washington D.C.
- Iqbal, S., Zakar, R., Zakar, M. Z., & Fischer, F. (2017). Factors associated with infants' and young children's (6-23 months) dietary diversity in Pakistan: Evidence from the demographic and health survey 2012-13. *Nutrition Journal*, 16(1).  
<https://doi.org/10.1186/s12937-017-0297-7>
- Jati, I. R. A. P., Widmer, C., Purwestri, R. C., Wirawan, N. N., Gola, U., Lambert, C., and Biesalski, H. K. (2014). Design and validation of a program to identify inadequate intake of iron, zinc, and vitamin A. *Nutrition*, 30(11–12), 1310–1317.  
<https://doi.org/10.1016/j.nut.2014.03.015>
- Jayatissa, R., Bekele, A., Piyasena, C. L., & Mahamithawa, S. (2006). Assessment of nutritional status of children under five years of age , pregnant women , and lactating women living in relief camps after the tsunami in Sri Lanka. *Food and Nutrition Bulletin*, 27(2), 144–152.
- Coates, J., Swindale, A., & Bilinsky, P. (2007). Household Food Insecurity Access Scale (HFIAS) for Measurement of Food Access: Indicator Guide. *Journal of Chemical Information and Modeling* (Vol. 53, pp. 1689–1699).  
<https://doi.org/10.1017/CBO9781107415324.004>
- Kandala, N., Fahrmeir, L., Klasen, S., and Priebe, J. (2009). Geo-additive Models of Childhood Undernutrition in Three Sub-Saharan African Countries 1. *Population, Space and Place*, 473(December 2008), 461–473.
- Kassa, T., Meshesha, B., Haji, Y., & Ebrahim, J. (2016). Appropriate complementary feeding



- practices and associated factors among mothers of children age 6-23 months in Southern Ethiopia, 2015. *BMC Pediatrics*, 16(1), 1–10. <https://doi.org/10.1186/s12887-016-0675-x>
- Kariuki, L. W., Lambert, C., Purwestri, R. C., Maundu, P., & Biesalski, H. K. (2017). Role of food taboos in energy, macro and micronutrient intake of pregnant women in western Kenya. *Nutrition and Food Science*, 47(6), 795–807. <https://doi.org/10.1108/NFS-09-2016-0146>
- Karl Knutsson, E., & Selinus, R. (1970). Fasting in Ethiopia. *The American Journal of Clinical Nutrition*, 23(7), 956–969.
- Khan, G. N., Ariff, S., Khan, U., Habib, A., Umer, M., Suhag, Z., ... Soofi, S. (2017). Determinants of infant and young child feeding practices by mothers in two rural districts of Sindh, Pakistan: A cross-sectional survey. *International Breastfeeding Journal*, 12(1), 1–8. <https://doi.org/10.1186/s13006-017-0131-z>
- Kiziltan, G., Karabudak, E., Tuncay, G., Avsar, F., Tuncay, P., Mungan, O., & Meral, P. (2005). Dietary intake and nutritional status of Turkish pregnant women during Ramadan. *Saudi Med J*, 26(11), 1782–1787.
- Koufakis, T., Karras, S., Antonopoulou, V., Angeloudi, E., Zebekakis, P., and Kotsa, K. (2017). Effects of Orthodox religious fasting on human health: a systematic review. *European Journal of Nutrition*, 56(8), 2439–2455. <https://doi.org/10.1007/s00394-017-1534-8>
- Kumera, G., Tsedal, E., & Ayana, M. (2018). Dietary diversity and associated factors among children of Orthodox Christian mothers/caregivers during the fasting season in Dejen District, North West Ethiopia. *Nutrition and Metabolism*. 15:16.

<https://doi.org/10.1186/s12986-018-0248-0>

- Kwok, T., Cheng, G., Lai, W. K., Poon, P., Woo, J., & Pang, C. P. (2004). Use of Fasting Urinary Methylmalonic Acid to Screen for Metabolic Vitamin B12 Deficiency in Older Persons. *Nutrition*, 20, 1–4. <https://doi.org/10.1016/j.nut.2004.06.001>
- Lambert, C., Eleraky, L., Mbwana, H., Kinabo, J., Biesalski, H. K., Riedel, S., ... Stuetz, W. (2018). Validation of a food group based nutrition software to assess nutrient intake in Tanzania. In *Global food security and food safety: The role of universities* (Vol. 30, p. 2018). Ghent, Belgium: Ghent University.
- Lander, R., Enkhjargal, T. S., Batjargal, J., Bolormaa, N., Enkhmyagmar, D., Tserendolgor, U., ... Gibson, R. S. (2010). Poor dietary quality of complementary foods is associated with multiple micronutrient deficiencies during early childhood in Mongolia. *Public Health Nutrition*, 13(9), 1304–1313. <https://doi.org/10.1017/S1368980009991856>
- Lazarou, C., and Matalas, A. (2010). A critical review of current evidence , perspectives and research implications of diet-related traditions of the Eastern Christian Orthodox Church on dietary intakes and health consequences. *International Journal of Food Sciences and Nutrition*, 61:7, 739-758. DOI: 10.3109/09637481003769782.
- Lebso, M., Anato, A., & Loha, E. (2017). Prevalence of anemia and associated factors among pregnant women in Southern Ethiopia: A community based cross-sectional study. *PLoS ONE*, 12(12), 1–11. <https://doi.org/10.1371/journal.pone.0188783>
- Leiper, J., Molla, A. M., and Molla, A. M. (2003). Effects on health of fluid restriction during fasting in Ramadan. *European Journal of Clinical Nutrition*, 57(2), S30–S38.

<https://doi.org/10.1038/sj.ejcn.1601899>

Ludwig, C., & Bau, A. (2007). *Nutrition Baseline Survey - Ethiopia -for the Global Programme: Food and Nutrition Security and Enhanced Resilience*. Addis Ababa, Ethiopia.

Lyu, L., Hsu, Y., Chen, H., Lo, C., & Lin, C. (2014). Comparisons of four dietary assessment methods during pregnancy in Taiwanese women. *Taiwanese Journal of Obstetrics & Gynecology*, 53(2), 162–169. <https://doi.org/10.1016/j.tjog.2014.04.007>

Maliwichi-nyirenda, C. P., Maliwichi, L. L., Maliwichi-nyirenda, C. P., and Maliwichi, L. L. (2017). Studies on Ethno-Medicine Food Related Taboos Observed During Pregnancy in Malawi Food Related Taboos Observed During Pregnancy in Malawi. *Studies on Ethno-Medicine*, 10(2), 263–268. <https://doi.org/10.1080/09735070.2016.11905496>

Makoka, D., & Masibo, P. K. (2015). Is there a threshold level of maternal education sufficient to reduce child undernutrition? Evidence from Malawi, Tanzania and Zimbabwe. *BMC Pediatrics*, 15, 1–10. <https://doi.org/10.1186/s12887-015-0406-8>

Marie-Claude; Kennedy, Gina; Lee, Warren T. K.; and Moursi, Mourad. 2015. Moving forward on choosing a standard operational indicator of women's dietary diversity. Rome, Italy: Food and Agricultural Organization of the United Nations (FAO). <http://www.fao.org/3/a-i4942e.pdf>

Masreha Tessema, Tefera Belachew, & Getahun Ersino. (2013). Feeding patterns and stunting during early childhood in rural communities of Sidama, south Ethiopia. *Pan African Medical Journal*, 14(75), 1–12. <https://doi.org/10.11604/pamj.2013.14.75.1630>

McGovern, M. E., Krishna, A., Aguayo, V. M., and Subramanian, S. V. (2017). A review of the

- evidence linking child stunting to economic outcomes. *International Journal of Epidemiology*, 46(4). <https://doi.org/10.1093/ije/dyx017>
- Meyer-Rochow, V. B. (2009). Food taboos: Their origins and purposes. *Journal of Ethnobiology and Ethnomedicine*, 5, 1–10. <https://doi.org/10.1186/1746-4269-5-18>
- Mekonnen, T. C., Workie, S. B., Yimer, T. M., & Mersha, W. F. (2017). Meal frequency and dietary diversity feeding practices among children 6–23 months of age in Wolaita Sodo town, Southern Ethiopia. *Journal of Health, Population and Nutrition*, 36(1), 18. <https://doi.org/10.1186/s41043-017-0097-x>
- Mekuria, G., Wubneh, Y., & Tewabe, T. (2017). Household dietary diversity and associated factors among residents of finote selam town , north west Ethiopia : a cross sectional study. *BMC Nutrition*, 3(28), 3–8. <https://doi.org/10.1186/s40795-017-0148-0>
- Mengistu, G., Moges, T., Samuel, A., & Baye, K. (2017). Energy and nutrient intake of infants and young children in pastoralist communities of Ethiopia. *Nutrition*, 41, 1–6. <https://doi.org/10.1016/j.nut.2017.02.012>
- Mesfin, A., Henry, C., Girma, M., & Whiting, S. J. (2016). Use of pulse crops in complementary feeding of 6-23-month-old infants and young children in Taba Kebele, Damot Gale District, Southern Ethiopia. *Journal of Public Health in Africa*, 6(1), 1–6. <https://doi.org/10.4081/jphia.2015.357>
- Meyer-Rochow, V. B. (2009). Food taboos: Their origins and purposes. *Journal of Ethnobiology and Ethnomedicine*, 5, 1–10. <https://doi.org/10.1186/1746-4269-5-18>

- Morseth, M. S., Torheim, L. E., Gebremariam, M. K., Chandyo, R. K., Ulak, M., Shrestha, S. K., ... Henjum, S. (2018). Tracking of infant and young child feeding practices among 9- to 24-month-old children in Nepal: The MAL-ED Birth Cohort Study. *Public Health Nutrition*, 21(2), 355–364. <https://doi.org/10.1017/S1368980017002294>
- Mosites, E. M., Rabinowitz, P. M., Thumbi, S. M., & Montgomery, J. M. (2015). The Relationship between Livestock Ownership and Child Stunting in Three Countries in Eastern Africa Using National Survey Data. *PLoS ONE*, 10(9), 1–10. <https://doi.org/10.1371/journal.pone.0136686>
- Müller, O., & Krawinkel, M. (2005). Malnutrition And Health In Developing Countries. *Canadian Medical Association Journal*, 173(3), 279–286. <https://doi.org/10.1503/cmaj.050342>
- Mulat, E., Alem, G., Woyraw, W., & Temesgen, H. (2019). Uptake of minimum acceptable diet among children aged 6 – 23 months in orthodox religion followers during fasting season in rural area , Dembecha, north West Ethiopia. *BMC Nutrition*, 5(18), 1–10.
- Mulugeta, A., Hagos, F., Kruseman, G., Linderhof, B., Stoecker, B., Abraha, Z., ... Samual, G. G. (2010). Factors Contributing to Child Malnutrition in Tigray , Northern Ethiopia. *East African Medical Journal*, 87(6), 248–254. <https://doi.org/10.4314/eamj.v87i6.63083>
- National Academy of Sciences. (1991). *Nutrition During Lactation* (4th Editio). Washington, D.C: National Academy Press.
- Naik, R. and R. Smith. 2015. Linkages Between Family Planning and Nutrition. Washington, DC: Futures Group, Health Policy Project

- Negash, C., Whiting, S. J., Henry, C. J., Belachew, T., & Hailemariam, T. G. (2015). Association between maternal and child nutritional status in Hula , rural southern Ethiopia : A cross sectional study. *PLoS ONE*, *10*(11), e0142301. <https://doi.org/10.1371/journal.pone.0142301>
- Neumann, C., Harris, D. M., & Rogers, L. M. (2002). Contribution of animal source foods in improving diet quality and function in children in the developing world. *Nutrition Research*, *22*, 193–220.
- Nguyen, P. H., Avula, R., Ruel, M. T., Saha, K. K., Ali, D., Tran, L. M., ... Rawat, R. (2013). Maternal and Child Dietary Diversity Are Associated in Bangladesh, Vietnam, and Ethiopia. *The Journal of Nutrition*, *143*, 1176–1183. <https://doi.org/10.3945/jn.112.172247>.
- Nigatu, M., Gebrehiwot, T. T., & Gemed, D. H. (2018). Household Food Insecurity, Low Dietary Diversity, and Early Marriage Were Predictors for Undernutrition among Pregnant Women Residing in Gambella, Ethiopia. *Advances in Public Health*, *2018*(1350195), 10. <https://doi.org/https://doi.org/10.1155/2018/1350195>
- Nkurunziza, S., Meessen, B., Van geertruyden, J. P., & Korachais, C. (2017). Determinants of stunting and severe stunting among Burundian children aged 6-23 months: Evidence from a national cross-sectional household survey, 2014. *BMC Pediatrics*. <https://doi.org/10.1186/s12887-017-0929-2>
- Nyaruhucha, C. N. M., Msuya, J. M., Mamiro, P. S., & Kerengi, A. J. (2006). Nutritional status and feeding practices of under-five children in Simanjiro District, Tanzania. *Tanzania Health Research Bulletin*, *8*(3), 162–167.
- Ocampo-Guirindola, M. L., Garcia-Malabad, C. J., Valdeabell-Maniego, M. L. M., & Punzalan,

- S. L. M. (2016). Association between dietary diversity score and nutritional status of Filipino children aged 6-23 months. *Philippine Journal of Science*, 145(1), 57–69.
- Ogechi, U. P., & Chilezie, O. V. (2017). Assessment of Dietary Diversity Score, Nutritional Status and Socio-demographic Characteristics of Under-5 Children in Some Rural Areas of Imo State, Nigeria. *Mal J Nutr*, 23(3), 425–435.
- Özaltın, E., Hill, K., & Subramanian, S. V. (2011). Association of Maternal Stature With Offspring Mortality, Underweight, and Stunting in Low- to Middle-Income Countries. *JAMA*, 303(15), 1507–1516. <https://doi.org/10.1001/jama.2010.450>.
- Papadaki, A., Vardavas, C., Hatzis, C., and Kafatos, A. (2007). Calcium, nutrient and food intake of Greek Orthodox Christian monks during a fasting and non-fasting week. *Public Health Nutrition*, 11(10), 1022–1029. <https://doi.org/10.1017/S1368980007001498>
- Picciano, M. F. (2003). Dietary Supplement Use in Women : Current Status and Future Directions Pregnancy and Lactation : Physiological Adjustments , Nutritional Requirements and the Role of Dietary Supplements. *J. Nutr.*, 133, 1997–2002.
- Rakicioğlu, N., Samur, G., Topçu, A., & Topçu, A. A. (2006). The effect of Ramadan on maternal nutrition and composition of breast milk. *Pediatrics International*, 48(3), 278–283. <https://doi.org/10.1111/j.1442-200X.2006.02204.x>
- Regassa, N., & Stoecker, B. J. (2012). Household food insecurity and hunger among households in Sidama district, southern Ethiopia. *Public Health Nutrition*, 15(7), 1276–1283. <https://doi.org/10.1017/S1368980011003119>
- Riang’a, R. M., Broerse, J., and Nangulu, A. K. (2017). Food beliefs and practices among the

- Kalenjin pregnant women in rural Uasin Gishu County, Kenya. *Journal of Ethnobiology and Ethnomedicine*, 13(1). <https://doi.org/10.1186/s13002-017-0157-8>
- Roba, K. T., O'Connor, T. P., Belachew, T., and O'Brien, N. M. (2015). Seasonal variation in nutritional status and anemia among lactating mothers in two agro-ecological zones of rural Ethiopia: A longitudinal study. *Nutrition*, 31(10), 1213–1218. <https://doi.org/10.1016/j.nut.2015.03.007>
- Roba, K. T., O'Connor, T. P., Belachew, T., and O'Brien, N. M. (2016a). Infant and Young Child Feeding (IYCF) Practices Among Mothers of Children Aged 6–23 Months in Two Agro-ecological Zones of Rural Ethiopia. *International Journal of Nutrition and Food Sciences*, 5(3), 185. <https://doi.org/10.11648/j.ijnfs.20160503.16>
- Roba, K. T., O'Connor, T. P., Belachew, T., and O'Brien, N. M. (2016b). Variations between post- and pre-harvest seasons in stunting, wasting, and infant and young child feeding (IYCF) practices among children 6-23 months of age in lowland and midland agro-ecological zones of rural Ethiopia. *Pan African Medical Journal*, 24, 1–9. <https://doi.org/10.11604/pamj.2016.24.163.9387>
- Sadeghirad, B., Motaghipisheh, S., Kolahdooz, F., Zahedi, M. J., and Haghdoost, A. A. (2014). Islamic fasting and weight loss: A systematic review and meta-analysis. *Public Health Nutrition*, 17(2), 396–406. <https://doi.org/10.1017/S1368980012005046>
- Sarri, K. O., Tzanakis, N. E., Linardakis, M. K., Mamalakis, G. D., & Kafatos, A. G. (2003). Effects of Greek orthodox christian church fasting on serum lipids and obesity. *BMC Public Health*, 8, 1–8.



- Sarri, K. O., Linardakis, M. K., Bervanaki, F. N., Tzanakis, N. E., & Kafatos, A. G. (2004). Greek Orthodox fasting rituals: a hidden characteristic of the Mediterranean diet of Crete. *British Journal of Nutrition*, 92, 277–284. <https://doi.org/10.1079/BJN20041197>
- Sarri, K. O., Kafatos, A. G., and Higgins, S. (2005). Is religious fasting related to iron status in Greek Orthodox Christians? *British Journal of Nutrition*, 2001, 198–203. <https://doi.org/10.1079/BJN20051472>
- Schlemmer, U., Frølich, W., Prieto, R. M., & Grases, F. (2009). Phytate in foods and significance for humans: food sources, intake, processing, bioavailability, protective role and analysis. *Mol Nutr Food Re*, 53(Suppl 2), S330-75. <https://doi.org/doi:10.1002/mnfr.200900099>.
- Schwei, R. J., Tesfay, H., Asfaw, F., Jogo, W., & Busse, H. (2017). Household dietary diversity, vitamin A consumption and food security in rural Tigray, Ethiopia. *Public Health Nutrition*, 20(9), 1540–1547. <https://doi.org/10.1017/S1368980017000350>
- Sellassie, S. H., and Mikael, B. (2019). Worship in the Ethiopian Orthodox church. Retrieved March 12, 2019, from <https://www.ethiopianorthodox.org/english/ethiopian/worship.html>
- Senarath, U., Godakandage, S. S. P., Jayawickrama, H., Siriwardena, I., & Dibley, M. J. (2012). Determinants of inappropriate complementary feeding practices in young children in Sri Lanka: Secondary data analysis of Demographic and Health Survey 2006-2007. *Maternal and Child Nutrition*, 8(SUPPL. 1), 60–77. <https://doi.org/10.1111/j.1740-8709.2011.00375.x>

- Silva, D. e, Valente, A., Borges, A., Almeida, F., Neves, E., & Afonso, C. (2017). Relationship between the mothers' nutritional status with that of a child population from São Tomé Príncipe, “ *Africa. Rev. Bras. Saúde Matern. Infant., Recife*, 17(2), 327–335.  
<https://doi.org/http://dx.doi.org/10.1590/1806-93042017000200007>
- Sitotaw, I. K., Kiday, H., & Adama, Y. (2017). Comparison of nutritional status and associated factors of lactating women between lowland and highland communities of District Raya, Alamata. *BMC Nutrition*, 3(61), 1–12. <https://doi.org/10.1186/s40795-017-0179-6>
- Sohnesen, T. P., Ambel, A. A., Fisker, P., Andrews, C., and Khan, Q. (2017). Small area estimation of child undernutrition in Ethiopian districts. *PLoS ONE*, 12(4), 1–17.  
<https://doi.org/https://doi.org/10.1371/journal.pone.0175445>
- Solomon, D., Aderaw, Z., & Tegegne, T. K. (2017). Minimum dietary diversity and associated factors among children aged 6-23 months in Addis Ababa, Ethiopia. *International Journal for Equity in Health*, 16(1), 1–9. <https://doi.org/10.1186/s12939-017-0680->
- Steyn, N. P., Nel, J. H., Nantel, G., Kennedy, G., & Labadarios, D. (2006). Food variety and dietary diversity scores in children : are they good indicators of dietary adequacy? *Public Health Nutrition*, 9(5), 644–650. <https://doi.org/10.1079/PHN2005912>.
- Stoecker, B. J., and Abebe, Y. (2014). Context for food-based approaches for improved feeding of infants and children in Ethiopia. *Food and Nutrition Bulletin*, 35(4), S209–S213.  
<https://doi.org/10.1177/15648265140354S308>.
- Tafa, M., Haidar, J., & Fekadu, H. (2015). Gynecology & Obstetrics Modern Family Planning Use and its Relationship with the Nutritional Status of Women of Child Bearing Age at

- Tena District , Arsi Zone Oromia Regional State , Ethiopia 2013 : A Community based Case Control Study. *Gynecology & Obstetrics*, 5(3), 1–10. <https://doi.org/10.4172/2161-0932.1000282>
- Tamiru, M. W., Tolessa, B. E., & Abera, S. F. (2015). Under Nutrition and Associated Factors Among Under-Five Age Children of Kunama Ethnic Groups in Tahtay Adiyabo Woreda , Tigray Regional State , Ethiopia : Community Based Study. *International Journal of Nutrition and Food Sciences*, 4(3), 277–288. <https://doi.org/10.11648/j.ijnfs.20150403.15>
- Tebekaw, Y., Teller, C., & ColÃ³n-Ramos, U. (2014). The burden of underweight and overweight among women in Addis Ababa, Ethiopia. *BMC Public Health*, 14(1126). <https://doi.org/10.1186/1471-2458-14-1126>
- Tassew, A. A., Tekle, D. Y., Belachew, A. B., & Adhena, B. M. (2019). Factors affecting feeding 6–23 months age children according to minimum acceptable diet in Ethiopia: A multilevel analysis of the Ethiopian Demographic Health Survey. *PLoS ONE*, 14(2). <https://doi.org/10.1371/journal.pone.0203098>
- Tegegne, M., Sileshi, S., Benti, T., Teshome, M., & Woldie, H. (2017). Factors associated with minimal meal frequency and dietary diversity practices among infants and young children in the predominantly agrarian society of Bale zone, Southeast Ethiopia: A community based cross sectional study. *Archives of Public Health*, 75(1), 1–11. <https://doi.org/10.1186/s13690-017-0216-6>
- Tesfaye, B., Sinclair, K., Wuehler, S. E., Moges, T., De-regil, L. M., & Dickin, K. L. (2018). Applying international guidelines for calcium supplementation to prevent pre-eclampsia : simulation of recommended dosages suggests risk of excess intake in Ethiopia. *Public*

- Health Nutrition*, 22(3), 531–541. <https://doi.org/10.1017/S1368980018002562>
- Tessema, M., De Groote, H., Brouwer, I. D., Feskens, E. J. M., Belachew, T., Zerfu, D., ...  
 Gunaratna, N. S. (2019). Soil zinc is associated with serum zinc but not with linear growth of children in Ethiopia. *Nutrients*, 11(221). <https://doi.org/10.3390/nu11020221>
- Tewodros, D. (2008). *Beef Cattle Production System and Opportunities for Market Orientation in Borena Zone, Southern*. MSc Thesis, Department of Animal Sciences. Haromaya University.Pp.139.
- Tezera, R., Sahile, Z., Yilma, D., Misganaw, E., & Mulu, E. (2018). Prevalence of anemia among school-age children in Ethiopia: A systematic review and meta-analysis. *Systematic Reviews*, 7(1). <https://doi.org/10.1186/s13643-018-0741-6>
- Tigga, P. L., & Sen, J. (2016). Maternal Body Mass Index Is Strongly Associated with Children Z -Scores for Height and BMI. *Journal of Anthropology*, 2016(6538235), 10.  
<https://doi.org/10.1155/2016/6538235>
- Tigray Region Agriculture and Rural Development Bureau. (2016). Food Security Status of Tigray Districts 2016.pdf.
- Tolulope, A., and Olubukola, B. (2015). Spatial Pattern of Household Food Insecurity and Childhood Malnutrition in Akinyele Local Government Area, Nigeria, (January).  
<https://doi.org/10.15640/ijhs.v3n1a15>
- Trepanowski, J. F., and Bloomer, R. J. (2010). The impact of religious fasting on human health. *Nutrition Journal*, 9(1), 57. <https://doi.org/10.1186/1475-2891-9-57>

- UNICEF. (1998). *The status of the world's children*. Geneva, Switzerland: Oxford University Press.
- UNICEF. (2015). *UNICEF's approach to scaling up nutrition for mothers and their children. Discussion paper. Programme Division, UNICEF, New York, June 2015.*
- UNICEF. (2017). *Leveles and trend in child malnutrition: UNICEF/WHO/World Bank Group joint estimates , key findings of the 2017 edition.*
- Vasilevski, V., Alban, S., Box, P. O., Carolan-olah, M., & Affiliation, M. (2016). Ethiopian pregnancy food taboos. *Journal of Clinical Nursing*, 25(19), 3069–3075. <https://doi.org/https://doi.org/10.1111/jocn.13319>
- Vollmer, S., Bommer, C., Krishna, A., Harttgen, K., & Subramanian, S. V. (2017). The association of parental education with childhood undernutrition in low- and middle-income countries : comparing the role of paternal and maternal education. *International Journal of Epidemiology*, 46(August 2016), 312–323. <https://doi.org/10.1093/ije/dyw133>
- Weldehaweria, N. B., Misgina, K. H., Weldu, M. G., Gebregiorgis, Y. S., Gebrezgi, B. H., Zewdie, S. W., ... Alemu, W. (2016). Dietary diversity and related factors among lactating women visiting public health facilities in Aksum town, Tigray, Northern Ethiopia. *BMC Nutrition*, 2(38). <https://doi.org/10.1186/s40795-016-0077-3>
- WFP. (2009). *Food Security and Vulnerability in Selected Towns of Tigray Region , Ethiopia.*
- WHO. (1998). Complementary feeding of young children in developing countries: a review of current scientific knowledge. Geneva, Switzerland.
- WHO. (2010). Indicators for assessing infant and young child feeding practices. *Part 3 Country*

- Profiles*, 1–47. [https://doi.org/ISBN 978 92 4 159975 7](https://doi.org/ISBN%20978%2092%204%20159975%207)
- WHO. (2014). Postnatal care of the mother and newborn 2013. *World Health Organization*, 1–72. [https://doi.org/978 92 4 150664 9](https://doi.org/978%2092%204%20150664%209)
- WHO. (2011). The Global Prevalence of Anaemia in 2011. *Document, World Heal*, 1–43.
- Williams, J., Mai, C. T., and Mulinare, J. (2015). National Birth Defects Prevention Month and Folic Acid Awareness Week — January 2015 Updated Estimates of Neural Tube Defects Prevented by Mandatory Folic Acid Fortification — United States , 1995 – 2011. *MMWR*, 64(1), 1–5.
- Workicho, A., Belachew, T., Feyissa, G. T., Wondafrash, B., Lachat, C., Verstraeten, R., & Kolsteren, P. (2016). Household dietary diversity and Animal Source Food consumption in Ethiopia: Evidence from the 2011 Welfare Monitoring Survey. *BMC Public Health*, 16(1192). <https://doi.org/10.1186/s12889-016-3861-8>
- Wu, X., & Li, L. (2012). Family size and maternal health: Evidence from the One-Child policy in China. *Journal of Population Economics*, 25(4), 1341–1364. <https://doi.org/10.1007/s00148-011-0361-0>
- Zerfu, T. A., Umeta, M., & Baye, K. (2016). Dietary habits, food taboos, and perceptions towards weight gain during pregnancy in Arsi, rural central Ethiopia: a qualitative cross-sectional study. *Journal of Health, Population, and Nutrition*, 35(1), 22. <https://doi.org/10.1186/s41043-016-0059-8>
- Zhang, J., Kang, H., Wang, L., Zhao, X., Liu, R., & He, L. (2016). A Prevalence Analysis to

Assess the Relationship between Maternal and Children' s Nutritional Status in North-Western Rural China Clinics in Mother and Child Health. *Clinics in Mother and Child Health*, 13(2), 100242. <https://doi.org/10.4172/2090-7214.1000242>

Zhao, W., Yu, K., Tan, S., Zheng, Y., Zhao, A., Wang, P., & Zhang, Y. (2017). Dietary diversity scores : an indicator of micronutrient inadequacy instead of obesity for Chinese children. *BMC Public Health*, 17(440). <https://doi.org/10.1186/s12889-017-4381-x>

## **Affidavit**

Annex 2 to the University of Hohenheim doctoral degree regulations for Dr. rer. nat.

Affidavit according to Sec. 7(7) of the University of Hohenheim doctoral degree regulations for Dr. rer. nat.

1. For the dissertation submitted on the topic

Dietary Intake, Nutritional Status of Lactating Women and their 6-23-Months-Old Children in Genta Afeshum District, Rural Ethiopia; Development and Validation of Calculator for Inadequate Micronutrient Intake (CIMI)

I hereby declare that I independently completed the work.

2. I only used the sources and aids documented and only made use of permissible assistance by third parties. In particular, I properly documented any contents which I used- either by directly quoting or paraphrasing - from other works.

3. I did not accept any assistance from a commercial doctoral agency or consulting firm.

4. I am aware of the meaning of this affidavit and the criminal penalties of an incorrect or incomplete affidavit.

I hereby confirm the correctness of the above declaration: I hereby affirm in lieu of oath that I have, to the best of my knowledge, declared nothing but the truth and have not omitted any information.

Place and Date

Signature



# **Curriculum Vitae**

## **Personal Information**

Full name: Beruk Berhanu Desalegn

Sex: Male

Nationality: Ethiopian

Date of Birth: July 26, 1985G.C

Place of Birth: Hawassa, Ethiopia

## **Contact Address**

School of Nutrition, Food Science and Technology

Hawassa University, Hawassa, SNNPR, Ethiopia

P.O. Box: 05

Phone Number: +251-941048918 or +251-916041191

Email: BerukB.Desalegn @uni-hohenheim.de

berhanuberuk@gmail.com

## **Educational Information**

- PhD student at University of Hohenheim, Germany (Since April, 2016)
- MSc student in Food Science and Post-Harvest Technology (2010-2013)
  - MSc Thesis Title: Formulation of Quality Protein Maize Based Nutritionally Improved Complementary Food: The Case of Shebedino Woreda, Southern Ethiopia
  - Thesis Result: Excellent
- BSc student in Rural Development and Family Sciences (2007-2010)

- BSc Thesis Title: Knowledge, Attitude and Practices of Family Planning Use in Hawella-Tulla Sub-city, Hawassa Town
- Diploma in Medical Laboratory Technology (2003-2004)

### **Work Experience**

- Lecturer and researcher (Since October, 2013)  
School of Nutrition, Food Science and Technology, Hawassa University
- Head, Gender Office, Food preparation and Dormitory Services
  - College of Agriculture, Hawassa University (2009-2013)
- Data Analyst
  - College of Agriculture, Hawassa University (2009)
- Medical Laboratory Technician
  - Ministry of Health of Ethiopia and Hawassa University (2005-2009)

### **Publications**

1. Desalegn, B.; Lambert, C. Low diet diversity and its associated factors among the mothers and their children in agroforestry land use systems of Sidama, Ethiopia: A community-based cross-sectional study. *Cogent Food and Agriculture* 2020, 6(1), 1818367. <https://doi.org/10.1080/23311932.2020.1818367>
2. Desalegn, B.; Lambert, C.; Riedel, S.; Negese, T.; Biesalski, H.K. Feeding Practices and Undernutrition in 6–23-Month-Old Children of Orthodox Christian Mothers in Rural Tigray, Ethiopia: Longitudinal Study. *Nutrients* 2019, 11 (1), 138. doi:10.3390/nu11010138
3. Desalegn, B.; Lambert, C.; Riedel, S.; Negese, T.; Biesalski, H.K. Ethiopian Orthodox Fasting and Lactating Mothers: Longitudinal Study on Dietary Pattern and Nutritional Status in Rural Tigray, Ethiopia. *International Journal of Environmental Research and Public Health* 2018, 15 (8), 1767. doi:10.3390/ijerph15081767.
4. Desalegn, B.B.; Abegaz, K.; Kinf, E. Assessment of Knowledge and Practices on Complementary Food Preparation and Child Feeding at Shebedino, Sidama, Southern

- Ethiopia. *International Journal of Food Science and Nutritional Engineering* 2015, 5(2), 82-87.
5. Desalegn, B.B.; Abegaz, K.; Kifle, E. Effect of Blending Ratio and Processing Technique on Physicochemical, Functional Properties and Sensory Acceptability of Quality Protein Maize (QPM) Based Complementary Food. *International Journal of Food Science and Nutritional Engineering* 2015, 5 (3), 121-129.
  6. Berhanu, B. Effect of Soaking and Germination on Proximate Composition, Mineral Bioavailability and Functional Properties of Chickpea Flour. *Food and Public Health* 2015, 5(4),108-113.
  7. Desalegn, B.B.; Tadesse, F.T.; Haile, D. Physical and Proximate Characterization of Anchote (*Coccinia abyssinica*) Accessions Grown Under Hawassa and Wondo Genet Conditions, Southern Ethiopia. *Food Science and Quality Management* 2015, 42, 62-74.
  8. Desalegn, B.B.; Abegaz, K.; Kifle, E. Formulation of Quality Protein Maize (QPM) Based Nutritionally Improved Complementary Food: The Case of Shebedino Woreda, Southern Ethiopia. *European Journal of Nutrition and Food Safety* 2015, 5(5), 5-416.
  9. Anchamo, A.A.; Pragma, S.; Bosha, B.; Desalegn, B.B. Effect of Breakfast Eating Patterns and Anthropometric Measurements on Cognitive Function of Early Adolescents in Rural Area of Sidama Zone, Southern Ethiopia. *Journal of Food and Nutrition Sciences* 2015, 3(6), 251-258. doi: 10.11648/j.jfns.20150306.19
  10. Dagem, A.; Haki, G.; Ali, K.; Desalegn, B.B.; Basore., D. Proximate, Mineral Composition and Sensory Acceptability of Home Made Noodles from Nettle (*Urtica simensis*) Leaves and Wheat Flour Blends. *International Journal of Food Science and Nutritional Engineering* 2016, 6(3), 55-61. doi:10.5923/j.food.20160603.02
  11. Desalegn, B.B.; Kifle, E.; Fikre, K.; Bosha, T. Stunting and Its Associated Factors in Under Five Years Old Children: The Case of Hawassa University Technology Villages, Southern Ethiopia. *IOSR Journal of Environmental Science, Toxicology and Food Technology (IOSR-JESTFT)* 2016, 10(11), 25-31. doi: 10.9790/2402-1011022531
  12. Desalegn, B.B.; Abegaz, K.; Kifle, E. Formulation of Quality Protein Maize (QPM) Based Nutritionally Improved Complementary Food: The Case of Shebedino Woreda,

- Southern Ethiopia. Proceedings of the 5th Annual Conference, Food and Nutrition Society of Ethiopia 2015, p. 55-68.
13. Desalegn, B.B.; Tadesse, F.T.; Haile, D. Physical and Proximate Characterization and Mineral Analysis of Anchote (*Coccinia abyssinica*) Accessions Grown Under Hawassa and Wondo Genet Conditions, Southern Ethiopia. Proceedings of the 37th Annual Research Review Workshop, Hawassa University, Hawassa, Ethiopia. 2016, p. 1-18.
  14. Desalegn, B.B.; Kinfu, E.; Fikre, K.; Bosha, T. Stunting and Its Associated Factors in Under Five Years Old Children: The Case of Hawassa University Technology Villages, Southern Ethiopia. Proceedings of the 37th Annual Research Review Workshop, Hawassa University, Hawassa, Ethiopia. 2016, p.45-58.
  15. Getahun, D.; Berhanu, B. Formulation of Flat Bread (*Kitta*) from Maize (*Zea Mays Linnaeus.*) and Amaranth (*Amaranthus Caudatus L.*): Evaluation of Physico-Chemical Properties, Nutritional, Sensory and Keeping Quality. International Journal of Food Science and Nutritional Engineering 2017, 7(6), 125-131. DOI: 10.5923/j.food.20170706.01
  16. Hassen, Y.; Mukisa, I.M.; Kurabachew, H.; Desalegn, B.B. Evaluation of Yeast and Lactic Acid Bacteria Starter Cultures for the Production of Rice Injera. Evaluation of Yeast and Lactic Acid Bacteria Starter Cultures for the Production of Rice Injera. J. Food Process. Technol. 2017, 9,721. doi: 10.4172/2157-7110.1000721
  17. Desalegn, B.B.; Desta, F.T. Effect of Blending Ratio on Proximate Composition, Physico-Chemical Property, and Sensory Acceptability of Injera Produced from Red Tef (*Eragrostis tef*) and Cassava (*Manihot Esculenta*). Food Science and Quality Management 2017, 68,6-10.
  18. Berhanu, B.; Reta, F.; Getahun, D.; Zegeye, M. Formulation of Flat Bread (*Kitta*) from Maize (*Zea Mays Linnaeus.*) and Amaranth (*Amaranthus Caudatus L.*): Evaluation of Physico-Chemical Properties, Nutritional, Sensory and Keeping Quality. Proceedings of the 38th Annual Research Review Workshop, College of Agriculture, Hawassa University, Hawassa, Ethiopia. 2017, p. 179-187.

19. Desalegn, B.B.; Abegaz, K.; Kinf, E. Assessment of Knowledge and Practices on Complementary Food Preparation and Child Feeding at Shebedino, Sidama, Southern Ethiopia. Book of Abstracts of Research Articles Published by Academic Staffs of Hawassa University. Volume 2. December, 2017. Hawassa University, Hawassa, Ethiopia.
20. Desalegn, B.B.; Abegaz, K.; Kinf, E. Effect of Blending Ratio and Processing Technique on Physicochemical, Functional Properties and Sensory Acceptability of Quality Protein Maize (QPM) Based Complementary Food. Book of Abstracts of Research Articles Published by Academic Staffs of Hawassa University. Volume 2. December, 2017. Hawassa University, Hawassa, Ethiopia.

#### **Scholarship and Research Grant Award**

- PhD research grant from German Federal Ministry for Economic Cooperation and Development (BMZ) through the Food Security Center/DAAD Scholarship Program
- PhD Scholarship Award from University of Hohenheim
- Training Scholarship Award from CDI, Wageningen University and Research, Wageningen, The Netherlands.
- Large Scale Thematic Research grant from Hawassa University (1 project)
- Medium and Small Scale Research grants from Hawassa University (8 projects)
- MSc research grant from NORHED and USAID/ENGINE project at Hawassa University